



# Full wwPDB X-ray Structure Validation Report ⓘ

May 23, 2020 – 11:45 am BST

PDB ID : 1VJ1  
Title : Crystal structure of putative NADPH-dependent oxidoreductase from *Mus musculus* at 2.10 Å resolution  
Authors : Joint Center for Structural Genomics (JCSG)  
Deposited on : 2003-12-03  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	363	<div> <div></div> <div>81%</div> <div>13%</div> <div>6%</div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2720 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called putative NADPH-dependent oxidoreductase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	341	Total	C	N	O	S	Se	0	2	0
			2567	1620	445	485	8	9			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-10	GLY	-	LEADER SEQUENCE	UNP Q8VDQ1
A	-9	SER	-	LEADER SEQUENCE	UNP Q8VDQ1
A	-8	ASP	-	LEADER SEQUENCE	UNP Q8VDQ1
A	-7	LYS	-	LEADER SEQUENCE	UNP Q8VDQ1
A	-6	ILE	-	LEADER SEQUENCE	UNP Q8VDQ1
A	-5	HIS	-	LEADER SEQUENCE	UNP Q8VDQ1
A	-4	HIS	-	LEADER SEQUENCE	UNP Q8VDQ1
A	-3	HIS	-	LEADER SEQUENCE	UNP Q8VDQ1
A	-2	HIS	-	LEADER SEQUENCE	UNP Q8VDQ1
A	-1	HIS	-	LEADER SEQUENCE	UNP Q8VDQ1
A	0	HIS	-	LEADER SEQUENCE	UNP Q8VDQ1
A	1	MSE	MET	MODIFIED RESIDUE	UNP Q8VDQ1
A	52	MSE	MET	MODIFIED RESIDUE	UNP Q8VDQ1
A	56	MSE	MET	MODIFIED RESIDUE	UNP Q8VDQ1
A	135	MSE	MET	MODIFIED RESIDUE	UNP Q8VDQ1
A	158	MSE	MET	MODIFIED RESIDUE	UNP Q8VDQ1
A	244	MSE	MET	MODIFIED RESIDUE	UNP Q8VDQ1
A	325	MSE	MET	MODIFIED RESIDUE	UNP Q8VDQ1
A	332	MSE	MET	MODIFIED RESIDUE	UNP Q8VDQ1
A	333	MSE	MET	MODIFIED RESIDUE	UNP Q8VDQ1

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	153	Total	O	0	0
			153	153		



- Molecule 1: putative NADPH-dependent oxidoreductase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	42.39 Å 91.81 Å 100.56 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.28 – 2.10 50.28 – 2.10	Depositor EDS
% Data completeness (in resolution range)	94.9 (50.28-2.10) 94.9 (50.28-2.10)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.82 (at 2.10 Å)	Xtriage
Refinement program	REFMAC 5.1.9999	Depositor
R, $R_{free}$	0.180 , 0.216 0.188 , 0.228	Depositor DCC
$R_{free}$ test set	1123 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	31.6	Xtriage
Anisotropy	0.427	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 49.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	2720	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.60% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.70	1/2612 (0.0%)	0.85	9/3517 (0.3%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	187	CYS	CB-SG	-6.40	1.71	1.82

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	24	ARG	NE-CZ-NH2	-9.04	115.78	120.30
1	A	24	ARG	NE-CZ-NH1	8.83	124.72	120.30
1	A	49	ASP	CB-CG-OD2	7.04	124.63	118.30
1	A	75	ASP	CB-CG-OD2	6.29	123.96	118.30
1	A	230	ASP	CB-CG-OD2	6.05	123.75	118.30
1	A	63	ASP	CB-CG-OD2	5.54	123.28	118.30
1	A	284	ARG	NE-CZ-NH2	-5.49	117.55	120.30
1	A	235	ASP	CB-CG-OD2	5.16	122.94	118.30
1	A	294	ASP	CB-CG-OD2	5.01	122.81	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2567	0	2542	22	0
2	A	153	0	0	0	1
All	All	2720	0	2542	22	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:244:MSE:HE2	1:A:248:SER:HB3	1.56	0.84
1:A:62:THR:HG22	1:A:64:TYR:H	1.58	0.67
1:A:158:MSE:HE2	1:A:180:CYS:SG	2.37	0.65
1:A:48:VAL:CG2	1:A:341[B]:GLN:HG2	2.30	0.61
1:A:1:MSE:HE2	1:A:30:LEU:HD23	1.83	0.60
1:A:275:ALA:O	1:A:279:GLU:HG2	2.08	0.54
1:A:47:SER:HB2	1:A:340:LYS:HZ1	1.74	0.52
1:A:93:GLY:HA2	1:A:351:LEU:HD11	1.92	0.51
1:A:93:GLY:CA	1:A:351:LEU:HD11	2.39	0.51
1:A:74:ALA:HB3	1:A:103:TRP:CE3	2.47	0.50
1:A:9:ASN:HB2	1:A:24:ARG:HB2	1.94	0.48
1:A:135:MSE:HE1	1:A:296:PHE:CE2	2.48	0.48
1:A:89:LYS:HG3	1:A:90:LEU:HG	1.94	0.47
1:A:122:VAL:HG12	1:A:133:ILE:HG13	1.96	0.47
1:A:195:PHE:CE1	1:A:199:GLU:HG3	2.51	0.46
1:A:236:ILE:HD12	1:A:236:ILE:H	1.82	0.45
1:A:47:SER:HB2	1:A:340:LYS:NZ	2.31	0.44
1:A:9:ASN:OD1	1:A:24:ARG:HD3	2.20	0.41
1:A:187:CYS:O	1:A:208:TYR:CD2	2.73	0.41
1:A:118:ASP:HA	1:A:119:PRO:HD3	1.94	0.41
1:A:128:TYR:CE1	1:A:314:VAL:HG21	2.56	0.41
1:A:251:ILE:HD12	1:A:251:ILE:N	2.35	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:421:HOH:O	2:A:503:HOH:O 2_564	2.19	0.01

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	339/363 (93%)	334 (98%)	5 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	274/292 (94%)	268 (98%)	6 (2%)	52	57

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	LYS
1	A	167	CYS
1	A	187	CYS
1	A	197	THR
1	A	253	CYS
1	A	290	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	69	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	332/363 (91%)	-0.17	4 (1%) 79 82	10, 19, 34, 53	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	63	ASP	3.6
1	A	232	VAL	3.2
1	A	-1	HIS	2.9
1	A	351	LEU	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.