



# Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 10:11 am BST

PDB ID : 2VJK  
Title : Formyl-CoA transferase with aspartyl-CoA thioester intermediate derived from oxalyl-CoA  
Authors : Berthold, C.L.; Toyota, C.G.; Richards, N.G.J.; Lindqvist, Y.  
Deposited on : 2007-12-11  
Resolution : 1.97 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	11647 (2.00-1.96)
Clashscore	141614	1014 (1.98-1.98)
Ramachandran outliers	138981	1006 (1.98-1.98)
Sidechain outliers	138945	1006 (1.98-1.98)
RSRZ outliers	127900	11410 (2.00-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	428	<div> <div>4%</div> <div>91%</div> <div>8%</div> </div>
1	B	428	<div> <div>16%</div> <div>93%</div> <div>6%</div> </div>

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7566 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called FORMYL-COENZYME A TRANSFERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	427	Total	C	N	O	S	0	9	0
			3352	2125	572	631	24			
1	B	427	Total	C	N	O	S	0	3	0
			3327	2106	570	628	23			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	186	ILE	MET	conflict	UNP O06644
B	186	ILE	MET	conflict	UNP O06644

- Molecule 2 is COENZYME A (three-letter code: COA) (formula:  $C_{21}H_{36}N_7O_{16}P_3S$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P S	0	0
			48	21	7	16	3 1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	S	0	0
			48	21	7	16	3	1		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Cl	0	0
			1	1		
3	A	2	Total	Cl	0	0
			2	2		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	440	Total	O	0	0
			440	440		
5	B	347	Total	O	0	0
			347	347		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: FORMYL-COENZYME A TRANSFERASE



#### • Molecule 1: FORMYL-COENZYME A TRANSFERASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	151.91Å 151.91Å 99.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.97 48.04 – 1.97	Depositor EDS
% Data completeness (in resolution range)	98.8 (30.00-1.97) 98.8 (48.04-1.97)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.73 (at 1.97Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.173 , 0.214 0.175 , 0.219	Depositor DCC
$R_{free}$ test set	3910 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	24.9	Xtriage
Anisotropy	0.007	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 50.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.034 for -k,-h,-l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7566	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.20% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: COA, MG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.40	0/3453	0.55	0/4667
1	B	0.39	0/3411	0.54	0/4612
All	All	0.39	0/6864	0.55	0/9279

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3352	0	3318	24	0
1	B	3327	0	3268	19	0
2	A	48	0	31	0	0
2	B	48	0	31	0	0
3	A	2	0	0	0	0
3	B	1	0	0	0	0
4	A	1	0	0	0	0
5	A	440	0	0	2	0
5	B	347	0	0	1	0
All	All	7566	0	6648	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 3.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:186:ILE:HG23	1:B:4:PRO:HA	1.41	1.01
1:B:422:LEU:HD22	1:B:427:VAL:HG21	1.57	0.85
1:A:319:ASP:OD1	5:A:2325:HOH:O	2.00	0.79
1:B:295:MET:HE2	1:B:345:ILE:HD12	1.67	0.73
1:B:295:MET:CE	1:B:345:ILE:HD12	2.28	0.63
1:A:233:ASN:ND2	5:A:2227:HOH:O	2.32	0.62
1:B:422:LEU:CD2	1:B:427:VAL:HG21	2.28	0.61
1:A:83:LEU:HD23	1:A:86:MET:CE	2.31	0.61
1:A:16:VAL:HG22	1:A:17:GLN:HE21	1.71	0.56
1:B:282:PHE:CE2	1:B:324:ILE:HD11	2.41	0.55
1:A:32[A]:ASN:HD21	1:A:68:ARG:HH21	1.54	0.55
1:A:124:VAL:HG23	1:A:176:LEU:HD22	1.89	0.54
1:B:83:LEU:HD23	1:B:107:PHE:CE1	2.45	0.52
1:A:102:LEU:HD22	1:A:107:PHE:HB2	1.91	0.51
1:A:13:PHE:HE2	1:A:74:MET:HE3	1.74	0.51
1:A:208:VAL:HG12	1:A:208:VAL:O	2.11	0.50
1:B:321:PHE:HA	1:B:324:ILE:HG22	1.93	0.50
1:A:83:LEU:HD23	1:A:86:MET:HE3	1.91	0.50
1:A:36:ILE:HG12	1:A:70[A]:ILE:HD11	1.94	0.50
1:A:133:ASN:HB3	1:A:136:LEU:HD12	1.95	0.49
1:A:45:THR:O	1:A:57[A]:SER:OG	2.31	0.48
1:B:16:VAL:HA	1:B:45:THR:HG23	1.96	0.47
1:B:295:MET:CE	1:B:345:ILE:CD1	2.92	0.47
1:A:34:ILE:CG2	1:A:70[A]:ILE:HD12	2.45	0.47
1:A:331:LYS:HB3	1:A:335:GLU:HG3	1.98	0.46
1:B:82:LEU:HD21	1:B:414:LEU:HD11	1.99	0.45
1:A:74:MET:CE	1:A:83:LEU:HD11	2.47	0.45
1:B:317:LEU:O	1:B:320:ILE:HG22	2.17	0.45
1:A:179:GLY:HA3	1:A:196:VAL:HG21	2.00	0.44
1:B:205:LEU:HA	1:B:208:VAL:HG22	2.00	0.44
1:B:422:LEU:HD22	1:B:427:VAL:CG2	2.39	0.44
1:B:345:ILE:HG23	1:B:346:PRO:HD2	2.01	0.43
1:A:35:LYS:O	1:A:70[A]:ILE:HD13	2.17	0.43
1:B:17:GLN:HG3	1:B:63:PHE:CE2	2.54	0.43
1:A:209:ARG:HD2	1:B:63:PHE:CZ	2.54	0.43
1:A:13:PHE:CE2	1:A:74:MET:HE3	2.54	0.43
1:B:317:LEU:O	1:B:320:ILE:CG2	2.67	0.42
1:A:34:ILE:HG22	1:A:70[A]:ILE:HD12	2.00	0.42

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:32:ASN:HD21	1:B:68:ARG:HH21	1.66	0.42
1:A:11:LEU:HD11	1:A:36:ILE:HD11	2.01	0.42
1:A:142:VAL:HG23	5:B:2274:HOH:O	2.19	0.42

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	434/428 (101%)	426 (98%)	7 (2%)	1 (0%)	47	38
1	B	428/428 (100%)	418 (98%)	8 (2%)	2 (0%)	29	16
All	All	862/856 (101%)	844 (98%)	15 (2%)	3 (0%)	34	29

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	140	GLU
1	B	16	VAL
1	A	16	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	359/351 (102%)	346 (96%)	13 (4%)	35	23
1	B	353/351 (101%)	348 (99%)	5 (1%)	67	62
All	All	712/702 (101%)	694 (98%)	18 (2%)	50	39

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	32[A]	ASN
1	A	32[B]	ASN
1	A	59	TYR
1	A	70[A]	ILE
1	A	70[B]	ILE
1	A	102	LEU
1	A	186	ILE
1	A	205	LEU
1	A	386	PHE
1	A	392	GLN
1	A	410	LYS
1	A	415	ASP
1	A	418	LYS
1	B	59	TYR
1	B	195	LYS
1	B	317	LEU
1	B	386	PHE
1	B	392	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	206	ASN
1	A	233	ASN
1	A	392	GLN
1	B	32	ASN
1	B	113	GLN
1	B	206	ASN
1	B	342	GLN
1	B	378	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	COA	A	1169	1	41,50,50	1.11	1 (2%)	52,75,75	1.06	2 (3%)
2	COA	B	1169	1	41,50,50	1.07	1 (2%)	52,75,75	1.05	3 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	COA	A	1169	1	-	6/44/64/64	0/3/3/3
2	COA	B	1169	1	-	3/44/64/64	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1169	COA	O9P-C9P	5.25	1.33	1.23

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1169	COA	O9P-C9P	5.10	1.33	1.23

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1169	COA	N3A-C2A-N1A	-4.75	121.25	128.68
2	B	1169	COA	N3A-C2A-N1A	-4.61	121.47	128.68
2	B	1169	COA	P2A-O3A-P1A	-2.96	122.67	132.83
2	A	1169	COA	C4A-C5A-N7A	-2.07	107.24	109.40
2	B	1169	COA	C4A-C5A-N7A	-2.05	107.27	109.40

There are no chirality outliers.


All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1169	COA	C5B-O5B-P1A-O1A
2	A	1169	COA	CCP-O6A-P2A-O4A
2	A	1169	COA	CCP-O6A-P2A-O5A
2	B	1169	COA	C5B-O5B-P1A-O3A
2	B	1169	COA	C5B-O5B-P1A-O2A
2	A	1169	COA	C4B-C3B-O3B-P3B
2	A	1169	COA	C2B-C3B-O3B-P3B
2	A	1169	COA	C3B-O3B-P3B-O9A
2	A	1169	COA	CCP-O6A-P2A-O3A

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	427/428 (99%)	0.13	17 (3%) 38 40	12, 22, 48, 69	0
1	B	427/428 (99%)	0.71	67 (15%) 2 2	11, 26, 70, 86	0
All	All	854/856 (99%)	0.42	84 (9%) 7 8	11, 24, 64, 86	0

All (84) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	305	PRO	7.9
1	B	301	TRP	7.1
1	B	424	ALA	5.9
1	A	415	ASP	5.4
1	B	290	PRO	5.4
1	B	415	ASP	5.4
1	B	417	ALA	5.2
1	B	307	TYR	5.0
1	B	299	PRO	5.0
1	B	241	ASN	4.9
1	B	320	ILE	4.8
1	B	306	ALA	4.7
1	B	323	PHE	4.7
1	B	343	TYR	4.4
1	B	422	LEU	4.2
1	B	288	MET	4.1
1	B	420	LYS	4.1
1	B	296	ILE	4.1
1	B	414	LEU	4.0
1	B	294	ASP	4.0
1	B	302	LYS	4.0
1	B	297	ASP	4.0
1	B	416	ASP	3.9
1	B	303	ASP	3.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	295	MET	3.8
1	B	309	THR	3.7
1	B	413	GLY	3.7
1	B	298	LYS	3.7
1	B	427	VAL	3.7
1	B	344	GLY	3.5
1	B	287	ASN	3.5
1	A	2	THR	3.5
1	B	232	PRO	3.5
1	B	421	GLU	3.4
1	A	413	GLY	3.3
1	B	428	VAL	3.2
1	B	81	GLU	3.2
1	B	341	ALA	3.1
1	B	342	GLN	3.1
1	B	419	ILE	3.1
1	B	289	TRP	3.0
1	A	424	ALA	3.0
1	B	236	PHE	3.0
1	B	426	GLN	2.9
1	A	427	VAL	2.9
1	B	412	LEU	2.9
1	B	339	TRP	2.8
1	B	425	LYS	2.8
1	B	285	ALA	2.8
1	A	414	LEU	2.8
1	A	420	LYS	2.7
1	B	314	VAL	2.7
1	A	425	LYS	2.6
1	B	418	LYS	2.6
1	A	111	TYR	2.5
1	B	308	ASN	2.5
1	B	292	ILE	2.5
1	B	240	GLY	2.5
1	A	426	GLN	2.5
1	B	239	ASP	2.5
1	B	310	PHE	2.5
1	A	422	LEU	2.4
1	B	77	PRO	2.4
1	B	242	PRO	2.4
1	B	286	ALA	2.4
1	B	2	THR	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	419	ILE	2.4
1	B	233	ASN	2.3
1	B	315	ASP	2.3
1	B	78	GLU	2.3
1	B	311	GLU	2.3
1	B	247	ASN	2.2
1	B	330	ASP	2.2
1	A	207	LEU	2.2
1	A	105	MET	2.2
1	B	293	CYS	2.2
1	B	238	ARG	2.2
1	A	412	LEU	2.1
1	A	421	GLU	2.1
1	B	300	GLU	2.1
1	A	75	LYS	2.1
1	B	82	LEU	2.1
1	B	167	LEU	2.1
1	B	282	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	COA	B	1169	48/48	0.90	0.15	23,48,54,54	0
2	COA	A	1169	48/48	0.94	0.13	30,43,45,46	0
4	MG	A	1431	1/1	0.96	0.05	42,42,42,42	0
3	CL	A	1432	1/1	0.98	0.05	27,27,27,27	0
3	CL	A	1430	1/1	0.99	0.08	24,24,24,24	0


*Continued on next page...*




*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	CL	B	1430	1/1	0.99	0.05	22,22,22,22	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.