



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 22, 2020 – 08:56 AM BST

PDB ID : 5VJB  
Title : Guanidine-II riboswitch P2 hairpin dimer with 5-bromoU substitution from *Pseudomonas aeruginosa*  
Authors : Reiss, C.W.; Strobel, S.A.  
Deposited on : 2017-04-19  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)
RNA backbone	3102	1000 (2.54-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	16	
1	B	16	
1	C	16	
1	D	16	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GAI	C	101	-	X	-	-

## 2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 1456 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called RNA (5'-R(\*GP\*CP\*GP\*GP\*GP\*GP\*AP\*CP\*GP\*AP\*CP\*CP\*CP\*(5BU)P\*GP\*C)-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	16	Total	Br	C	N	O	P	0	0	0
			343	1	153	65	109	15			
1	B	16	Total	Br	C	N	O	P	0	0	0
			343	1	153	65	109	15			
1	C	16	Total	Br	C	N	O	P	0	0	0
			343	1	153	65	109	15			
1	D	16	Total	Br	C	N	O	P	0	0	0
			343	1	153	65	109	15			

- Molecule 2 is GUANIDINE (three-letter code: GAI) (formula: CH<sub>5</sub>N<sub>3</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	N	0	0
			4	1	3		
2	B	1	Total	C	N	0	0
			4	1	3		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	C	1	Total	C	N	0	0
			4	1	3		
2	D	1	Total	C	N	0	0
			4	1	3		

- Molecule 3 is SPERMINE (three-letter code: SPM) (formula:  $C_{10}H_{26}N_4$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	N	0	0
			14	10	4		
3	B	1	Total	C	N	0	0
			14	10	4		
3	C	1	Total	C	N	0	0
			14	10	4		
3	D	1	Total	C	N	0	0
			14	10	4		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Mg	0	0
			1	1		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total 2	O 2	0	0
5	B	5	Total 5	O 5	0	0
5	C	1	Total 1	O 1	0	0
5	D	3	Total 3	O 3	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: RNA (5'-R(\*GP\*CP\*GP\*GP\*GP\*GP\*AP\*CP\*GP\*AP\*CP\*CP\*CP\*(5BU)P\*GP\*C)-3')

Chain A: 




- Molecule 1: RNA (5'-R(\*GP\*CP\*GP\*GP\*GP\*GP\*AP\*CP\*GP\*AP\*CP\*CP\*CP\*(5BU)P\*GP\*C)-3')

Chain B: 




- Molecule 1: RNA (5'-R(\*GP\*CP\*GP\*GP\*GP\*GP\*AP\*CP\*GP\*AP\*CP\*CP\*CP\*(5BU)P\*GP\*C)-3')

Chain C: 



- Molecule 1: RNA (5'-R(\*GP\*CP\*GP\*GP\*GP\*GP\*AP\*CP\*GP\*AP\*CP\*CP\*CP\*(5BU)P\*GP\*C)-3')

Chain D: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	50.23Å 60.54Å 71.91Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 2.10 38.66 – 2.07	Depositor EDS
% Data completeness (in resolution range)	98.3 (40.00-2.10) 98.3 (38.66-2.07)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.65 (at 2.06Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, $R_{free}$	0.216 , 0.238 0.226 , 0.242	Depositor DCC
$R_{free}$ test set	634 reflections (4.64%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	46.8	Xtriage
Anisotropy	0.201	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.25 , 30.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	1456	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.63% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 5BU, MG, SPM, GAI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.82	1/360 (0.3%)	0.98	1/561 (0.2%)
1	B	0.73	0/360	1.04	1/561 (0.2%)
1	C	1.39	5/360 (1.4%)	0.92	1/561 (0.2%)
1	D	0.75	1/360 (0.3%)	0.96	0/561
All	All	0.96	7/1440 (0.5%)	0.97	3/2244 (0.1%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	10	A	O3'-P	-8.02	1.51	1.61
1	C	9	G	O3'-P	-7.48	1.52	1.61
1	C	7	A	O3'-P	-5.40	1.54	1.61
1	D	8	C	O3'-P	-5.38	1.54	1.61
1	A	9	G	O3'-P	-5.33	1.54	1.61
1	C	12	C	O3'-P	-5.24	1.54	1.61
1	C	8	C	O3'-P	-5.06	1.55	1.61

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	3	G	O5'-P-OP2	-9.45	97.20	105.70
1	C	9	G	C4'-C3'-O3'	7.89	128.77	113.00
1	A	7	A	O5'-P-OP1	-7.54	98.91	105.70

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	343	0	176	5	0
1	B	343	0	176	1	0
1	C	343	0	176	1	0
1	D	343	0	176	0	0
2	A	4	0	4	0	0
2	B	4	0	4	0	0
2	C	4	0	4	0	0
2	D	4	0	4	0	0
3	A	14	0	26	3	0
3	B	14	0	26	0	0
3	C	14	0	26	0	0
3	D	14	0	26	0	0
4	B	1	0	0	0	0
5	A	2	0	0	0	0
5	B	5	0	0	0	0
5	C	1	0	0	0	0
5	D	3	0	0	0	0
All	All	1456	0	824	7	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (7) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:6:G:N7	3:A:102:SPM:H61	2.17	0.58
1:A:10:A:N3	1:A:10:A:H2'	2.25	0.51
1:A:8:C:OP1	3:A:102:SPM:H71	2.16	0.46
1:A:6:G:N7	3:A:102:SPM:C6	2.79	0.45
1:B:7:A:C5	1:B:8:C:C5	3.06	0.43
1:A:14:5BU:O5'	1:A:14:5BU:H6	2.20	0.42
1:C:7:A:C6	1:C:8:C:C5	3.10	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

### 5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

### 5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	15/16 (93%)	1 (6%)	1 (6%)
1	B	15/16 (93%)	0	0
1	C	15/16 (93%)	1 (6%)	2 (13%)
1	D	15/16 (93%)	1 (6%)	1 (6%)
All	All	60/64 (93%)	3 (5%)	4 (6%)

All (3) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	A	3	G
1	C	10	A
1	D	10	A

All (4) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	A	10	A
1	C	9	G
1	C	10	A
1	D	10	A

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	5BU	B	14	1	15,22,23	2.20	4 (26%)	17,32,35	2.65	2 (11%)
1	5BU	A	14	1	15,22,23	1.68	3 (20%)	17,32,35	1.82	2 (11%)
1	5BU	D	14	1	15,22,23	2.11	5 (33%)	17,32,35	2.61	4 (23%)
1	5BU	C	14	1	15,22,23	1.48	1 (6%)	17,32,35	1.99	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	5BU	B	14	1	-	0/5/25/26	0/2/2/2
1	5BU	A	14	1	-	0/5/25/26	0/2/2/2
1	5BU	D	14	1	-	0/5/25/26	0/2/2/2
1	5BU	C	14	1	-	0/5/25/26	0/2/2/2

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	14	5BU	C4-C5	6.54	1.46	1.38
1	D	14	5BU	C4-C5	5.96	1.46	1.38
1	C	14	5BU	C4-C5	4.37	1.44	1.38
1	A	14	5BU	C2'-C1'	-4.32	1.47	1.53
1	B	14	5BU	O4'-C1'	2.91	1.45	1.41
1	B	14	5BU	C2'-C1'	-2.82	1.49	1.53
1	D	14	5BU	C2'-C1'	-2.74	1.49	1.53
1	A	14	5BU	C4-C5	2.71	1.41	1.38
1	B	14	5BU	C2-N3	-2.51	1.33	1.38
1	D	14	5BU	BR-C5	-2.34	1.83	1.90
1	D	14	5BU	O4'-C1'	2.22	1.44	1.41
1	D	14	5BU	O4'-C4'	-2.21	1.40	1.45
1	A	14	5BU	C2-N3	-2.04	1.34	1.38

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	14	5BU	C4-N3-C2	8.31	122.16	115.14
1	D	14	5BU	C4-N3-C2	7.16	121.19	115.14

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	14	5BU	C4-N3-C2	6.36	120.51	115.14
1	D	14	5BU	C5-C4-N3	-6.16	116.26	123.64
1	A	14	5BU	C4-N3-C2	5.65	119.91	115.14
1	B	14	5BU	C5-C4-N3	-5.62	116.91	123.64
1	C	14	5BU	C5-C4-N3	-3.94	118.92	123.64
1	A	14	5BU	C5-C4-N3	-3.66	119.26	123.64
1	D	14	5BU	C3'-C2'-C1'	-3.30	96.02	100.98
1	C	14	5BU	C5-C6-N1	2.18	122.79	119.97
1	D	14	5BU	BR-C5-C4	-2.15	118.24	121.50

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	14	5BU	1	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 1 is monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SPM	D	102	-	13,13,13	0.53	0	12,12,12	0.81	0
2	GAI	B	101	-	3,3,3	2.84	2 (66%)	3,3,3	0.45	0
2	GAI	C	101	-	3,3,3	1.66	1 (33%)	3,3,3	1.86	2 (66%)
2	GAI	A	101	-	3,3,3	2.88	2 (66%)	3,3,3	1.47	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SPM	A	102	-	13,13,13	0.45	0	12,12,12	0.87	0
3	SPM	C	102	-	13,13,13	0.58	0	12,12,12	0.87	0
2	GAI	D	101	-	3,3,3	1.56	0	3,3,3	1.01	0
3	SPM	B	102	-	13,13,13	0.63	0	12,12,12	0.71	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SPM	D	102	-	-	4/11/11/11	-
3	SPM	A	102	-	-	6/11/11/11	-
3	SPM	C	102	-	-	6/11/11/11	-
3	SPM	B	102	-	-	8/11/11/11	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	101	GAI	C-N3	-4.32	1.28	1.36
2	B	101	GAI	C-N2	-3.85	1.29	1.36
2	A	101	GAI	C-N2	-2.42	1.31	1.36
2	B	101	GAI	C-N3	-2.40	1.31	1.36
2	C	101	GAI	C-N1	2.08	1.35	1.30

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	101	GAI	N3-C-N2	2.37	121.77	116.13
2	C	101	GAI	N3-C-N1	-2.18	115.74	121.92

There are no chirality outliers.

All (24) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	102	SPM	N5-C6-C7-C8
3	C	102	SPM	C7-C8-C9-N10
3	D	102	SPM	N10-C11-C12-C13
3	A	102	SPM	C3-C4-N5-C6
3	B	102	SPM	C7-C8-C9-N10

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
3	A	102	SPM	N10-C11-C12-C13
3	B	102	SPM	N10-C11-C12-C13
3	C	102	SPM	C8-C9-N10-C11
3	C	102	SPM	C12-C11-N10-C9
3	B	102	SPM	C8-C9-N10-C11
3	A	102	SPM	C7-C8-C9-N10
3	A	102	SPM	C12-C11-N10-C9
3	B	102	SPM	C7-C6-N5-C4
3	D	102	SPM	C12-C11-N10-C9
3	B	102	SPM	N1-C2-C3-C4
3	D	102	SPM	C2-C3-C4-N5
3	A	102	SPM	N1-C2-C3-C4
3	C	102	SPM	N1-C2-C3-C4
3	B	102	SPM	C2-C3-C4-N5
3	C	102	SPM	C7-C6-N5-C4
3	C	102	SPM	C3-C4-N5-C6
3	B	102	SPM	C11-C12-C13-N14
3	D	102	SPM	C6-C7-C8-C9
3	A	102	SPM	N5-C6-C7-C8

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	102	SPM	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	15/16 (93%)	-0.52	0 100 100	34, 41, 46, 58	0
1	B	15/16 (93%)	-0.51	0 100 100	36, 39, 44, 60	0
1	C	15/16 (93%)	-0.61	0 100 100	36, 39, 43, 55	0
1	D	15/16 (93%)	-0.58	0 100 100	36, 40, 44, 44	0
All	All	60/64 (93%)	-0.55	0 100 100	34, 41, 46, 60	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	5BU	A	14	21/22	0.88	0.15	35,43,50,61	0
1	5BU	C	14	21/22	0.91	0.12	37,40,47,54	0
1	5BU	D	14	21/22	0.94	0.12	40,44,50,51	0
1	5BU	B	14	21/22	0.94	0.14	34,38,43,48	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.



## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	SPM	C	102	14/14	0.72	0.31	55,65,96,102	0
3	SPM	D	102	14/14	0.74	0.29	54,63,76,78	0
3	SPM	A	102	14/14	0.79	0.21	52,58,64,67	0
3	SPM	B	102	14/14	0.81	0.43	48,72,91,95	0
2	GAI	C	101	4/4	0.91	0.16	32,35,36,39	0
4	MG	B	103	1/1	0.92	0.21	30,30,30,30	0
2	GAI	B	101	4/4	0.92	0.20	38,39,43,43	0
2	GAI	D	101	4/4	0.94	0.23	30,34,35,38	0
2	GAI	A	101	4/4	0.94	0.18	34,36,37,39	0

## 6.5 Other polymers

There are no such residues in this entry.