



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 22, 2020 – 01:12 AM BST

PDB ID : 6VJB
Title : Crystal structure of a catalytically inactive CXC Chemokine-degrading protease SpyCEP from *Streptococcus pyogenes*
Authors : Malito, E.; Rouse, S.
Deposited on : 2020-01-15
Resolution : 2.24 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.24 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2391 (2.26-2.22)
Clashscore	141614	2539 (2.26-2.22)
Ramachandran outliers	138981	2489 (2.26-2.22)
Sidechain outliers	138945	2490 (2.26-2.22)
RSRZ outliers	127900	2353 (2.26-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1589	<div> <div>5%</div> <div> <div></div> <div>76%</div> <div>8%</div> <div>15%</div> </div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 11195 atoms, of which 24 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative cell envelope proteinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1346	Total	C	N	O	S	0	0	0
			10461	6591	1782	2063	25			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	33	MET	-	initiating methionine	UNP Q9A180
A	151	ALA	ASP	engineered mutation	UNP Q9A180
A	617	ALA	SER	engineered mutation	UNP Q9A180
A	1615	GLU	-	expression tag	UNP Q9A180
A	1616	HIS	-	expression tag	UNP Q9A180
A	1617	HIS	-	expression tag	UNP Q9A180
A	1618	HIS	-	expression tag	UNP Q9A180
A	1619	HIS	-	expression tag	UNP Q9A180
A	1620	HIS	-	expression tag	UNP Q9A180
A	1621	HIS	-	expression tag	UNP Q9A180

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	Ca	0	0
			3	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	691	Total	O	0	0
			691	691		

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- [illegible]

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	140.50Å 121.75Å 106.64Å 90.00° 111.78° 90.00°	Depositor
Resolution (Å)	44.51 – 2.24 65.24 – 2.24	Depositor EDS
% Data completeness (in resolution range)	96.5 (44.51-2.24) 96.5 (65.24-2.24)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.41 (at 2.25Å)	Xtriage
Refinement program	BUSTER 2.11.7	Depositor
R, R_{free}	0.190 , 0.244 0.203 , 0.261	Depositor DCC
R_{free} test set	3911 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	35.2	Xtriage
Anisotropy	0.234	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 55.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	11195	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	1/10660 (0.0%)	0.69	0/14420

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	404	SER	C-N	8.28	1.50	1.34

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	10461	0	10283	76	0
2	A	16	24	24	1	0
3	A	3	0	0	0	0
4	A	691	0	0	3	0
All	All	11171	24	10307	76	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:549:SER:HA	1:A:552:ASN:HB2	1.68	0.76
1:A:1312:LEU:N	1:A:1312:LEU:HD12	2.01	0.75
1:A:320:ILE:N	1:A:320:ILE:HD12	2.06	0.70
1:A:585:THR:CG2	1:A:587:ASP:OD1	2.40	0.70
1:A:321:MET:HE1	4:A:2262:HOH:O	1.94	0.68
1:A:1493:SER:OG	1:A:1560:THR:HG23	1.94	0.67
1:A:1023:LEU:HD13	1:A:1118:ALA:HB2	1.77	0.66
1:A:320:ILE:H	1:A:320:ILE:HD12	1.60	0.66
1:A:501:ILE:HG23	1:A:512:VAL:HG21	1.81	0.62
1:A:825:VAL:H	1:A:1008:GLN:HE22	1.48	0.62
1:A:1538:GLY:H	1:A:1560:THR:HG22	1.64	0.61
1:A:1148:LEU:HD23	1:A:1180:LEU:HG	1.84	0.60
1:A:875:ILE:HG12	1:A:1077:VAL:HG12	1.84	0.59
1:A:320:ILE:CD1	1:A:320:ILE:H	2.16	0.58
1:A:585:THR:HG23	1:A:587:ASP:OD1	2.04	0.58
1:A:1546:LEU:HD11	1:A:1552:ILE:HG12	1.85	0.58
1:A:675:GLN:OE1	1:A:678:GLY:HA2	2.05	0.56
1:A:385:TYR:HD2	1:A:958:LYS:HD3	1.72	0.55
1:A:348:GLY:HA2	1:A:404:SER:O	2.07	0.54
1:A:470:LYS:HG3	1:A:553:GLY:HA2	1.91	0.53
1:A:429:VAL:HG12	1:A:429:VAL:O	2.09	0.53
1:A:977:PHE:CD2	1:A:1014:MET:HE1	2.45	0.52
1:A:1523:GLN:HE21	1:A:1528:PRO:HB3	1.75	0.52
1:A:1434:GLU:HB3	1:A:1444:ILE:HB	1.92	0.52
1:A:1500:ASP:HB2	1:A:1571:ARG:HA	1.91	0.51
1:A:1405:LEU:HD23	1:A:1485:PHE:HB2	1.93	0.51
1:A:1423:ARG:HB2	1:A:1451:THR:HG22	1.93	0.51
1:A:320:ILE:CD1	1:A:320:ILE:N	2.73	0.50
1:A:1188:PHE:CD2	1:A:1314:SER:HB3	2.46	0.50
1:A:1421:LEU:HD11	1:A:1429:PRO:HB3	1.93	0.49
1:A:1537:GLU:HA	1:A:1560:THR:CG2	2.41	0.49
1:A:693:THR:HG22	1:A:715:THR:HB	1.94	0.49
1:A:1042:ASP:HB2	1:A:1048:VAL:HG23	1.94	0.49
1:A:1312:LEU:N	1:A:1312:LEU:CD1	2.73	0.49
1:A:1421:LEU:HB2	1:A:1455:LEU:HD11	1.96	0.48
1:A:777:GLN:HG2	4:A:1860:HOH:O	2.14	0.48
1:A:201:ASN:HA	1:A:314:ARG:HB3	1.96	0.48
1:A:1215:VAL:HG22	1:A:1265:VAL:HG22	1.97	0.47
1:A:706:ILE:HD11	1:A:815:GLY:HA3	1.96	0.47
1:A:457:SER:HB2	1:A:466:PHE:HE2	1.80	0.46
1:A:452:LYS:HE2	1:A:455:LYS:HD2	1.97	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:199:ALA:HB1	1:A:209:ILE:HG21	1.98	0.46
1:A:1321:GLU:HB2	1:A:1375:LEU:HB3	1.97	0.46
1:A:147:VAL:HB	1:A:628:LEU:HD11	1.98	0.46
1:A:651:LYS:HE2	4:A:2244:HOH:O	2.15	0.46
1:A:1387:THR:HG22	1:A:1390:ASP:H	1.81	0.46
1:A:153:GLY:HA3	1:A:211:GLU:HB2	1.97	0.46
1:A:1073:LYS:HD2	1:A:1333:PHE:CE1	2.52	0.45
1:A:1214:THR:HG22	1:A:1216:ASN:HD21	1.80	0.45
1:A:544:PHE:CZ	1:A:549:SER:HB2	2.51	0.45
1:A:429:VAL:HG12	1:A:432:LEU:HB2	1.99	0.45
1:A:447:GLU:OE2	1:A:453:ASP:HB2	2.17	0.44
1:A:1217:VAL:HA	1:A:1262:GLN:O	2.17	0.44
1:A:515:PHE:HB3	1:A:544:PHE:CE2	2.53	0.44
1:A:1187:PHE:HB3	1:A:1281:ILE:HD13	1.98	0.43
1:A:1214:THR:HG22	1:A:1216:ASN:ND2	2.33	0.43
1:A:432:LEU:HD21	1:A:551:LEU:HD12	2.00	0.43
1:A:1513:THR:HG23	1:A:1515:GLN:H	1.84	0.43
1:A:693:THR:HG23	1:A:694:GLY:O	2.18	0.43
1:A:486:ILE:HG12	1:A:511:GLY:HA3	2.00	0.42
1:A:488:LEU:HD11	1:A:549:SER:OG	2.19	0.42
1:A:1004:GLY:H	2:A:1702:EDO:H22	1.85	0.42
1:A:470:LYS:HA	1:A:488:LEU:HB3	2.01	0.41
1:A:1095:PRO:HD2	1:A:1104:PHE:CE1	2.55	0.41
1:A:872:THR:HG21	1:A:1342:ILE:HD12	2.03	0.41
1:A:1214:THR:CG2	1:A:1216:ASN:HD21	2.34	0.41
1:A:1073:LYS:HD2	1:A:1333:PHE:HE1	1.86	0.41
1:A:426:LEU:HD12	1:A:566:SER:HB3	2.03	0.41
1:A:859:THR:HA	1:A:922:GLY:HA2	2.03	0.41
1:A:1154:MET:HE2	1:A:1182:LYS:HE2	2.03	0.41
1:A:429:VAL:HG22	1:A:558:SER:O	2.20	0.41
1:A:1551:ARG:HH21	1:A:1573:VAL:HG11	1.85	0.41
1:A:192:ILE:CD1	1:A:199:ALA:HB3	2.51	0.41
1:A:150:ILE:HG12	1:A:313:MET:HB2	2.03	0.40
1:A:590:LEU:HD23	1:A:700:SER:HB2	2.03	0.40
1:A:1513:THR:HG22	1:A:1517:GLN:H	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1322/1589 (83%)	1288 (97%)	34 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1146/1354 (85%)	1108 (97%)	38 (3%)	38	43

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	166	VAL
1	A	273	ASP
1	A	369	LYS
1	A	431	GLU
1	A	448	SER
1	A	450	ASP
1	A	453	ASP
1	A	540	ILE
1	A	544	PHE
1	A	551	LEU
1	A	552	ASN
1	A	566	SER
1	A	585	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	587	ASP
1	A	693	THR
1	A	697	ASN
1	A	810	ASN
1	A	866	SER
1	A	870	VAL
1	A	872	THR
1	A	964	ILE
1	A	1022	VAL
1	A	1061	ASN
1	A	1096	LEU
1	A	1155	THR
1	A	1210	TYR
1	A	1221	ASP
1	A	1294	ARG
1	A	1312	LEU
1	A	1313	ASP
1	A	1362	SER
1	A	1451	THR
1	A	1482	GLN
1	A	1512	LYS
1	A	1513	THR
1	A	1560	THR
1	A	1569	SER
1	A	1575	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	465	GLN
1	A	810	ASN
1	A	1008	GLN
1	A	1216	ASN
1	A	1402	ASN
1	A	1432	ASN
1	A	1523	GLN
1	A	1536	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 3 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	A	1703	-	3,3,3	0.60	0	2,2,2	0.25	0
2	EDO	A	1702	-	3,3,3	0.65	0	2,2,2	0.24	0
2	EDO	A	1704	-	3,3,3	0.14	0	2,2,2	0.32	0
2	EDO	A	1701	-	3,3,3	0.58	0	2,2,2	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	1703	-	-	0/1/1/1	-
2	EDO	A	1702	-	-	1/1/1/1	-
2	EDO	A	1704	-	-	1/1/1/1	-
2	EDO	A	1701	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1704	EDO	O1-C1-C2-O2
2	A	1702	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1702	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1346/1589 (84%)	0.17	75 (5%) 24 23	24, 48, 91, 121	0

All (75) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	501	ILE	6.2
1	A	1213	LEU	5.9
1	A	1209	VAL	5.3
1	A	1210	TYR	5.3
1	A	542	HIS	4.9
1	A	469	VAL	4.8
1	A	1072	TYR	4.6
1	A	1207	ASN	4.5
1	A	1275	HIS	4.3
1	A	1239	SER	4.2
1	A	1125	PRO	4.1
1	A	321	MET	3.9
1	A	473	THR	3.6
1	A	544	PHE	3.6
1	A	1236	ALA	3.5
1	A	1214	THR	3.5
1	A	539	PHE	3.4
1	A	1265	VAL	3.3
1	A	558	SER	3.3
1	A	1326	ALA	3.1
1	A	451	PHE	3.1
1	A	452	LYS	3.1
1	A	1212	ASP	3.1
1	A	540	ILE	3.1
1	A	1301	VAL	3.0
1	A	489	ILE	3.0
1	A	1202	PHE	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1208	ASN	3.0
1	A	1276	GLN	2.9
1	A	1043	ARG	2.9
1	A	465	GLN	2.8
1	A	1237	SER	2.8
1	A	499	GLU	2.8
1	A	1238	VAL	2.8
1	A	557	GLY	2.7
1	A	1325	LEU	2.7
1	A	1432	ASN	2.7
1	A	512	VAL	2.7
1	A	1332	LYS	2.7
1	A	1241	ILE	2.7
1	A	503	LEU	2.7
1	A	1313	ASP	2.6
1	A	322	GLY	2.6
1	A	458	LEU	2.6
1	A	1407	LEU	2.6
1	A	459	GLY	2.6
1	A	1069	ASN	2.5
1	A	548	MET	2.5
1	A	511	GLY	2.5
1	A	1240	ALA	2.4
1	A	1131	THR	2.4
1	A	468	TYR	2.4
1	A	474	ASP	2.4
1	A	1074	TYR	2.4
1	A	1235	GLY	2.4
1	A	323	SER	2.4
1	A	456	ASP	2.4
1	A	1224	GLN	2.4
1	A	1226	GLN	2.3
1	A	551	LEU	2.2
1	A	1168	LEU	2.2
1	A	475	ALA	2.2
1	A	555	GLY	2.2
1	A	550	GLN	2.2
1	A	1367	VAL	2.2
1	A	1150	ASP	2.2
1	A	320	ILE	2.2
1	A	1234	ALA	2.1
1	A	1433	LEU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	449	VAL	2.1
1	A	1277	LYS	2.0
1	A	1156	GLN	2.0
1	A	561	PHE	2.0
1	A	454	ILE	2.0
1	A	1227	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	EDO	A	1704	4/4	0.82	0.24	15,17,18,18	0
2	EDO	A	1703	4/4	0.87	0.13	69,73,76,76	0
2	EDO	A	1702	4/4	0.87	0.13	40,41,41,41	0
2	EDO	A	1701	4/4	0.96	0.07	52,53,55,55	0
3	CA	A	1705	1/1	0.97	0.06	48,48,48,48	0
3	CA	A	1706	1/1	0.97	0.09	35,35,35,35	0
3	CA	A	1707	1/1	0.98	0.04	59,59,59,59	0

6.5 Other polymers [i](#)

There are no such residues in this entry.