



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 17, 2022 – 12:07 PM JST

PDB ID : 7VJY
Title : Crystal Structure of Sars-Cov-2 Mpro at 1.90 Å resolution-1
Authors : DeMirici, H.; Johnson, A.J.
Deposited on : 2021-09-29
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.25
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.25


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	306	<div> <div>25%</div> <div>81%</div> <div>16%</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2487 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3C-like proteinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	306	Total	C	N	O	S	0	7	0
			2427	1535	413	454	25			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	60	Total	O	0	0
			60	60		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 3C-like proteinase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	114.00Å 53.50Å 45.00Å 90.00° 102.00° 90.00°	Depositor
Resolution (Å)	48.24 – 1.90 48.24 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.24-1.90) 100.0 (48.24-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.08 (at 1.90Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.233 , 0.278 0.228 , 0.272	Depositor DCC
R_{free} test set	2000 reflections (9.50%)	wwPDB-VP
Wilson B-factor (Å ²)	35.2	Xtriage
Anisotropy	0.167	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 54.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.39$, $\langle L^2 \rangle = 0.23$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2487	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.49	1/2480 (0.0%)	0.67	0/3367

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	145	CYS	CB-SG	-5.10	1.73	1.81

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	276[A]	MET	Peptide
1	A	276[B]	MET	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2427	0	2371	34	0
2	A	60	0	0	7	0
All	All	2487	0	2371	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:21:THR:HG1	1:A:26:THR:HG1	0.99	0.94
1:A:145:CYS:SG	2:A:459:HOH:O	2.39	0.80
1:A:279:ARG:HE	1:A:280:THR:H	1.25	0.80
1:A:226:THR:HG23	1:A:229:ASP:H	1.55	0.72
1:A:5:LYS:O	2:A:401:HOH:O	2.09	0.70
1:A:262:LEU:HB2	2:A:427:HOH:O	1.95	0.66
1:A:245:ASP:OD1	2:A:402:HOH:O	2.13	0.65
1:A:197:ASP:OD2	2:A:403:HOH:O	2.16	0.62
1:A:17:MET:HG3	1:A:117:CYS:SG	2.42	0.59
1:A:201:THR:H	1:A:240:GLU:HG2	1.69	0.58
1:A:279:ARG:HE	1:A:280:THR:N	2.00	0.57
1:A:118:TYR:CE1	1:A:144:SER:HB3	2.39	0.57
1:A:201:THR:H	1:A:240:GLU:CG	2.20	0.53
1:A:279:ARG:HA	1:A:279:ARG:NE	2.24	0.53
1:A:5:LYS:HA	1:A:291:PHE:CZ	2.47	0.50
1:A:36:VAL:HG21	1:A:68:VAL:HG11	1.95	0.48
1:A:256:GLN:OE1	2:A:404:HOH:O	2.20	0.48
1:A:12:LYS:NZ	2:A:410:HOH:O	2.44	0.47
1:A:102:LYS:HG3	1:A:104:VAL:HG13	1.98	0.46
1:A:83:GLN:O	1:A:86:VAL:HG12	2.16	0.46
1:A:226:THR:CG2	1:A:229:ASP:HB2	2.46	0.46
1:A:22:CYS:SG	1:A:61:LYS:HE2	2.56	0.45
1:A:163:HIS:CE1	1:A:172:HIS:HB3	2.52	0.45
1:A:276[A]:MET:O	1:A:277:ASN:HB2	2.17	0.44
1:A:22:CYS:HB2	1:A:42:VAL:HG22	1.99	0.44
1:A:200:ILE:HA	1:A:240:GLU:HG2	1.98	0.44
1:A:113:SER:O	1:A:149:GLY:HA2	2.18	0.42
1:A:114:VAL:O	1:A:125:VAL:HA	2.18	0.42
1:A:63:ASN:OD1	1:A:80:HIS:ND1	2.44	0.42
1:A:175:THR:HG22	1:A:181:PHE:HA	2.01	0.42
1:A:212:VAL:HG22	1:A:220:LEU:HD21	2.02	0.41
1:A:187:ASP:OD1	1:A:187:ASP:N	2.54	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:19:GLN:O	1:A:68:VAL:HA	2.21	0.41
1:A:31:TRP:CD2	1:A:95:ASN:HB2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	311/306 (102%)	297 (96%)	10 (3%)	4 (1%)	12 4

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	277	ASN
1	A	6[A]	MET
1	A	6[B]	MET
1	A	154	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	270/263 (103%)	255 (94%)	15 (6%)	21 11

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	LYS
1	A	34	ASP
1	A	73	VAL
1	A	87	LEU
1	A	165[A]	MET
1	A	165[B]	MET
1	A	181	PHE
1	A	189[A]	GLN
1	A	189[B]	GLN
1	A	222	ARG
1	A	229	ASP
1	A	263	ASP
1	A	276[A]	MET
1	A	276[B]	MET
1	A	279	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	306/306 (100%)	1.64	78 (25%) 0 0	20, 35, 64, 89	0

All (78) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	154	TYR	7.8
1	A	72	ASN	7.3
1	A	306	GLN	6.8
1	A	79	GLY	6.4
1	A	222	ARG	6.2
1	A	277	ASN	6.0
1	A	24	THR	6.0
1	A	74	GLN	5.9
1	A	73	VAL	5.5
1	A	276[A]	MET	5.2
1	A	23	GLY	4.9
1	A	153	ASP	4.9
1	A	223	PHE	4.8
1	A	22	CYS	4.8
1	A	75	LEU	4.8
1	A	279	ARG	4.2
1	A	78	ILE	4.2
1	A	91	VAL	4.1
1	A	229	ASP	3.7
1	A	65	ASN	3.7
1	A	62	SER	3.5
1	A	70	ALA	3.5
1	A	226	THR	3.4
1	A	155	ASP	3.4
1	A	64	HIS	3.3
1	A	67	LEU	3.2
1	A	235	MET	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	303	VAL	3.1
1	A	45	THR	3.1
1	A	95	ASN	3.0
1	A	92	ASP	2.9
1	A	77	VAL	2.8
1	A	13	VAL	2.8
1	A	232	LEU	2.8
1	A	227	LEU	2.8
1	A	46	SER	2.8
1	A	89	LEU	2.7
1	A	60	ARG	2.7
1	A	208	LEU	2.6
1	A	213	ILE	2.6
1	A	59	ILE	2.5
1	A	42	VAL	2.5
1	A	114	VAL	2.5
1	A	32	LEU	2.5
1	A	272	LEU	2.5
1	A	142	ASN	2.5
1	A	247	VAL	2.4
1	A	274	ASN	2.4
1	A	220	LEU	2.4
1	A	80	HIS	2.4
1	A	27	LEU	2.3
1	A	297[A]	VAL	2.3
1	A	141	LEU	2.3
1	A	135	THR	2.3
1	A	94	ALA	2.3
1	A	278	GLY	2.2
1	A	233	VAL	2.2
1	A	234	ALA	2.2
1	A	296	VAL	2.2
1	A	230	PHE	2.2
1	A	304	THR	2.2
1	A	165[A]	MET	2.2
1	A	160	CYS	2.2
1	A	117	CYS	2.2
1	A	193	ALA	2.1
1	A	221	ASN	2.1
1	A	271	LEU	2.1
1	A	85	CYS	2.1
1	A	219	PHE	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	280	THR	2.1
1	A	156	CYS	2.1
1	A	265	CYS	2.1
1	A	115	LEU	2.1
1	A	301	SER	2.1
1	A	212	VAL	2.1
1	A	21	THR	2.0
1	A	294	PHE	2.0
1	A	262	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.