



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 10, 2022 – 12:42 PM JST

PDB ID : 7VOX
Title : The crystal structure of human forkhead box protein A in complex with DNA
2
Authors : Choi, Y.; Yoon, H.J.; Lee, H.H.
Deposited on : 2021-10-15
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at <http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.29
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.29


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	D	16	<div> <div></div> <div>50%50%</div> </div>
1	F	16	<div> <div></div> <div>56%44%</div> </div>
2	E	16	<div> <div></div> <div>56%44%</div> </div>
2	G	16	<div> <div></div> <div>62%38%</div> </div>
3	A	102	<div> <div>3%</div> <div>76%7%17%</div> </div>
3	B	102	<div> <div>10%</div> <div>73%8%19%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	C	102	<div><div></div><div>11%</div><div>77%</div><div>6%</div><div>17%</div></div>
3	H	102	<div><div></div><div>3%</div><div>76%</div><div>6%</div><div>17%</div></div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4564 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(P*TP*CP*GP*AP*TP*AP*AP*TP*AP*AP*AP*TP*AP*TP*TP*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	D	16	Total	C	N	O	P	0	0	0
			328	159	57	96	16			
1	F	16	Total	C	N	O	P	0	0	0
			328	159	57	96	16			

- Molecule 2 is a DNA chain called DNA (5'-D(P*AP*AP*AP*TP*AP*TP*TP*TP*AP*TP*TP*AP*TP*CP*GP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	16	Total	C	N	O	P	0	0	0
			328	159	57	96	16			
2	G	16	Total	C	N	O	P	0	0	0
			328	159	57	96	16			

- Molecule 3 is a protein called Hepatocyte nuclear factor 3-alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	85	Total	C	N	O	S	0	0	0
			704	453	122	125	4			
3	B	83	Total	C	N	O	S	0	0	0
			692	447	119	122	4			
3	C	85	Total	C	N	O	S	0	0	0
			704	453	122	125	4			
3	H	85	Total	C	N	O	S	0	0	0
			704	453	122	125	4			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	163	GLY	-	expression tag	UNP P55317
A	164	ALA	-	expression tag	UNP P55317

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	165	MET	-	expression tag	UNP P55317
A	166	GLY	-	expression tag	UNP P55317
A	167	SER	-	expression tag	UNP P55317
B	163	GLY	-	expression tag	UNP P55317
B	164	ALA	-	expression tag	UNP P55317
B	165	MET	-	expression tag	UNP P55317
B	166	GLY	-	expression tag	UNP P55317
B	167	SER	-	expression tag	UNP P55317
C	163	GLY	-	expression tag	UNP P55317
C	164	ALA	-	expression tag	UNP P55317
C	165	MET	-	expression tag	UNP P55317
C	166	GLY	-	expression tag	UNP P55317
C	167	SER	-	expression tag	UNP P55317
H	163	GLY	-	expression tag	UNP P55317
H	164	ALA	-	expression tag	UNP P55317
H	165	MET	-	expression tag	UNP P55317
H	166	GLY	-	expression tag	UNP P55317
H	167	SER	-	expression tag	UNP P55317

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Mg 1 1	0	0
4	B	1	Total Mg 1 1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	D	39	Total O 39 39	0	0
5	E	33	Total O 33 33	0	0
5	F	44	Total O 44 44	0	0
5	G	45	Total O 45 45	0	0
5	A	83	Total O 83 83	0	0
5	B	65	Total O 65 65	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	64	Total 64	O 64	0	0
5	H	73	Total 73	O 73	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA (5'-D(P*TP*CP*GP*AP*TP*AP*AP*TP*AP*AP*AP*TP*AP*TP*TP*T)-3')

Chain D: 



- Molecule 1: DNA (5'-D(P*TP*CP*GP*AP*TP*AP*AP*TP*AP*AP*AP*TP*AP*TP*TP*T)-3')

Chain F: 



- Molecule 2: DNA (5'-D(P*AP*AP*AP*TP*AP*TP*TP*TP*AP*TP*TP*AP*TP*CP*GP*A)-3')

Chain E: 





- Molecule 2: DNA (5'-D(P*AP*AP*AP*TP*AP*TP*TP*TP*AP*TP*TP*AP*TP*CP*GP*A)-3')

Chain G: 




- Molecule 3: Hepatocyte nuclear factor 3-alpha



Chain A: 





- Molecule 3: Hepatocyte nuclear factor 3-alpha



- Molecule 3: Hepatocyte nuclear factor 3-alpha



- Molecule 3: Hepatocyte nuclear factor 3-alpha



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	138.55Å 144.29Å 75.03Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.98 – 2.10 43.98 – 2.10	Depositor EDS
% Data completeness (in resolution range)	93.0 (43.98-2.10) 93.0 (43.98-2.10)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.90 (at 2.10Å)	Xtriage
Refinement program	PHENIX 1.13_2998	Depositor
R, R_{free}	0.186 , 0.212 0.186 , 0.212	Depositor DCC
R_{free} test set	1914 reflections (4.65%)	wwPDB-VP
Wilson B-factor (Å ²)	25.3	Xtriage
Anisotropy	0.064	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 64.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.110 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4564	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	D	0.63	0/367	1.08	0/564
1	F	0.66	0/367	1.08	0/564
2	E	0.65	0/367	1.03	0/564
2	G	0.63	0/367	1.03	0/564
3	A	0.27	0/728	0.43	0/989
3	B	0.30	0/716	0.49	0/973
3	C	0.30	0/728	0.52	0/989
3	H	0.32	0/728	0.53	1/989 (0.1%)
All	All	0.44	0/4368	0.75	1/6196 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	H	209	GLN	CA-CB-CG	5.27	124.99	113.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	328	0	184	8	1
1	F	328	0	184	9	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	E	328	0	184	7	0
2	G	328	0	184	12	0
3	A	704	0	681	7	0
3	B	692	0	672	5	1
3	C	704	0	681	8	0
3	H	704	0	681	11	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	83	0	0	1	0
5	B	65	0	0	1	0
5	C	64	0	0	2	0
5	D	39	0	0	2	0
5	E	33	0	0	3	0
5	F	44	0	0	2	0
5	G	45	0	0	4	0
5	H	73	0	0	2	0
All	All	4564	0	3451	51	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 51 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:G:1:DA:P	3:C:213:ARG:CZ	2.57	0.91
2:G:1:DA:P	3:C:213:ARG:NH2	2.50	0.85
1:F:9:DA:OP2	3:H:213:ARG:NE	2.13	0.82
1:D:12:DT:OP2	5:D:101:HOH:O	2.01	0.77
1:D:9:DA:H2''	1:D:10:DA:H5''	1.70	0.74

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:11:DA:O3'	3:B:240:LYS:NZ[5_445]	2.08	0.12

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	83/102 (81%)	81 (98%)	2 (2%)	0	100	100
3	B	81/102 (79%)	80 (99%)	1 (1%)	0	100	100
3	C	83/102 (81%)	81 (98%)	2 (2%)	0	100	100
3	H	83/102 (81%)	81 (98%)	2 (2%)	0	100	100
All	All	330/408 (81%)	323 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	78/91 (86%)	78 (100%)	0	100	100
3	B	77/91 (85%)	76 (99%)	1 (1%)	69	75
3	C	78/91 (86%)	78 (100%)	0	100	100
3	H	78/91 (86%)	77 (99%)	1 (1%)	69	75
All	All	311/364 (85%)	309 (99%)	2 (1%)	86	90

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	B	240	LYS
3	H	213	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	D	16/16 (100%)	-0.03	0 100 100	23, 31, 64, 68	0
1	F	16/16 (100%)	-0.10	0 100 100	18, 26, 59, 61	0
2	E	16/16 (100%)	-0.08	0 100 100	17, 31, 59, 65	0
2	G	16/16 (100%)	-0.07	0 100 100	19, 26, 57, 67	0
3	A	85/102 (83%)	0.09	3 (3%) 44 50	12, 23, 57, 90	0
3	B	83/102 (81%)	0.59	10 (12%) 4 5	13, 27, 107, 147	0
3	C	85/102 (83%)	0.60	11 (12%) 3 4	14, 25, 118, 134	0
3	H	85/102 (83%)	0.15	3 (3%) 44 50	13, 24, 66, 92	0
All	All	402/472 (85%)	0.29	27 (6%) 17 22	12, 27, 90, 147	0

The worst 5 of 27 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	B	234	SER	10.7
3	B	236	ASP	10.5
3	C	235	PRO	10.1
3	C	236	ASP	8.4
3	C	252	ASN	7.8

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no monosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MG	A	301	1/1	0.99	0.17	10,10,10,10	0
4	MG	B	301	1/1	0.99	0.18	7,7,7,7	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around MG A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around MG B 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.