



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 14, 2020 – 07:21 am BST

PDB ID : 1VPH
Title : CRYSTAL STRUCTURE OF a YbjQ-like protein of unknown function (SSO2532) FROM SULFOLOBUS SOLFATARICUS P2 AT 1.76 Å RESOLUTION
Authors : Joint Center for Structural Genomics (JCSG)
Deposited on : 2004-11-03
Resolution : 1.76 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	149	<div> <div>5%</div> <div> <div></div> <div>84%</div> <div>8%</div> <div>7%</div> </div> </div>
1	B	149	<div> <div>4%</div> <div> <div></div> <div>87%</div> <div>5%</div> <div>8%</div> </div> </div>
1	C	149	<div> <div>7%</div> <div> <div></div> <div>86%</div> <div>11%</div> <div></div> </div> </div>
1	D	149	<div> <div>5%</div> <div> <div></div> <div>89%</div> <div></div> <div>7%</div> </div> </div>
1	E	149	<div> <div>7%</div> <div> <div></div> <div>87%</div> <div>5%</div> <div>7%</div> </div> </div>
1	F	149	<div> <div>9%</div> <div> <div></div> <div>85%</div> <div>10%</div> <div></div> </div> </div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7049 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called hypothetical protein SSO2532.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	138	Total	C	N	O	S	0	2	0
			1102	703	191	205	3			
1	B	137	Total	C	N	O	S	0	3	0
			1093	699	183	208	3			
1	C	144	Total	C	N	O	S	0	2	0
			1151	731	201	216	3			
1	D	138	Total	C	N	O	S	0	2	0
			1108	704	190	211	3			
1	E	138	Total	C	N	O	S	0	1	0
			1098	699	190	206	3			
1	F	143	Total	C	N	O	S	0	3	0
			1147	730	204	210	3			

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	MET	-	LEADER SEQUENCE	UNP Q97VS8
A	-10	GLY	-	LEADER SEQUENCE	UNP Q97VS8
A	-9	SER	-	LEADER SEQUENCE	UNP Q97VS8
A	-8	ASP	-	LEADER SEQUENCE	UNP Q97VS8
A	-7	LYS	-	LEADER SEQUENCE	UNP Q97VS8
A	-6	ILE	-	LEADER SEQUENCE	UNP Q97VS8
A	-5	HIS	-	LEADER SEQUENCE	UNP Q97VS8
A	-4	HIS	-	LEADER SEQUENCE	UNP Q97VS8
A	-3	HIS	-	LEADER SEQUENCE	UNP Q97VS8
A	-2	HIS	-	LEADER SEQUENCE	UNP Q97VS8
A	-1	HIS	-	LEADER SEQUENCE	UNP Q97VS8
A	0	HIS	-	LEADER SEQUENCE	UNP Q97VS8
B	-11	MET	-	LEADER SEQUENCE	UNP Q97VS8
B	-10	GLY	-	LEADER SEQUENCE	UNP Q97VS8
B	-9	SER	-	LEADER SEQUENCE	UNP Q97VS8
B	-8	ASP	-	LEADER SEQUENCE	UNP Q97VS8
B	-7	LYS	-	LEADER SEQUENCE	UNP Q97VS8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-6	ILE	-	LEADER SEQUENCE	UNP Q97VS8
B	-5	HIS	-	LEADER SEQUENCE	UNP Q97VS8
B	-4	HIS	-	LEADER SEQUENCE	UNP Q97VS8
B	-3	HIS	-	LEADER SEQUENCE	UNP Q97VS8
B	-2	HIS	-	LEADER SEQUENCE	UNP Q97VS8
B	-1	HIS	-	LEADER SEQUENCE	UNP Q97VS8
B	0	HIS	-	LEADER SEQUENCE	UNP Q97VS8
C	-11	MET	-	LEADER SEQUENCE	UNP Q97VS8
C	-10	GLY	-	LEADER SEQUENCE	UNP Q97VS8
C	-9	SER	-	LEADER SEQUENCE	UNP Q97VS8
C	-8	ASP	-	LEADER SEQUENCE	UNP Q97VS8
C	-7	LYS	-	LEADER SEQUENCE	UNP Q97VS8
C	-6	ILE	-	LEADER SEQUENCE	UNP Q97VS8
C	-5	HIS	-	LEADER SEQUENCE	UNP Q97VS8
C	-4	HIS	-	LEADER SEQUENCE	UNP Q97VS8
C	-3	HIS	-	LEADER SEQUENCE	UNP Q97VS8
C	-2	HIS	-	LEADER SEQUENCE	UNP Q97VS8
C	-1	HIS	-	LEADER SEQUENCE	UNP Q97VS8
C	0	HIS	-	LEADER SEQUENCE	UNP Q97VS8
D	-11	MET	-	LEADER SEQUENCE	UNP Q97VS8
D	-10	GLY	-	LEADER SEQUENCE	UNP Q97VS8
D	-9	SER	-	LEADER SEQUENCE	UNP Q97VS8
D	-8	ASP	-	LEADER SEQUENCE	UNP Q97VS8
D	-7	LYS	-	LEADER SEQUENCE	UNP Q97VS8
D	-6	ILE	-	LEADER SEQUENCE	UNP Q97VS8
D	-5	HIS	-	LEADER SEQUENCE	UNP Q97VS8
D	-4	HIS	-	LEADER SEQUENCE	UNP Q97VS8
D	-3	HIS	-	LEADER SEQUENCE	UNP Q97VS8
D	-2	HIS	-	LEADER SEQUENCE	UNP Q97VS8
D	-1	HIS	-	LEADER SEQUENCE	UNP Q97VS8
D	0	HIS	-	LEADER SEQUENCE	UNP Q97VS8
E	-11	MET	-	LEADER SEQUENCE	UNP Q97VS8
E	-10	GLY	-	LEADER SEQUENCE	UNP Q97VS8
E	-9	SER	-	LEADER SEQUENCE	UNP Q97VS8
E	-8	ASP	-	LEADER SEQUENCE	UNP Q97VS8
E	-7	LYS	-	LEADER SEQUENCE	UNP Q97VS8
E	-6	ILE	-	LEADER SEQUENCE	UNP Q97VS8
E	-5	HIS	-	LEADER SEQUENCE	UNP Q97VS8
E	-4	HIS	-	LEADER SEQUENCE	UNP Q97VS8
E	-3	HIS	-	LEADER SEQUENCE	UNP Q97VS8
E	-2	HIS	-	LEADER SEQUENCE	UNP Q97VS8
E	-1	HIS	-	LEADER SEQUENCE	UNP Q97VS8

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	0	HIS	-	LEADER SEQUENCE	UNP Q97VS8
F	-11	MET	-	LEADER SEQUENCE	UNP Q97VS8
F	-10	GLY	-	LEADER SEQUENCE	UNP Q97VS8
F	-9	SER	-	LEADER SEQUENCE	UNP Q97VS8
F	-8	ASP	-	LEADER SEQUENCE	UNP Q97VS8
F	-7	LYS	-	LEADER SEQUENCE	UNP Q97VS8
F	-6	ILE	-	LEADER SEQUENCE	UNP Q97VS8
F	-5	HIS	-	LEADER SEQUENCE	UNP Q97VS8
F	-4	HIS	-	LEADER SEQUENCE	UNP Q97VS8
F	-3	HIS	-	LEADER SEQUENCE	UNP Q97VS8
F	-2	HIS	-	LEADER SEQUENCE	UNP Q97VS8
F	-1	HIS	-	LEADER SEQUENCE	UNP Q97VS8
F	0	HIS	-	LEADER SEQUENCE	UNP Q97VS8

- Molecule 2 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total K 1 1	0	0
2	D	1	Total K 1 1	0	0
2	C	1	Total K 1 1	0	0
2	F	1	Total K 1 1	0	0

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O P 5 4 1	0	0
3	A	1	Total O P 5 4 1	0	0
3	B	1	Total O P 5 4 1	0	0
3	C	1	Total O P 5 4 1	0	0
3	C	1	Total O P 5 4 1	0	0
3	D	1	Total O P 5 4 1	0	0
3	D	1	Total O P 5 4 1	0	0
3	D	1	Total O P 5 4 1	0	0
3	E	1	Total O P 5 4 1	0	0
3	F	1	Total O P 5 4 1	0	0
3	F	1	Total O P 5 4 1	0	0

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	1	Total Cl 1 1	0	0


- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	54	Total	O	0	0
			54	54		
6	B	60	Total	O	0	0
			60	60		
6	C	48	Total	O	0	0
			48	48		
6	D	48	Total	O	0	0
			48	48		
6	E	44	Total	O	0	0
			44	44		
6	F	32	Total	O	0	0
			32	32		



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	79.86 Å 92.18 Å 114.13 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	16.22 – 1.76 16.22 – 1.76	Depositor EDS
% Data completeness (in resolution range)	90.7 (16.22-1.76) 90.8 (16.22-1.76)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.07 (at 1.76 Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.189 , 0.222 0.196 , 0.229	Depositor DCC
R_{free} test set	3789 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	22.0	Xtriage
Anisotropy	0.606	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 55.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7049	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 70.73 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.9676e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, K, EDO, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.81	0/1132	0.80	1/1533 (0.1%)
1	B	0.80	0/1126	0.77	0/1525
1	C	0.83	0/1186	0.84	0/1608
1	D	0.84	0/1138	0.84	0/1539
1	E	0.84	0/1124	0.82	1/1523 (0.1%)
1	F	0.76	0/1191	0.81	0/1614
All	All	0.81	0/6897	0.81	2/9342 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	110	ARG	NE-CZ-NH1	5.46	123.03	120.30
1	A	1	MET	N-CA-C	-5.03	97.43	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1102	0	1077	13	0
1	B	1093	0	1058	8	0
1	C	1151	0	1088	9	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	1108	0	1081	11	0
1	E	1098	0	1068	7	0
1	F	1147	0	1097	14	0
2	A	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	F	1	0	0	0	0
3	A	10	0	0	0	0
3	B	5	0	0	0	0
3	C	10	0	0	0	0
3	D	15	0	0	1	0
3	E	5	0	0	0	0
3	F	10	0	0	1	0
4	C	1	0	0	0	0
5	D	4	0	6	1	0
6	A	54	0	0	1	0
6	B	60	0	0	0	0
6	C	48	0	0	0	0
6	D	48	0	0	0	0
6	E	44	0	0	1	0
6	F	32	0	0	0	0
All	All	7049	0	6475	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 49 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:28:ILE:HG22	1:A:134:ILE:HD11	1.51	0.93
1:D:1:MET:CE	1:E:137:GLU:HB3	1.99	0.92
1:A:10[A]:VAL:HG21	1:A:118:LEU:HD22	1.64	0.80
1:F:13:ALA:N	1:F:17:GLU:OE2	2.26	0.66
1:D:1:MET:HE1	1:E:137:GLU:HB3	1.76	0.66

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	138/149 (93%)	133 (96%)	4 (3%)	1 (1%)	22	8
1	B	138/149 (93%)	133 (96%)	5 (4%)	0	100	100
1	C	144/149 (97%)	140 (97%)	4 (3%)	0	100	100
1	D	138/149 (93%)	133 (96%)	5 (4%)	0	100	100
1	E	137/149 (92%)	132 (96%)	5 (4%)	0	100	100
1	F	144/149 (97%)	140 (97%)	4 (3%)	0	100	100
All	All	839/894 (94%)	811 (97%)	27 (3%)	1 (0%)	51	33

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1	MET

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	117/130 (90%)	116 (99%)	1 (1%)	78	67
1	B	115/130 (88%)	114 (99%)	1 (1%)	78	67
1	C	120/130 (92%)	118 (98%)	2 (2%)	60	42
1	D	119/130 (92%)	116 (98%)	3 (2%)	47	25
1	E	116/130 (89%)	111 (96%)	5 (4%)	29	9
1	F	121/130 (93%)	117 (97%)	4 (3%)	38	15

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	708/780 (91%)	692 (98%)	16 (2%)	55 28

5 of 16 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	1	MET
1	E	3[A]	VAL
1	F	1	MET
1	D	98[B]	GLU
1	F	26	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	F	15	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 17 ligands modelled in this entry, 5 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PO4	D	141	-	4,4,4	0.96	0	6,6,6	1.00	0
3	PO4	F	139	-	4,4,4	1.24	0	6,6,6	1.55	1 (16%)
3	PO4	D	140	-	4,4,4	1.26	1 (25%)	6,6,6	1.05	0
3	PO4	D	139	-	4,4,4	1.91	1 (25%)	6,6,6	1.49	1 (16%)
3	PO4	A	140	-	4,4,4	0.83	0	6,6,6	0.76	0
3	PO4	C	140	-	4,4,4	0.56	0	6,6,6	0.82	0
3	PO4	F	140	-	4,4,4	1.09	0	6,6,6	0.93	0
3	PO4	E	138	-	4,4,4	1.50	1 (25%)	6,6,6	1.78	2 (33%)
3	PO4	B	138	-	4,4,4	1.01	0	6,6,6	0.89	0
5	EDO	D	142	-	3,3,3	0.40	0	2,2,2	0.22	0
3	PO4	A	139	-	4,4,4	2.13	1 (25%)	6,6,6	1.07	0
3	PO4	C	139	-	4,4,4	0.98	0	6,6,6	1.35	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	D	142	-	-	1/1/1/1	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	139	PO4	P-O2	-3.25	1.44	1.54
3	D	139	PO4	P-O4	-2.74	1.46	1.54
3	E	138	PO4	P-O3	-2.29	1.47	1.54
3	D	140	PO4	P-O2	-2.09	1.48	1.54

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	138	PO4	O3-P-O2	3.51	119.25	107.97
3	F	139	PO4	O3-P-O2	3.18	118.17	107.97
3	E	138	PO4	O3-P-O1	-2.47	101.86	110.89
3	C	139	PO4	O4-P-O2	2.44	115.81	107.97
3	D	139	PO4	O4-P-O3	2.35	115.51	107.97

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	D	142	EDO	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	139	PO4	1	0
3	D	140	PO4	1	0
5	D	142	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	138/149 (92%)	0.40	8 (5%) 23 28	18, 22, 33, 54	0
1	B	137/149 (91%)	0.18	6 (4%) 34 40	17, 22, 33, 37	0
1	C	144/149 (96%)	0.59	11 (7%) 13 18	18, 22, 32, 38	0
1	D	138/149 (92%)	0.36	7 (5%) 28 34	18, 22, 33, 44	0
1	E	138/149 (92%)	0.41	11 (7%) 12 16	17, 22, 32, 45	0
1	F	143/149 (95%)	0.64	13 (9%) 9 11	18, 22, 32, 37	0
All	All	838/894 (93%)	0.43	56 (6%) 17 23	17, 22, 33, 54	0

The worst 5 of 56 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	82	ILE	4.3
1	C	13	ALA	4.3
1	B	126	SER	4.0
1	E	126	SER	4.0
1	A	113	TRP	3.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PO4	A	140	5/5	0.91	0.24	44,46,52,57	0
3	PO4	D	141	5/5	0.93	0.20	41,41,56,58	0
5	EDO	D	142	4/4	0.94	0.13	23,31,34,34	0
3	PO4	C	140	5/5	0.96	0.23	36,37,44,46	0
3	PO4	D	140	5/5	0.96	0.18	28,28,33,38	0
3	PO4	F	139	5/5	0.96	0.11	33,34,35,42	0
3	PO4	D	139	5/5	0.97	0.07	19,22,30,33	0
3	PO4	F	140	5/5	0.97	0.17	30,30,33,44	0
2	K	A	138	1/1	0.97	0.07	22,22,22,22	0
3	PO4	E	138	5/5	0.97	0.10	21,23,27,30	0
3	PO4	B	138	5/5	0.97	0.09	18,26,27,35	0
4	CL	C	141	1/1	0.97	0.06	48,48,48,48	0
2	K	F	138	1/1	0.98	0.07	28,28,28,28	0
3	PO4	C	139	5/5	0.98	0.08	24,30,41,42	0
2	K	D	138	1/1	0.98	0.04	24,24,24,24	0
2	K	C	138	1/1	0.99	0.10	27,27,27,27	0
3	PO4	A	139	5/5	0.99	0.05	20,23,29,31	0

6.5 Other polymers ⓘ

There are no such residues in this entry.