



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 12, 2022 – 04:44 PM JST

PDB ID : 7VPG
Title : Crystal structure of the C-terminal tail of SARS-CoV-1 Orf6 complex with human nucleoporin pair Rae1-Nup98
Authors : Li, T.; Guo, H.; Yang, T.; Wen, Y.; Ji, X.
Deposited on : 2021-10-17
Resolution : 2.49 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.25
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.25

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	378	<div><div></div><div><div>78%</div><div>11%</div><div>•</div><div>10%</div></div></div>
1	C	378	<div>%<div><div></div><div>77%</div><div>11%</div><div>•</div><div>11%</div></div></div>
1	E	378	<div>%<div><div></div><div>78%</div><div>10%</div><div>•</div><div>11%</div></div></div>
1	G	378	<div>%<div><div></div><div>74%</div><div>14%</div><div>•</div><div>10%</div></div></div>
2	B	67	<div><div>3%</div><div></div><div>66%</div><div>12%</div><div>22%</div></div>
2	D	67	<div><div></div><div>63%</div><div>9%</div><div>28%</div></div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	67	<div><div>%</div><div><div></div><div></div><div></div></div><div>70%10%19%</div></div>
2	H	67	<div><div>4%</div><div><div></div><div></div><div></div></div><div>57%19%24%</div></div>
3	I	22	<div><div>14%</div><div><div></div><div></div><div></div></div><div>36%9%55%</div></div>
3	J	22	<div><div>14%</div><div><div></div><div></div><div></div></div><div>27%18%55%</div></div>
3	K	22	<div><div>5%</div><div><div></div><div></div><div></div></div><div>23%23%55%</div></div>
3	X	22	<div><div>9%</div><div><div></div><div></div><div></div></div><div>41%18%41%</div></div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 13136 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called mRNA export factor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	341	Total	C	N	O	S	0	1	0
			2704	1706	476	504	18			
1	C	338	Total	C	N	O	S	0	0	0
			2674	1689	469	498	18			
1	E	338	Total	C	N	O	S	0	0	0
			2674	1689	468	499	18			
1	G	339	Total	C	N	O	S	0	1	0
			2686	1697	470	501	18			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	369	HIS	-	expression tag	UNP P78406
A	370	HIS	-	expression tag	UNP P78406
A	371	HIS	-	expression tag	UNP P78406
A	372	HIS	-	expression tag	UNP P78406
A	373	HIS	-	expression tag	UNP P78406
A	374	HIS	-	expression tag	UNP P78406
A	375	HIS	-	expression tag	UNP P78406
A	376	HIS	-	expression tag	UNP P78406
A	377	HIS	-	expression tag	UNP P78406
A	378	HIS	-	expression tag	UNP P78406
C	369	HIS	-	expression tag	UNP P78406
C	370	HIS	-	expression tag	UNP P78406
C	371	HIS	-	expression tag	UNP P78406
C	372	HIS	-	expression tag	UNP P78406
C	373	HIS	-	expression tag	UNP P78406
C	374	HIS	-	expression tag	UNP P78406
C	375	HIS	-	expression tag	UNP P78406
C	376	HIS	-	expression tag	UNP P78406
C	377	HIS	-	expression tag	UNP P78406
C	378	HIS	-	expression tag	UNP P78406
E	369	HIS	-	expression tag	UNP P78406

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	370	HIS	-	expression tag	UNP P78406
E	371	HIS	-	expression tag	UNP P78406
E	372	HIS	-	expression tag	UNP P78406
E	373	HIS	-	expression tag	UNP P78406
E	374	HIS	-	expression tag	UNP P78406
E	375	HIS	-	expression tag	UNP P78406
E	376	HIS	-	expression tag	UNP P78406
E	377	HIS	-	expression tag	UNP P78406
E	378	HIS	-	expression tag	UNP P78406
G	369	HIS	-	expression tag	UNP P78406
G	370	HIS	-	expression tag	UNP P78406
G	371	HIS	-	expression tag	UNP P78406
G	372	HIS	-	expression tag	UNP P78406
G	373	HIS	-	expression tag	UNP P78406
G	374	HIS	-	expression tag	UNP P78406
G	375	HIS	-	expression tag	UNP P78406
G	376	HIS	-	expression tag	UNP P78406
G	377	HIS	-	expression tag	UNP P78406
G	378	HIS	-	expression tag	UNP P78406

- Molecule 2 is a protein called Isoform 3 of Nuclear pore complex protein Nup98-Nup96.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	52	Total	C	N	O	S	0	0	0
			412	252	70	87	3			
2	D	48	Total	C	N	O	S	0	0	0
			384	235	66	81	2			
2	F	54	Total	C	N	O	S	0	0	0
			423	259	72	89	3			
2	H	51	Total	C	N	O	S	0	0	0
			405	247	69	86	3			

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	147	MET	-	initiating methionine	UNP P52948
B	148	HIS	-	expression tag	UNP P52948
B	149	HIS	-	expression tag	UNP P52948
B	150	HIS	-	expression tag	UNP P52948
B	151	HIS	-	expression tag	UNP P52948
B	152	HIS	-	expression tag	UNP P52948
B	153	HIS	-	expression tag	UNP P52948

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	154	HIS	-	expression tag	UNP P52948
B	155	HIS	-	expression tag	UNP P52948
B	156	HIS	-	expression tag	UNP P52948
B	157	HIS	-	expression tag	UNP P52948
D	147	MET	-	initiating methionine	UNP P52948
D	148	HIS	-	expression tag	UNP P52948
D	149	HIS	-	expression tag	UNP P52948
D	150	HIS	-	expression tag	UNP P52948
D	151	HIS	-	expression tag	UNP P52948
D	152	HIS	-	expression tag	UNP P52948
D	153	HIS	-	expression tag	UNP P52948
D	154	HIS	-	expression tag	UNP P52948
D	155	HIS	-	expression tag	UNP P52948
D	156	HIS	-	expression tag	UNP P52948
D	157	HIS	-	expression tag	UNP P52948
F	147	MET	-	initiating methionine	UNP P52948
F	148	HIS	-	expression tag	UNP P52948
F	149	HIS	-	expression tag	UNP P52948
F	150	HIS	-	expression tag	UNP P52948
F	151	HIS	-	expression tag	UNP P52948
F	152	HIS	-	expression tag	UNP P52948
F	153	HIS	-	expression tag	UNP P52948
F	154	HIS	-	expression tag	UNP P52948
F	155	HIS	-	expression tag	UNP P52948
F	156	HIS	-	expression tag	UNP P52948
F	157	HIS	-	expression tag	UNP P52948
H	147	MET	-	initiating methionine	UNP P52948
H	148	HIS	-	expression tag	UNP P52948
H	149	HIS	-	expression tag	UNP P52948
H	150	HIS	-	expression tag	UNP P52948
H	151	HIS	-	expression tag	UNP P52948
H	152	HIS	-	expression tag	UNP P52948
H	153	HIS	-	expression tag	UNP P52948
H	154	HIS	-	expression tag	UNP P52948
H	155	HIS	-	expression tag	UNP P52948
H	156	HIS	-	expression tag	UNP P52948
H	157	HIS	-	expression tag	UNP P52948

- Molecule 3 is a protein called ORF6 protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	X	13	Total	C	N	O	S	0	0	0
			109	66	13	29	1			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	I	10	Total	C	N	O	S	0	0	0
			86	52	10	23	1			
3	J	10	Total	C	N	O	S	0	0	0
			86	52	10	23	1			
3	K	10	Total	C	N	O	S	0	0	0
			86	52	10	23	1			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	116	Total	O	0	0
			116	116		
4	B	18	Total	O	0	0
			18	18		
4	C	76	Total	O	0	0
			76	76		
4	D	3	Total	O	0	0
			3	3		
4	E	91	Total	O	0	0
			91	91		
4	F	13	Total	O	0	0
			13	13		
4	G	80	Total	O	0	0
			80	80		
4	H	2	Total	O	0	0
			2	2		
4	X	3	Total	O	0	0
			3	3		
4	I	1	Total	O	0	0
			1	1		
4	J	4	Total	O	0	0
			4	4		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: mRNA export factor



- Molecule 1: mRNA export factor





- Molecule 1: mRNA export factor







● Molecule 3: ORF6 protein



● Molecule 3: ORF6 protein



● Molecule 3: ORF6 protein



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	149.61Å 103.30Å 134.94Å 90.00° 95.96° 90.00°	Depositor
Resolution (Å)	47.44 – 2.49 47.44 – 2.49	Depositor EDS
% Data completeness (in resolution range)	97.9 (47.44-2.49) 87.3 (47.44-2.49)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.59 (at 2.48Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.185 , 0.231 0.184 , 0.229	Depositor DCC
R_{free} test set	1963 reflections (2.79%)	wwPDB-VP
Wilson B-factor (Å ²)	34.5	Xtriage
Anisotropy	0.430	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 38.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	13136	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.53	1/2784 (0.0%)	0.75	5/3784 (0.1%)
1	C	0.42	0/2751	0.64	0/3739
1	E	0.44	0/2751	0.66	0/3739
1	G	0.42	0/2766	0.67	1/3760 (0.0%)
2	B	0.41	0/416	0.65	0/557
2	D	0.40	0/388	0.61	0/519
2	F	0.46	0/427	0.66	0/572
2	H	0.37	0/409	0.59	0/547
3	I	0.45	0/87	0.64	0/117
3	J	0.46	0/87	0.72	0/117
3	K	0.46	0/87	0.62	0/117
3	X	0.43	0/110	0.69	0/148
All	All	0.45	1/13063 (0.0%)	0.67	6/17716 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	E	0	2
1	G	0	2
All	All	0	6

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	350	LYS	CB-CG	5.62	1.67	1.52

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	350	LYS	CB-CG-CD	9.72	136.88	111.60
1	A	350	LYS	CA-CB-CG	8.01	131.03	113.40
1	G	348	GLN	CA-CB-CG	6.46	127.62	113.40
1	A	252	LYS	CD-CE-NZ	-6.17	97.50	111.70
1	A	252	LYS	N-CA-CB	-5.04	101.53	110.60

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	347	PRO	Peptide
1	A	348	GLN	Peptide
1	E	76	GLN	Sidechain,Peptide
1	G	347	PRO	Peptide
1	G	348	GLN	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2704	0	2601	24	0
1	C	2674	0	2568	24	0
1	E	2674	0	2566	21	1
1	G	2686	0	2581	41	1
2	B	412	0	406	9	0
2	D	384	0	376	5	0
2	F	423	0	418	4	0
2	H	405	0	397	8	0
3	I	86	0	65	0	0
3	J	86	0	65	1	0
3	K	86	0	65	2	0
3	X	109	0	87	4	0
4	A	116	0	0	1	0
4	B	18	0	0	1	0
4	C	76	0	0	0	0
4	D	3	0	0	1	0
4	E	91	0	0	2	0
4	F	13	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	G	80	0	0	3	0
4	H	2	0	0	0	0
4	I	1	0	0	0	0
4	J	4	0	0	0	0
4	X	3	0	0	0	0
All	All	13136	0	12195	131	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 131 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:348:GLN:OE1	1:A:351:ASN:ND2	2.10	0.84
1:E:211:LEU:HD13	1:E:235:SER:HB3	1.66	0.78
1:E:140:ASN:ND2	4:E:401:HOH:O	2.17	0.75
1:A:193:ILE:HG12	1:A:206[A]:ARG:HD3	1.68	0.73
2:H:168:THR:HG22	2:H:185:LYS:HD3	1.70	0.73

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:63:ALA:O	1:G:348:GLN:NE2[2_556]	2.11	0.09

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	338/378 (89%)	329 (97%)	9 (3%)	0	100	100
1	C	334/378 (88%)	326 (98%)	8 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	334/378 (88%)	328 (98%)	6 (2%)	0	100	100
1	G	336/378 (89%)	328 (98%)	7 (2%)	1 (0%)	41	61
2	B	48/67 (72%)	47 (98%)	1 (2%)	0	100	100
2	D	44/67 (66%)	43 (98%)	1 (2%)	0	100	100
2	F	50/67 (75%)	49 (98%)	1 (2%)	0	100	100
2	H	47/67 (70%)	46 (98%)	1 (2%)	0	100	100
3	I	8/22 (36%)	8 (100%)	0	0	100	100
3	J	8/22 (36%)	8 (100%)	0	0	100	100
3	K	8/22 (36%)	8 (100%)	0	0	100	100
3	X	11/22 (50%)	11 (100%)	0	0	100	100
All	All	1566/1868 (84%)	1531 (98%)	34 (2%)	1 (0%)	51	73

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	G	348	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	298/327 (91%)	288 (97%)	10 (3%)	37	63
1	C	294/327 (90%)	280 (95%)	14 (5%)	25	48
1	E	294/327 (90%)	283 (96%)	11 (4%)	34	60
1	G	296/327 (90%)	284 (96%)	12 (4%)	30	55
2	B	48/61 (79%)	48 (100%)	0	100	100
2	D	44/61 (72%)	44 (100%)	0	100	100
2	F	49/61 (80%)	47 (96%)	2 (4%)	30	55
2	H	47/61 (77%)	46 (98%)	1 (2%)	53	78

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	I	10/22 (46%)	8 (80%)	2 (20%)	1	2
3	J	10/22 (46%)	8 (80%)	2 (20%)	1	2
3	K	10/22 (46%)	7 (70%)	3 (30%)	0	0
3	X	13/22 (59%)	12 (92%)	1 (8%)	13	25
All	All	1413/1640 (86%)	1355 (96%)	58 (4%)	30	55

5 of 58 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	205	ARG
3	K	53	ASP
2	F	158	THR
3	J	61	ASP
2	H	171	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	366	ASN
2	H	181	ASN
1	E	25	ASN
1	E	263	ASN
1	E	285	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	341/378 (90%)	-0.37	1 (0%) 94 94	24, 35, 54, 84	0
1	C	338/378 (89%)	-0.06	5 (1%) 73 75	30, 45, 65, 90	0
1	E	338/378 (89%)	-0.17	4 (1%) 79 80	27, 40, 61, 95	0
1	G	339/378 (89%)	-0.22	5 (1%) 73 75	28, 45, 67, 98	0
2	B	52/67 (77%)	0.02	2 (3%) 40 43	30, 44, 77, 87	0
2	D	48/67 (71%)	-0.01	0 100 100	38, 53, 77, 82	0
2	F	54/67 (80%)	-0.08	1 (1%) 66 69	33, 45, 69, 78	0
2	H	51/67 (76%)	0.27	3 (5%) 22 23	41, 59, 98, 105	0
3	I	10/22 (45%)	0.96	3 (30%) 0 0	49, 63, 83, 90	0
3	J	10/22 (45%)	0.93	3 (30%) 0 0	44, 63, 84, 86	0
3	K	10/22 (45%)	0.71	1 (10%) 7 6	49, 63, 85, 85	0
3	X	13/22 (59%)	0.87	2 (15%) 2 1	41, 55, 89, 92	0
All	All	1604/1868 (85%)	-0.15	30 (1%) 66 69	24, 43, 68, 105	0

The worst 5 of 30 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	X	50	SER	5.6
1	C	366	ASN	4.0
1	G	263	ASN	3.6
2	H	172	THR	3.5
1	C	263	ASN	3.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.