



Full wwPDB EM Validation Report ⓘ

Nov 13, 2022 – 02:34 AM EST

PDB ID : 6VQC
EMDB ID : EMD-21348
Title : Mammalian V-ATPase from rat brain membrane-embedded Vo region rotational state 1 (from focused refinement)
Authors : Abbas, Y.M.; Rubinstein, J.L.
Deposited on : 2020-02-04
Resolution : 3.80 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.













Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	247	
2	L	119	
3	a	838	
4	b	205	
5	c	463	
6	d	351	
7	e	81	
8	f	98	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
9	g	155	
9	h	155	
9	i	155	
9	j	155	
9	k	155	
9	l	155	
9	m	155	
9	n	155	
9	o	155	
10	p	350	

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 20991 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATPase H⁺-transporting V1 subunit D.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	H	104	Total	C	N	O	S	0	0
			803	509	138	153	3		

- Molecule 2 is a protein called V-type proton ATPase subunit F.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	L	110	Total	C	N	O	S	0	0
			875	553	157	163	2		

- Molecule 3 is a protein called V-type proton ATPase 116 kDa subunit a isoform 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	a	441	Total	C	N	O	S	0	0
			3495	2337	556	579	23		

- Molecule 4 is a protein called ATPase, H⁺ transporting, V0 subunit B (Predicted), isoform CRA_a.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	b	203	Total	C	N	O	S	0	0
			1503	996	237	259	11		

- Molecule 5 is a protein called V-type proton ATPase subunit S1.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	c	41	Total	C	N	O	S	0	0
			337	228	51	54	4		

- Molecule 6 is a protein called V-type proton ATPase subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	d	350	Total	C	N	O	S	0	0
			2833	1829	460	530	14		

- Molecule 7 is a protein called V-type proton ATPase subunit e 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	e	78	Total	C	N	O	S	0	0
			621	428	98	92	3		

- Molecule 8 is a protein called Ribonuclease K.


Mol	Chain	Residues	Atoms				AltConf	Trace
8	f	84	Total	C	N	O	0	0
			480	309	85	86		

- Molecule 9 is a protein called V-type proton ATPase 16 kDa proteolipid subunit.


Mol	Chain	Residues	Atoms					AltConf	Trace
9	g	150	Total	C	N	O	S	0	0
			1068	699	171	190	8		
9	h	150	Total	C	N	O	S	0	0
			1068	699	171	190	8		
9	i	150	Total	C	N	O	S	0	0
			1068	699	171	190	8		
9	j	150	Total	C	N	O	S	0	0
			1068	699	171	190	8		
9	k	150	Total	C	N	O	S	0	0
			1068	699	171	190	8		
9	l	150	Total	C	N	O	S	0	0
			1068	699	171	190	8		
9	m	150	Total	C	N	O	S	0	0
			1068	699	171	190	8		
9	n	150	Total	C	N	O	S	0	0
			1068	699	171	190	8		
9	o	150	Total	C	N	O	S	0	0
			1068	699	171	190	8		

- Molecule 10 is a protein called Renin receptor.


Mol	Chain	Residues	Atoms					AltConf	Trace
10	p	52	Total	C	N	O	S	0	0
			432	290	63	76	3		




- Molecule 7: V-type proton ATPase subunit e 2




- Molecule 8: Ribonuclease K




- Molecule 9: V-type proton ATPase 16 kDa proteolipid subunit



- Molecule 9: V-type proton ATPase 16 kDa proteolipid subunit




- Molecule 9: V-type proton ATPase 16 kDa proteolipid subunit




- Molecule 9: V-type proton ATPase 16 kDa proteolipid subunit







- Molecule 9: V-type proton ATPase 16 kDa proteolipid subunit




- Molecule 9: V-type proton ATPase 16 kDa proteolipid subunit




- Molecule 9: V-type proton ATPase 16 kDa proteolipid subunit





- Molecule 9: V-type proton ATPase 16 kDa proteolipid subunit



- Molecule 9: V-type proton ATPase 16 kDa proteolipid subunit



- Molecule 10: Renin receptor



<div>◆</div> <div>Y335</div> <div>D336</div> <div>S337</div> <div>I338</div> <div>I339</div> <div>Y340</div> <div>R341</div> <div>M342</div> <div>T343</div>	LEU	ASP	GLU
	VAL	LEU	THR
	ASP	LEU	PRO
	ALA	PHE	VAL
	LEU	LEU	VAL
	GLN	SER	LEU
	LYS	GLU	GLN
	PHE	LEU	LEU
	ALA	GLN	ALA
	ASP	VAL	PRO
<div>◆</div> <div>ASN</div> <div>GLN</div> <div>LYS</div> <div>ILE</div> <div>ARG</div> <div>MET</div> <div>ASP</div>	ASP	HIS	GLU
	MET	ASP	GLU
	THR	ILE	ASP
	SER	SER	VAL
	LEU	SER	THR
	THR	SER	MET
	GLY	LEU	VAL
	GLY	LEU	VAL
	ASN	SER	GLY
	ALA	ARG	LYS
<div>◆</div> <div>VAL</div> <div>GLU</div> <div>VAL</div> <div>THR</div> <div>THR</div> <div>VAL</div> <div>VAL</div> <div>THR</div> <div>VAL</div>	VAL	HIS	ALA
	VAL	SER	ASN
	GLU	SER	ASN
	LEU	PRO	ASN
	VAL	ASP	VAL
	VAL	THR	THR
	THR	LEU	GLN
	THR	SER	LEU
	VAL	LEU	ARG
	LYS	GLU	ASN
<div>◆</div> <div>SER</div> <div>P292</div> <div>N300</div> <div>G316</div> <div>D332</div> <div>P333</div> <div>G334</div>	SER	ALA	ASN
	THR	ALA	ASN
	GLN	ARG	ASN
	GLU	THR	ASN
	ASN	GLY	LEU
	THR	GLU	SER
	GLN	ASP	PRO
	GLN	SER	LEU
	SER	GLN	ASN
	P292	PHE	LEU

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	90648	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	43	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	FEI FALCON III (4k x 4k)	Depositor
Maximum map value	2.773	Depositor
Minimum map value	-1.050	Depositor
Average map value	-0.002	Depositor
Map value standard deviation	0.069	Depositor
Recommended contour level	0.5	Depositor
Map size (\AA)	360.4, 360.4, 360.4	wwPDB
Map dimensions	340, 340, 340	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.06, 1.06, 1.06	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	H	0.84	0/812	0.64	0/1091
2	L	0.80	0/889	0.74	0/1199
3	a	0.52	0/3603	0.49	0/4888
4	b	0.51	0/1537	0.49	0/2088
5	c	0.48	0/347	0.54	0/466
6	d	0.73	0/2899	0.53	0/3927
7	e	0.47	0/645	0.44	0/887
8	f	0.40	0/490	0.50	0/676
9	g	0.50	0/1083	0.56	0/1466
9	h	0.47	0/1083	0.52	0/1466
9	i	0.49	0/1083	0.52	0/1466
9	j	0.47	0/1083	0.49	0/1466
9	k	0.49	0/1083	0.53	0/1466
9	l	0.48	0/1083	0.51	0/1466
9	m	0.49	0/1083	0.55	0/1466
9	n	0.47	0/1083	0.51	0/1466
9	o	0.49	0/1083	0.51	0/1466
10	p	0.49	0/445	0.60	0/609
All	All	0.56	0/21414	0.53	0/29025

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	803	0	828	4	0
2	L	875	0	883	12	0
3	a	3495	0	3434	0	0
4	b	1503	0	1551	0	0
5	c	337	0	338	0	0
6	d	2833	0	2770	0	0
7	e	621	0	639	0	0
8	f	480	0	288	0	0
9	g	1068	0	1136	0	0
9	h	1068	0	1136	0	0
9	i	1068	0	1136	0	0
9	j	1068	0	1136	0	0
9	k	1068	0	1136	0	0
9	l	1068	0	1136	0	0
9	m	1068	0	1136	0	0
9	n	1068	0	1136	0	0
9	o	1068	0	1136	0	0
10	p	432	0	428	0	0
All	All	20991	0	21383	13	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:58:ARG:HG2	2:L:60:ASP:H	1.65	0.61
2:L:85:ILE:N	2:L:86:PRO:CD	2.67	0.58
2:L:85:ILE:N	2:L:86:PRO:HD2	2.23	0.54
1:H:68:SER:OG	1:H:137:ASN:ND2	2.41	0.53
1:H:55:LYS:NZ	2:L:99:ASP:OD2	2.32	0.53
2:L:112:MET:O	2:L:113:PHE:C	2.55	0.43
2:L:60:ASP:N	2:L:60:ASP:OD1	2.52	0.42
1:H:148:LEU:HG	2:L:89:LEU:HD22	2.01	0.42
2:L:9:ALA:CB	2:L:61:ILE:HG21	2.51	0.41
2:L:85:ILE:HB	2:L:86:PRO:HD3	2.01	0.41
1:H:148:LEU:HD23	2:L:89:LEU:HD13	2.02	0.41
2:L:107:ARG:O	2:L:108:ARG:C	2.58	0.40
2:L:9:ALA:HB2	2:L:61:ILE:HG21	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	102/247 (41%)	97 (95%)	5 (5%)	0	100	100
2	L	108/119 (91%)	101 (94%)	5 (5%)	2 (2%)	8	42
3	a	435/838 (52%)	424 (98%)	11 (2%)	0	100	100
4	b	201/205 (98%)	197 (98%)	4 (2%)	0	100	100
5	c	39/463 (8%)	39 (100%)	0	0	100	100
6	d	348/351 (99%)	328 (94%)	19 (6%)	1 (0%)	41	74
7	e	76/81 (94%)	75 (99%)	1 (1%)	0	100	100
8	f	80/98 (82%)	77 (96%)	3 (4%)	0	100	100
9	g	148/155 (96%)	146 (99%)	2 (1%)	0	100	100
9	h	148/155 (96%)	145 (98%)	3 (2%)	0	100	100
9	i	148/155 (96%)	145 (98%)	3 (2%)	0	100	100
9	j	148/155 (96%)	146 (99%)	2 (1%)	0	100	100
9	k	148/155 (96%)	146 (99%)	2 (1%)	0	100	100
9	l	148/155 (96%)	145 (98%)	3 (2%)	0	100	100
9	m	148/155 (96%)	146 (99%)	2 (1%)	0	100	100
9	n	148/155 (96%)	145 (98%)	2 (1%)	1 (1%)	22	60
9	o	148/155 (96%)	143 (97%)	4 (3%)	1 (1%)	22	60
10	p	50/350 (14%)	47 (94%)	2 (4%)	1 (2%)	7	41
All	All	2771/4147 (67%)	2692 (97%)	73 (3%)	6 (0%)	50	79

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	L	61	ILE
6	d	119	GLN
9	o	10	TYR
10	p	300	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	L	100	ALA
9	n	10	TYR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	84/212 (40%)	84 (100%)	0	100	100
2	L	94/100 (94%)	93 (99%)	1 (1%)	73	85
3	a	362/743 (49%)	362 (100%)	0	100	100
4	b	156/158 (99%)	154 (99%)	2 (1%)	69	82
5	c	36/395 (9%)	35 (97%)	1 (3%)	43	68
6	d	305/306 (100%)	304 (100%)	1 (0%)	92	96
7	e	64/68 (94%)	64 (100%)	0	100	100
8	f	14/83 (17%)	14 (100%)	0	100	100
9	g	109/113 (96%)	108 (99%)	1 (1%)	78	88
9	h	109/113 (96%)	108 (99%)	1 (1%)	78	88
9	i	109/113 (96%)	109 (100%)	0	100	100
9	j	109/113 (96%)	109 (100%)	0	100	100
9	k	109/113 (96%)	109 (100%)	0	100	100
9	l	109/113 (96%)	109 (100%)	0	100	100
9	m	109/113 (96%)	108 (99%)	1 (1%)	78	88
9	n	109/113 (96%)	106 (97%)	3 (3%)	43	68
9	o	109/113 (96%)	108 (99%)	1 (1%)	78	88
10	p	47/313 (15%)	47 (100%)	0	100	100
All	All	2143/3395 (63%)	2131 (99%)	12 (1%)	86	92

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	L	60	ASP
4	b	58	ILE
4	b	70	TYR
5	c	448	ARG
6	d	24	LYS
9	g	141	LEU
9	h	139	GLU
9	m	61	MET
9	n	139	GLU
9	n	141	LEU
9	n	143	LEU
9	o	139	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	H	137	ASN
3	a	751	GLN
6	d	297	HIS
6	d	340	HIS
9	m	123	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-21348. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y



Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 170



Y Index: 170




Z Index: 170


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 183



Y Index: 171




Z Index: 247


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views [i](#)


6.4.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.5. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.5 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 134 nm^3 ; this corresponds to an approximate mass of 121 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.263 Å⁻¹


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-21348 and PDB model 6VQC. Per-residue inclusion information can be found in section [3](#) on page [6](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.5 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.5).







































9.4 Atom inclusion ⓘ



At the recommended contour level, 83% of all backbone atoms, 63% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.5) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.6287	 0.3690
H	 0.6692	 0.3820
L	 0.6418	 0.3750
a	 0.6062	 0.3590
b	 0.6168	 0.3700
c	 0.5982	 0.3950
d	 0.6533	 0.3910
e	 0.6039	 0.3530
f	 0.6415	 0.3270
g	 0.6341	 0.3680
h	 0.6370	 0.3770
i	 0.6436	 0.3750
j	 0.6417	 0.3830
k	 0.6028	 0.3650
l	 0.6256	 0.3590
m	 0.6209	 0.3610
n	 0.6370	 0.3550
o	 0.6294	 0.3590
p	 0.6010	 0.3750

