



Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 10:13 pm BST

PDB ID : 5VUJ
Title : Structure of rat neuronal nitric oxide synthase heme domain in complex with
7-(((3-(Dimethylamino)benzyl)amino)methyl)quinolin-2-amine
Authors : Li, H.; Poulos, T.L.
Deposited on : 2017-05-19
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	422	<div> <div>4%</div> <div> <div></div> <div>83%</div> <div>13%</div> <div></div> </div> <div></div> </div>
1	B	422	<div> <div>3%</div> <div> <div></div> <div>90%</div> <div>7%</div> <div></div> </div> <div></div> </div>

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 7147 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitric oxide synthase, brain.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	407	Total	C	N	O	S	0	3	0
			3322	2127	566	607	22			
1	B	411	Total	C	N	O	S	0	2	0
			3354	2146	574	613	21			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula: $C_9H_{15}N_5O_3$).



Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	N	0	0
			23	19	4		

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Zn	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	117	Total	O	0	0
			117	117		
7	B	179	Total	O	0	0
			179	179		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nitric oxide synthase, brain



- Molecule 1: Nitric oxide synthase, brain



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.72Å 110.42Å 165.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.84 – 1.95 46.84 – 1.95	Depositor EDS
% Data completeness (in resolution range)	73.7 (46.84-1.95) 73.7 (46.84-1.95)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.26 (at 1.95Å)	Xtriage
Refinement program	PHENIX 1.8.2_1309	Depositor
R, R_{free}	0.177 , 0.231 0.177 , 0.231	Depositor DCC
R_{free} test set	2515 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	23.2	Xtriage
Anisotropy	0.111	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 55.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7147	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, ZN, ACT, H4B, JHT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.38	0/3421	0.55	0/4641
1	B	0.40	0/3453	0.56	0/4681
All	All	0.39	0/6874	0.55	0/9322

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3322	0	3232	34	0
1	B	3354	0	3270	21	0
2	A	43	0	30	4	0
2	B	43	0	30	4	0
3	A	17	0	15	0	0
3	B	17	0	15	0	0
4	A	23	0	0	2	0
4	B	23	0	0	2	0
5	A	4	0	3	0	0
5	B	4	0	3	0	0
6	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	117	0	0	0	0
7	B	179	0	0	1	0
All	All	7147	0	6598	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:596:ARG:NH2	1:A:600:ASP:OD2	2.03	0.91
2:B:801:HEM:HBB2	2:B:801:HEM:HHC	1.62	0.80
1:B:596:ARG:NH2	1:B:600:ASP:OD1	2.23	0.70
1:B:478:GLN:HB2	1:B:481:ARG:HG3	1.73	0.70
2:A:801:HEM:HMC2	2:A:801:HEM:HBC2	1.73	0.70
1:A:350:THR:N	1:A:353:GLN:OE1	2.27	0.67
2:B:801:HEM:HBC2	2:B:801:HEM:HMC2	1.80	0.64
1:B:322:LEU:HD13	1:B:699:ARG:HH21	1.63	0.64
1:A:478:GLN:HB2	1:A:481:ARG:HG3	1.84	0.60
2:A:801:HEM:HHC	2:A:801:HEM:HBB2	1.86	0.57
1:B:388:ILE:O	1:B:392:SER:N	2.35	0.56
1:A:455:LEU:HD12	1:A:587:TRP:HB3	1.88	0.56
1:B:475:TRP:HB2	1:B:523:LEU:HB3	1.87	0.55
1:A:480:ILE:HD13	1:A:541:VAL:HG13	1.91	0.52
2:A:801:HEM:HBA1	4:A:803:JHT:C22	2.40	0.52
1:B:487:GLN:OE1	1:B:514:ARG:NH2	2.45	0.50
1:A:687:ILE:HD12	1:B:626:LYS:HB3	1.95	0.49
1:A:567:VAL:HG21	4:A:803:JHT:C13	2.44	0.48
1:A:596:ARG:HH21	1:B:692:HIS:CE1	2.32	0.47
2:B:801:HEM:HBA1	4:B:803:JHT:C22	2.45	0.47
1:A:465:ARG:NH2	1:A:471:ASP:OD2	2.45	0.47
1:A:596:ARG:HH21	1:B:692:HIS:HE1	1.62	0.47
1:A:598:TYR:O	1:A:606:ILE:HG12	2.15	0.47
1:A:393:THR:OG1	1:A:394:TYR:N	2.46	0.46
1:A:355:PHE:HD1	1:A:388:ILE:HD12	1.79	0.46
1:A:330:ILE:HD11	1:B:696:LEU:HD22	1.98	0.45
1:A:455:LEU:HD12	1:A:587:TRP:CB	2.46	0.45
1:A:674:ALA:HB3	1:A:695:MET:HB3	1.98	0.45
1:A:596:ARG:NH2	1:B:692:HIS:CE1	2.85	0.45
2:B:801:HEM:O1A	7:B:901:HOH:O	2.21	0.45
1:A:686:SER:HA	1:A:691:PHE:CG	2.52	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:475:TRP:HB2	1:A:523:LEU:HB3	1.99	0.44
1:A:523:LEU:HD22	1:A:531:PRO:HB2	1.98	0.44
1:A:476:ASN:OD1	1:A:476:ASN:N	2.50	0.44
1:A:469:LYS:HA	1:A:469:LYS:HD2	1.78	0.44
1:A:330:ILE:HD11	1:B:696:LEU:HB3	1.99	0.44
1:A:362:LEU:HA	1:A:362:LEU:HD23	1.89	0.43
1:A:391:THR:O	1:A:392:SER:OG	2.32	0.43
1:B:525:GLN:HG3	1:B:529:ASN:O	2.19	0.43
1:A:626:LYS:HB3	1:B:687:ILE:HD12	2.00	0.43
1:A:706:TYR:OH	2:A:801:HEM:O1D	2.26	0.42
1:B:353:GLN:HB3	1:B:353:GLN:HE21	1.62	0.42
1:B:511:LYS:HD3	1:B:511:LYS:HA	1.85	0.42
1:B:353:GLN:HG2	1:B:353:GLN:H	1.54	0.42
1:A:485:TYR:HB3	1:A:514:ARG:NH1	2.34	0.42
1:A:596:ARG:CZ	1:A:600:ASP:OD2	2.67	0.42
1:A:525:GLN:HG3	1:A:529:ASN:O	2.20	0.41
1:A:548:HIS:CG	1:A:549:PRO:HD2	2.54	0.41
1:B:567:VAL:HG21	4:B:803:JHT:C13	2.51	0.41
1:A:403:TYR:CE1	1:A:407:HIS:CE1	3.09	0.41
1:B:362:LEU:HD11	1:B:384:VAL:HG21	2.03	0.41
1:B:473:ARG:HD3	1:B:580:SER:HB2	2.03	0.41
1:B:494:GLY:O	1:B:496:PRO:HD3	2.22	0.40
1:A:476:ASN:HD21	1:A:522:LEU:HA	1.85	0.40
1:A:515:GLY:N	1:A:518:ASP:OD2	2.43	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	405/422 (96%)	391 (96%)	14 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	409/422 (97%)	400 (98%)	9 (2%)	0	100	100
All	All	814/844 (96%)	791 (97%)	23 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	365/377 (97%)	353 (97%)	12 (3%)	38	26
1	B	368/377 (98%)	360 (98%)	8 (2%)	52	44
All	All	733/754 (97%)	713 (97%)	20 (3%)	44	34

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	337	LEU
1	A	470	HIS
1	A	476	ASN
1	A	489	ASP
1	A	507	GLN
1	A	547	ARG
1	A	550	LYS
1	A	555	LYS
1	A	620	LYS
1	A	645	LYS
1	A	656	GLU
1	A	715	VAL
1	B	320	SER
1	B	321	THR
1	B	332	MET
1	B	353	GLN
1	B	390	SER
1	B	392	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	423	LYS
1	B	547	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	353	GLN
1	B	440	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 1 is monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HEM	B	801	1	27,50,50	2.21	6 (22%)	17,82,82	1.55	2 (11%)
5	ACT	B	804	-	1,3,3	1.33	0	0,3,3	0.00	-
4	JHT	A	803	-	25,25,25	0.92	0	33,34,34	0.85	1 (3%)
5	ACT	A	804	-	1,3,3	1.34	0	0,3,3	0.00	-

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	H4B	A	802	-	16,18,18	0.78	0	11,26,26	2.74	6 (54%)
2	HEM	A	801	1	27,50,50	2.19	6 (22%)	17,82,82	1.84	5 (29%)
4	JHT	B	803	-	25,25,25	0.86	0	33,34,34	0.95	1 (3%)
3	H4B	B	802	-	16,18,18	0.87	0	11,26,26	2.47	5 (45%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	B	801	1	-	1/6/54/54	-
4	JHT	A	803	-	-	5/10/10/10	0/3/3/3
3	H4B	A	802	-	-	0/8/17/17	0/2/2/2
2	HEM	A	801	1	-	0/6/54/54	-
4	JHT	B	803	-	-	6/10/10/10	0/3/3/3
3	H4B	B	802	-	-	0/8/17/17	0/2/2/2

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	HEM	C3D-C2D	5.09	1.52	1.37
2	B	801	HEM	C3B-C2B	-4.86	1.33	1.40
2	A	801	HEM	C3B-C2B	-4.79	1.33	1.40
2	B	801	HEM	C3D-C2D	4.77	1.51	1.37
2	B	801	HEM	C3C-C2C	-4.28	1.34	1.40
2	B	801	HEM	C3C-CAC	4.04	1.56	1.47
2	A	801	HEM	C3C-CAC	3.86	1.55	1.47
2	A	801	HEM	C3B-CAB	3.69	1.55	1.47
2	A	801	HEM	C3C-C2C	-3.65	1.35	1.40
2	B	801	HEM	C3B-CAB	3.52	1.55	1.47
2	A	801	HEM	CAA-C2A	2.46	1.55	1.52
2	B	801	HEM	CAA-C2A	2.08	1.55	1.52

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	802	H4B	C4-C4A-C8A	5.01	119.02	114.57
3	A	802	H4B	C4-C4A-C8A	4.86	118.89	114.57
3	A	802	H4B	C4-C4A-N5	4.05	122.52	119.12
2	A	801	HEM	CBD-CAD-C3D	-4.00	105.11	112.48

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	801	HEM	CBD-CAD-C3D	-3.89	105.30	112.48
3	A	802	H4B	N3-C2-N1	-3.14	120.49	125.42
3	B	802	H4B	N3-C2-N1	-3.02	120.67	125.42
3	A	802	H4B	C2-N1-C8A	2.95	121.16	114.54
3	B	802	H4B	C4-N3-C2	2.95	120.62	115.93
2	A	801	HEM	C1D-C2D-C3D	-2.94	104.95	107.00
2	A	801	HEM	CBA-CAA-C2A	-2.89	107.15	112.49
3	A	802	H4B	C4-N3-C2	2.89	120.52	115.93
2	B	801	HEM	C1D-C2D-C3D	-2.79	105.06	107.00
3	A	802	H4B	C4A-N5-C6	-2.74	113.70	121.16
2	A	801	HEM	CAD-CBD-CGD	-2.67	108.18	112.67
3	B	802	H4B	C2-N1-C8A	2.63	120.42	114.54
3	B	802	H4B	C4-C4A-N5	2.42	121.15	119.12
4	B	803	JHT	C08-C09-N10	-2.33	106.75	112.80
2	A	801	HEM	C4A-C3A-C2A	2.26	108.57	107.00
4	A	803	JHT	C05-C04-N02	-2.07	118.83	121.63

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	801	HEM	C2A-CAA-CBA-CGA
4	A	803	JHT	C12-C11-N10-C09
4	B	803	JHT	C12-C11-N10-C09
4	A	803	JHT	C05-C04-N02-C03
4	A	803	JHT	C23-C04-N02-C01
4	A	803	JHT	C23-C04-N02-C03
4	B	803	JHT	C05-C04-N02-C03
4	A	803	JHT	C05-C04-N02-C01
4	B	803	JHT	C23-C04-N02-C01
4	B	803	JHT	C23-C04-N02-C03
4	B	803	JHT	C05-C04-N02-C01
4	B	803	JHT	C08-C09-N10-C11

There are no ring outliers.

4 monomers are involved in 10 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	801	HEM	4	0
4	A	803	JHT	2	0
2	A	801	HEM	4	0

Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	803	JHT	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand JHT A 803



Ligand HEM A 801





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	407/422 (96%)	0.24	18 (4%) 34 44	15, 42, 84, 122	0
1	B	411/422 (97%)	-0.01	11 (2%) 54 63	15, 33, 69, 93	0
All	All	818/844 (96%)	0.12	29 (3%) 44 53	15, 37, 79, 122	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	355	PHE	5.1
1	A	716	TRP	4.3
1	A	488	PRO	4.1
1	B	352	ASP	4.0
1	A	300	PHE	3.8
1	A	322	LEU	3.7
1	B	300	PHE	3.6
1	A	321	THR	3.3
1	A	352	ASP	3.1
1	B	321	THR	3.0
1	B	348	VAL	2.9
1	A	715	VAL	2.9
1	A	370	LYS	2.9
1	B	718	GLY	2.9
1	B	389	GLU	2.8
1	B	350	THR	2.8
1	A	388	ILE	2.8
1	A	551	PHE	2.5
1	A	713	THR	2.4
1	A	351	LYS	2.4
1	B	323	GLU	2.3
1	A	391	THR	2.3
1	B	338	PRO	2.2
1	A	554	PHE	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	619	ARG	2.2
1	B	307[A]	GLU	2.1
1	B	337	LEU	2.1
1	A	550	LYS	2.0
1	A	299	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	H4B	A	802	17/17	0.89	0.19	29,48,58,62	0
3	H4B	B	802	17/17	0.90	0.14	22,44,54,55	0
4	JHT	B	803	23/23	0.92	0.19	23,42,96,98	0
4	JHT	A	803	23/23	0.96	0.17	18,38,100,104	0
5	ACT	B	804	4/4	0.97	0.17	37,48,48,54	0
5	ACT	A	804	4/4	0.98	0.19	45,47,48,52	0
2	HEM	B	801	43/43	0.99	0.12	12,21,49,64	0
2	HEM	A	801	43/43	0.99	0.11	12,21,52,62	0
6	ZN	B	805	1/1	1.00	0.07	32,32,32,32	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around JHT B 803:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



**Electron density around JHT A 803:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM B 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM A 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.