



# Full wwPDB X-ray Structure Validation Report ⓘ

May 24, 2021 – 06:04 PM EDT

PDB ID : 1VXR  
Title : O-ETHYLMETHYLPHOSPHONYLATED ACETYLCHOLINESTERASE  
OBTAINED BY REACTION WITH O-ETHYL-S-[2-[BIS(1-METHYLETHYL)  
AMINO]ETHYL] METHYLPHOSPHONOTHIOATE (VX)  
Authors : Millard, C.B.; Koellner, G.; Silman, I.; Sussman, J.L.  
Deposited on : 1999-04-21  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.18


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	537	
2	B	2	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	A	9010	X	-	-	-

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 4655 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN (ACETYLCHOLINESTERASE).


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	532	Total	C	N	O	S	0	0	0
			4244	2721	720	781	22			

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is O-ETHYLMETHYLPHOSPHONIC ACID ESTER GROUP (three-letter code: VX) (formula:  $C_3H_9O_3P$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	P	0	0
			6	3	2	1		

- Molecule 5 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula:  $C_6H_{13}NO_4S$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
5	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	339	Total	O	0	0
			339	339		

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

#### • Molecule 1: PROTEIN (ACETYLCHOLINESTERASE)



#### • Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	112.84Å 112.84Å 137.36Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.20	Depositor
% Data completeness (in resolution range)	89.9 (30.00-2.20)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.06	Depositor
Refinement program	CNS	Depositor
R, $R_{free}$	0.189 , 0.230	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	4655	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: VX, MES, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.04	4/4367 (0.1%)	1.01	21/5929 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	446	PHE	CE1-CZ	5.61	1.48	1.37
1	A	206	VAL	CB-CG2	5.45	1.64	1.52
1	A	408	VAL	CB-CG1	-5.16	1.42	1.52
1	A	260	GLU	CG-CD	5.13	1.59	1.51

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	243	ARG	NE-CZ-NH2	-12.08	114.26	120.30
1	A	216	ARG	NE-CZ-NH2	-8.95	115.83	120.30
1	A	243	ARG	NE-CZ-NH1	8.81	124.70	120.30
1	A	388	ARG	NE-CZ-NH2	-8.40	116.10	120.30
1	A	216	ARG	NE-CZ-NH1	8.01	124.31	120.30
1	A	388	ARG	NE-CZ-NH1	7.09	123.85	120.30
1	A	221	ARG	NE-CZ-NH1	7.08	123.84	120.30
1	A	88	ARG	NE-CZ-NH2	-6.70	116.95	120.30
1	A	182	ASP	CB-CG-OD1	6.51	124.16	118.30

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	494	LEU	CA-CB-CG	6.48	130.21	115.30
1	A	83	MET	CG-SD-CE	6.13	110.01	100.20
1	A	518	VAL	CB-CA-C	-6.05	99.91	111.40
1	A	518	VAL	CG1-CB-CG2	5.63	119.90	110.90
1	A	243	ARG	CG-CD-NE	-5.42	100.41	111.80
1	A	221	ARG	NE-CZ-NH2	-5.38	117.61	120.30
1	A	517	ARG	NE-CZ-NH1	-5.30	117.65	120.30
1	A	220	ARG	NE-CZ-NH2	-5.26	117.67	120.30
1	A	175	MET	CG-SD-CE	-5.25	91.81	100.20
1	A	285	ASP	N-CA-C	-5.24	96.86	111.00
1	A	217	ASP	CB-CG-OD1	5.09	122.88	118.30
1	A	358	LEU	CA-CB-CG	5.07	126.95	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	63	TYR	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4244	0	4094	68	0
2	B	28	0	25	0	0
3	A	14	0	13	0	0
4	A	6	0	8	0	0
5	A	24	0	26	2	0
6	A	339	0	0	7	0
All	All	4655	0	4166	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:499:GLU:HG2	6:A:1276:HOH:O	1.81	0.79
1:A:310:ASN:OD1	1:A:410:LYS:NZ	2.21	0.73
1:A:455:GLU:CD	1:A:455:GLU:H	1.94	0.69
1:A:383:ASN:C	1:A:383:ASN:HD22	1.98	0.66
1:A:139:GLU:O	1:A:140:GLU:HG2	1.97	0.64
1:A:110:THR:OG1	1:A:478:LYS:HG2	1.98	0.64
1:A:468:ARG:HD3	6:A:1105:HOH:O	2.00	0.62
1:A:453:VAL:HG22	1:A:456:LEU:HG	1.82	0.61
1:A:50:PRO:HA	1:A:175:MET:HE3	1.83	0.60
1:A:216:ARG:HB3	1:A:315:LYS:HB2	1.83	0.60
1:A:344:GLU:OE1	1:A:346:LYS:HE3	2.03	0.58
1:A:468:ARG:CD	6:A:1105:HOH:O	2.52	0.57
1:A:452:LEU:HD22	1:A:463:GLU:HG3	1.88	0.56
1:A:47:ARG:HG3	6:A:1198:HOH:O	2.05	0.55
1:A:522:VAL:HG13	1:A:526:GLN:OE1	2.07	0.55
1:A:197:PHE:HA	1:A:223:ILE:O	2.08	0.54
1:A:139:GLU:C	1:A:140:GLU:HG2	2.28	0.53
1:A:260:GLU:H	1:A:260:GLU:CD	2.11	0.53
1:A:357:LYS:HD2	6:A:1288:HOH:O	2.09	0.53
1:A:49:GLU:O	1:A:175:MET:HE1	2.09	0.53
1:A:223:ILE:HA	1:A:320:LEU:O	2.08	0.53
1:A:302:PRO:HD2	1:A:308:MET:SD	2.49	0.52
1:A:474:ALA:O	1:A:478:LYS:HG3	2.09	0.52
1:A:36:ALA:HB2	1:A:175:MET:HE2	1.92	0.52
1:A:419:TYR:CZ	1:A:494:LEU:HD13	2.45	0.52
1:A:461:GLU:H	1:A:461:GLU:CD	2.12	0.51
1:A:197:PHE:CB	1:A:223:ILE:HB	2.42	0.50
1:A:531:LEU:C	1:A:531:LEU:HD23	2.32	0.50
1:A:247:GLU:HG2	5:A:8010:MES:O1S	2.12	0.50
1:A:221:ARG:HD3	1:A:318:GLN:OE1	2.13	0.49
1:A:383:ASN:ND2	1:A:386:LYS:H	2.11	0.49
1:A:452:LEU:HD13	1:A:467:ARG:NH2	2.28	0.49
1:A:500:GLN:HE22	5:A:8020:MES:H52	1.80	0.47
1:A:211:LEU:CD1	1:A:305:LEU:HD22	2.44	0.47
1:A:456:LEU:O	1:A:457:ASN:HB2	2.15	0.47
1:A:408:VAL:HG23	1:A:418:THR:HG21	1.97	0.46
1:A:36:ALA:HB2	1:A:175:MET:CE	2.46	0.46
1:A:110:THR:CB	1:A:478:LYS:HG2	2.45	0.46
1:A:197:PHE:HB3	1:A:223:ILE:HB	1.98	0.46
1:A:39:PRO:HA	1:A:43:MET:HE2	1.99	0.45
1:A:517:ARG:HG2	1:A:517:ARG:NH1	2.31	0.44
1:A:383:ASN:C	1:A:383:ASN:ND2	2.68	0.44

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:258:SER:HB2	1:A:260:GLU:OE1	2.17	0.44
1:A:408:VAL:CG2	1:A:418:THR:HG21	2.47	0.44
1:A:82:GLU:HG2	6:A:1171:HOH:O	2.17	0.44
1:A:366:LEU:HD23	1:A:535:THR:HG21	2.00	0.44
1:A:220:ARG:HG3	1:A:221:ARG:HG3	1.99	0.43
1:A:146:LEU:C	1:A:146:LEU:HD12	2.37	0.43
1:A:357:LYS:HD3	1:A:368:LEU:HD11	2.00	0.43
1:A:127:LEU:HD12	1:A:130:TYR:CE2	2.54	0.43
1:A:24:SER:O	1:A:24:SER:OG	2.30	0.43
1:A:207:GLY:HA3	1:A:229:PRO:HD3	2.01	0.43
1:A:360:VAL:HG12	1:A:363:ALA:HB2	2.00	0.43
1:A:499:GLU:CG	6:A:1276:HOH:O	2.55	0.43
1:A:219:PHE:O	1:A:315:LYS:HE2	2.18	0.43
1:A:517:ARG:HG2	1:A:517:ARG:HH11	1.84	0.43
1:A:451:PRO:HA	1:A:458:TYR:CD1	2.54	0.42
1:A:211:LEU:HD12	1:A:305:LEU:HD22	2.00	0.42
1:A:484:GLU:HB2	1:A:487:SER:CB	2.50	0.42
1:A:452:LEU:CD2	1:A:463:GLU:HG3	2.49	0.42
1:A:515:ARG:O	1:A:518:VAL:HG22	2.19	0.42
1:A:484:GLU:HB2	1:A:487:SER:HB2	2.02	0.42
1:A:185:GLN:HA	1:A:189:GLY:O	2.20	0.41
1:A:27:ILE:HD11	1:A:133:LYS:HB2	2.02	0.41
1:A:255:ASN:HB3	1:A:261:GLU:HG2	2.02	0.41
1:A:228:SER:HB2	1:A:229:PRO:HD2	2.01	0.41
1:A:366:LEU:CD2	1:A:535:THR:HG21	2.52	0.40
1:A:449:GLY:HA2	1:A:466:SER:OG	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	530/537 (99%)	503 (95%)	27 (5%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	465/469 (99%)	442 (95%)	23 (5%)	25	31

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	16	MET
1	A	129	VAL
1	A	140	GLU
1	A	161	SER
1	A	197	PHE
1	A	216	ARG
1	A	261	GLU
1	A	287	ILE
1	A	288	PHE
1	A	330	PHE
1	A	340	SER
1	A	349	ARG
1	A	354	SER
1	A	358	LEU
1	A	383	ASN
1	A	391	LEU
1	A	430	LEU
1	A	440	HIS
1	A	453	VAL
1	A	455	GLU
1	A	473	TRP
1	A	516	LEU
1	A	518	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	68	GLN
1	A	374	GLN
1	A	383	ASN
1	A	500	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	NAG	B	1	2,1	14,14,15	0.88	0	17,19,21	0.90	1 (5%)
2	NAG	B	2	2	14,14,15	1.31	2 (14%)	17,19,21	0.99	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	B	2	2	-	2/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2	NAG	C1-C2	2.51	1.56	1.52
2	B	2	NAG	C4-C5	2.08	1.57	1.53

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	NAG	C2-N2-C7	-2.77	118.96	122.90
2	B	1	NAG	C2-N2-C7	-2.00	120.05	122.90

There are no chirality outliers.


All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1	NAG	O5-C5-C6-O6
2	B	1	NAG	C4-C5-C6-O6
2	B	2	NAG	O5-C5-C6-O6
2	B	2	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	MES	A	8010	-	12,12,12	8.49	9 (75%)	14,16,16	2.17	3 (21%)
3	NAG	A	9010	1	14,14,15	1.27	1 (7%)	17,19,21	1.18	1 (5%)
4	VX	A	999	1	2,5,6	0.57	0	1,5,8	0.47	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	MES	A	8020	-	12,12,12	7.87	8 (66%)	14,16,16	2.46	5 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	MES	A	8010	-	-	4/6/14/14	0/1/1/1
3	NAG	A	9010	1	1/1/5/7	2/6/23/26	0/1/1/1
4	VX	A	999	1	-	0/1/3/4	-
5	MES	A	8020	-	-	3/6/14/14	0/1/1/1

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	8010	MES	C8-S	-21.45	1.47	1.77
5	A	8020	MES	C8-S	-19.77	1.49	1.77
5	A	8010	MES	O2S-S	11.42	1.78	1.45
5	A	8010	MES	O1S-S	11.31	1.78	1.45
5	A	8020	MES	O2S-S	10.97	1.77	1.45
5	A	8020	MES	O1S-S	10.60	1.76	1.45
5	A	8010	MES	O3S-S	9.99	1.82	1.47
5	A	8020	MES	O3S-S	8.66	1.78	1.47
5	A	8020	MES	C7-C8	-4.27	1.41	1.52
5	A	8010	MES	C7-C8	-4.14	1.41	1.52
5	A	8010	MES	C7-N4	-3.49	1.39	1.47
5	A	8020	MES	C7-N4	-3.08	1.40	1.47
3	A	9010	NAG	O5-C5	2.63	1.48	1.43
5	A	8020	MES	O1-C6	2.42	1.52	1.42
5	A	8020	MES	O1-C2	2.35	1.52	1.42
5	A	8010	MES	C3-C2	-2.22	1.41	1.50
5	A	8010	MES	C5-C6	-2.19	1.41	1.50
5	A	8010	MES	O1-C6	2.14	1.51	1.42

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	8020	MES	O1S-S-C8	5.75	113.83	106.92
5	A	8010	MES	O1S-S-C8	4.98	112.92	106.92
5	A	8020	MES	O2S-S-C8	3.99	111.72	106.92

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	8010	MES	O2S-S-C8	3.62	111.28	106.92
5	A	8020	MES	O3S-S-C8	3.46	111.37	105.77
3	A	9010	NAG	C2-N2-C7	-3.31	118.19	122.90
5	A	8010	MES	O3S-S-C8	2.66	110.07	105.77
5	A	8020	MES	O2S-S-O1S	-2.38	105.71	113.95
5	A	8020	MES	O3S-S-O1S	-2.08	106.19	111.27

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	A	9010	NAG	C1

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	8010	MES	C7-C8-S-O2S
5	A	8020	MES	C7-C8-S-O1S
5	A	8020	MES	C7-C8-S-O2S
3	A	9010	NAG	O5-C5-C6-O6
3	A	9010	NAG	C4-C5-C6-O6
5	A	8010	MES	C7-C8-S-O3S
5	A	8020	MES	C7-C8-S-O3S
5	A	8010	MES	N4-C7-C8-S
5	A	8010	MES	C7-C8-S-O1S

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	8010	MES	1	0
5	A	8020	MES	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

### 6.4 Ligands

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers

EDS was not executed - this section is therefore empty.