



Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2020 – 07:39 pm BST

PDB ID : 1VZ4
Title : Fe-Succinate Complex of AtsK
Authors : Mueller, I.; Stueckl, A.C.; Uson, I.; Kertesz, M.
Deposited on : 2004-05-14
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	301	 56% 22% • • 17%
1	D	301	 66% 18% • 14%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3937 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called PUTATIVE ALKYLSULFATASE ATSK.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	249	Total	C	N	O	0	0	0
			1896	1200	347	349			
1	D	259	Total	C	N	O	0	0	0
			1975	1250	359	366			

- Molecule 2 is FE (II) ION (three-letter code: FE2) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Fe	0	0
			1	1		
2	D	1	Total	Fe	0	0
			1	1		

- Molecule 3 is SUCCINIC ACID (three-letter code: SIN) (formula: C₄H₆O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total 8	C 4	O 4	0	0
3	D	1	Total 8	C 4	O 4	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	20	Total 20	O 20	0	0
4	D	28	Total 28	O 28	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	I 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	71.84Å 141.00Å 160.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.50 50.00 – 1.89	Depositor EDS
% Data completeness (in resolution range)	99.2 (50.00-2.50) 64.9 (50.00-1.89)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.36 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.2.0003	Depositor
R, R_{free}	0.227 , 0.287 0.232 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	30.7	Xtriage
Anisotropy	1.151	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 42.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3937	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FE2, SIN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.36	10/1940 (0.5%)	1.28	15/2652 (0.6%)
1	D	1.29	5/2021 (0.2%)	1.24	14/2762 (0.5%)
All	All	1.33	15/3961 (0.4%)	1.26	29/5414 (0.5%)

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	265	TYR	CE2-CZ	10.35	1.52	1.38
1	D	211	GLN	CG-CD	7.14	1.67	1.51
1	D	111	VAL	CA-CB	6.93	1.69	1.54
1	D	81	HIS	C-O	6.03	1.34	1.23
1	A	265	TYR	CE1-CZ	-5.98	1.30	1.38
1	A	249	TRP	CB-CG	-5.84	1.39	1.50
1	A	288	VAL	CB-CG2	5.78	1.65	1.52
1	A	76	GLY	N-CA	5.31	1.54	1.46
1	A	124	ARG	CG-CD	5.31	1.65	1.51
1	A	247	TRP	CB-CG	5.26	1.59	1.50
1	D	167	ASP	CB-CG	5.20	1.62	1.51
1	A	138	ALA	CA-CB	5.07	1.63	1.52
1	D	216	VAL	CB-CG2	5.04	1.63	1.52
1	A	145	GLN	CG-CD	5.01	1.62	1.51
1	A	133	GLY	C-O	5.01	1.31	1.23

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	37	LEU	CB-CG-CD1	-8.83	95.99	111.00
1	D	64	ASP	CB-CG-OD1	8.60	126.04	118.30
1	D	38	ASP	CB-CG-OD2	8.55	125.99	118.30
1	A	269	ASP	CB-CG-OD2	8.04	125.53	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	167	ASP	CB-CG-OD1	7.78	125.30	118.30
1	A	36	ASP	CB-CG-OD2	6.60	124.24	118.30
1	A	274	PRO	N-CD-CG	-6.32	93.72	103.20
1	A	29	ARG	NE-CZ-NH1	6.24	123.42	120.30
1	D	258	ASP	CB-CG-OD2	6.22	123.90	118.30
1	A	29	ARG	NE-CZ-NH2	-5.95	117.33	120.30
1	A	38	ASP	CB-CG-OD2	5.87	123.58	118.30
1	A	124	ARG	NE-CZ-NH1	-5.81	117.40	120.30
1	A	65	ASP	CB-CG-OD2	5.62	123.36	118.30
1	D	88	ASP	CB-CG-OD2	5.57	123.31	118.30
1	A	167	ASP	N-CA-C	5.50	125.86	111.00
1	D	197	PRO	N-CD-CG	-5.46	95.01	103.20
1	D	81	HIS	N-CA-C	5.43	125.67	111.00
1	A	53	LYS	CD-CE-NZ	-5.41	99.25	111.70
1	D	157	LYS	CB-CA-C	5.38	121.17	110.40
1	D	13	LEU	CA-CB-CG	5.37	127.64	115.30
1	D	269	ASP	CB-CG-OD1	5.34	123.10	118.30
1	D	134	ASP	CB-CG-OD1	5.27	123.05	118.30
1	D	147	LEU	CB-CG-CD1	-5.25	102.08	111.00
1	D	156	ASP	CB-CG-OD1	5.23	123.00	118.30
1	A	49	LEU	CA-CB-CG	5.15	127.14	115.30
1	A	135	THR	OG1-CB-CG2	-5.06	98.36	110.00
1	A	35	PRO	N-CD-CG	-5.05	95.63	103.20
1	A	224	LEU	CB-CG-CD2	5.02	119.53	111.00
1	A	156	ASP	CB-CG-OD2	5.01	122.81	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1896	0	1804	36	0
1	D	1975	0	1882	22	0
2	A	1	0	0	0	0
2	D	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	8	0	4	0	0
3	D	8	0	4	0	0
4	A	20	0	0	0	0
4	D	28	0	0	3	0
All	All	3937	0	3694	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:24:ILE:CG1	1:D:24:ILE:CD1	1.77	1.57
1:D:153:GLU:HG2	4:D:2013:HOH:O	1.51	1.10
1:A:15:LEU:HD12	1:A:15:LEU:H	1.46	0.80
1:A:126:VAL:HG21	1:A:278:ARG:HG3	1.68	0.75
1:A:224:LEU:O	1:A:228:GLN:HG3	1.87	0.73
1:A:15:LEU:HD11	1:A:44:ALA:HB1	1.73	0.70
1:D:24:ILE:CB	1:D:24:ILE:CD1	2.72	0.67
1:A:164:ASN:OD1	1:A:192:TYR:HB2	1.95	0.67
1:A:154:LEU:O	1:A:158:LEU:HG	1.97	0.65
1:D:42:VAL:HG13	1:D:75:LEU:HD23	1.80	0.62
1:D:242:GLU:OE1	1:D:242:GLU:N	2.29	0.61
1:D:121:SER:HB3	1:D:281:THR:HG22	1.81	0.61
1:A:111:VAL:HG12	1:A:114:VAL:HG22	1.83	0.61
1:D:164:ASN:OD1	1:D:192:TYR:HB2	2.02	0.58
1:D:122:ILE:HG12	1:D:256:ILE:HD12	1.87	0.57
1:D:50:VAL:O	1:D:53:LYS:HE3	2.05	0.56
1:D:229:HIS:HE1	4:D:2018:HOH:O	1.89	0.56
1:A:147:LEU:HB2	1:A:152:ARG:HG2	1.87	0.56
1:D:195:GLU:O	1:D:295:SER:HB3	2.05	0.56
1:A:33:LEU:HD22	1:A:71:PHE:CE1	2.42	0.55
1:A:146:GLU:OE2	1:A:237:HIS:ND1	2.25	0.54
1:A:287:PRO:HB2	1:A:295:SER:OG	2.07	0.54
1:D:123:LEU:HD11	1:D:277:VAL:HB	1.91	0.53
1:A:97:ASP:C	1:A:102:GLN:HG3	2.30	0.52
1:A:121:SER:HB2	1:A:257:TRP:CE2	2.47	0.50
1:A:54:VAL:HG23	1:A:257:TRP:HB3	1.95	0.48
1:D:111:VAL:HG13	1:D:118:PRO:HD2	1.96	0.48
1:D:205:SER:HB2	1:D:207:GLU:HG3	1.95	0.48
1:A:219:ILE:HG23	1:A:219:ILE:HD13	1.60	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:279:ARG:HD3	1:A:281:THR:HG23	1.95	0.47
1:A:279:ARG:HD3	1:A:281:THR:CG2	2.46	0.46
1:A:126:VAL:CG2	1:A:278:ARG:HG3	2.43	0.46
1:A:46:GLN:O	1:A:50:VAL:HG23	2.15	0.46
1:A:111:VAL:O	1:A:114:VAL:HG22	2.16	0.45
1:A:38:ASP:O	1:A:40:ALA:N	2.49	0.45
1:D:27:GLU:OE2	1:D:58:ARG:HG3	2.16	0.45
1:A:107:TRP:HB3	1:A:263:GLN:OE1	2.16	0.45
1:D:97:ASP:HB2	4:D:2003:HOH:O	2.17	0.45
1:A:123:LEU:HD11	1:A:277:VAL:CG2	2.47	0.45
1:D:76:GLY:O	1:D:78:PRO:HD3	2.17	0.45
1:A:62:HIS:HE2	1:A:63:LEU:HD13	1.82	0.44
1:A:64:ASP:O	1:A:65:ASP:C	2.55	0.44
1:A:58:ARG:O	1:A:60:GLN:HG2	2.17	0.43
1:A:76:GLY:O	1:A:78:PRO:HD3	2.19	0.43
1:A:103:ARG:NH1	1:A:271:GLY:O	2.51	0.43
1:D:107:TRP:CD1	1:D:265:TYR:HD1	2.38	0.42
1:A:144:TYR:O	1:A:145:GLN:C	2.58	0.42
1:A:50:VAL:O	1:A:53:LYS:NZ	2.44	0.42
1:D:122:ILE:HG12	1:D:256:ILE:CD1	2.48	0.42
1:D:97:ASP:CG	1:D:98:GLY:H	2.23	0.42
1:A:111:VAL:HB	1:A:118:PRO:HD2	2.01	0.41
1:D:285:GLU:H	1:D:285:GLU:CD	2.22	0.41
1:A:123:LEU:HD11	1:A:277:VAL:HG23	2.02	0.41
1:A:133:GLY:HA2	1:A:270:TYR:CD2	2.56	0.41
1:A:111:VAL:HG12	1:A:111:VAL:O	2.21	0.41
1:A:144:TYR:CE2	1:A:152:ARG:HD2	2.55	0.41
1:A:24:ILE:HG13	1:A:25:GLY:N	2.36	0.40
1:D:105:ASN:HB3	1:D:266:ALA:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	241/301 (80%)	223 (92%)	15 (6%)	3 (1%)	13	24
1	D	251/301 (83%)	235 (94%)	14 (6%)	2 (1%)	19	35
All	All	492/602 (82%)	458 (93%)	29 (6%)	5 (1%)	15	28

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	39	ALA
1	A	165	GLU
1	D	223	SER
1	A	80	ALA
1	D	80	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	187/244 (77%)	173 (92%)	14 (8%)	13	26
1	D	196/244 (80%)	179 (91%)	17 (9%)	10	20
All	All	383/488 (78%)	352 (92%)	31 (8%)	11	23

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	LEU
1	A	24	ILE
1	A	32	LYS
1	A	43	GLU
1	A	61	THR
1	A	67	SER
1	A	103	ARG
1	A	118	PRO
1	A	145	GLN
1	A	149	GLU
1	A	163	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	200	ARG
1	A	224	LEU
1	A	265	TYR
1	D	29	ARG
1	D	32	LYS
1	D	34	SER
1	D	43	GLU
1	D	53	LYS
1	D	63	LEU
1	D	103	ARG
1	D	111	VAL
1	D	121	SER
1	D	163	SER
1	D	211	GLN
1	D	218	ARG
1	D	223	SER
1	D	265	TYR
1	D	274	PRO
1	D	276	ILE
1	D	295	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	228	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SIN	A	1300	2	1,7,7	0.47	0	2,8,8	1.97	1 (50%)
3	SIN	D	1302	2	1,7,7	0.23	0	2,8,8	1.87	1 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SIN	A	1300	2	-	0/1/5/5	-
3	SIN	D	1302	2	-	0/1/5/5	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1300	SIN	C3-C2-C1	-2.76	108.04	112.67
3	D	1302	SIN	C3-C2-C1	-2.51	108.47	112.67

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	249/301 (82%)	-0.42	0 100 100	33, 52, 69, 86	0
1	D	259/301 (86%)	-0.45	0 100 100	36, 50, 68, 77	0
All	All	508/602 (84%)	-0.43	0 100 100	33, 51, 68, 86	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SIN	A	1300	8/8	0.94	0.16	48,51,57,61	0
3	SIN	D	1302	8/8	0.96	0.15	43,51,54,55	0
2	FE2	D	1301	1/1	1.00	0.16	45,45,45,45	0
2	FE2	A	1299	1/1	1.00	0.15	50,50,50,50	0

6.5 Other polymers

There are no such residues in this entry.