



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 20, 2021 – 10:08 AM EDT

PDB ID : 6VZO
Title : Crystal structure of human PPARgamma ligand binding domain (Protein delipidated by denature and refold)
Authors : Shang, J.; Kojetin, D.J.
Deposited on : 2020-02-28
Resolution : 2.27 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.23.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.23.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6980 (2.30-2.26)
Clashscore	141614	7711 (2.30-2.26)
Ramachandran outliers	138981	7597 (2.30-2.26)
Sidechain outliers	138945	7598 (2.30-2.26)
RSRZ outliers	127900	6849 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	<div> <div>8%</div> <div> <div></div> <div>75%</div> <div>18%</div> <div>• 5%</div> </div> </div>
1	B	275	<div> <div>5%</div> <div> <div></div> <div>78%</div> <div>12%</div> <div>• 8%</div> </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4392 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Peroxisome proliferator-activated receptor gamma.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	260	Total	C	N	O	S	0	0	0
			2083	1346	339	388	10			
1	B	254	Total	C	N	O	S	0	0	0
			2039	1320	334	376	9			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	138	Total	O	0	0
			138	138		
2	B	132	Total	O	0	0
			132	132		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Peroxisome proliferator-activated receptor gamma



- Molecule 1: Peroxisome proliferator-activated receptor gamma



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	93.04Å 62.10Å 118.85Å 90.00° 102.22° 90.00°	Depositor
Resolution (Å)	58.08 – 2.27 58.08 – 2.27	Depositor EDS
% Data completeness (in resolution range)	99.5 (58.08-2.27) 99.5 (58.08-2.27)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.81 (at 2.27Å)	Xtriage
Refinement program	PHENIX 1.11.1_2575	Depositor
R, R_{free}	0.218 , 0.271 0.223 , 0.267	Depositor DCC
R_{free} test set	1540 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	22.7	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 46.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	4392	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.53	0/2117	0.62	0/2850
1	B	0.53	1/2072 (0.0%)	0.64	0/2787
All	All	0.53	1/4189 (0.0%)	0.63	0/5637

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	358	LYS	C-N	8.13	1.49	1.34

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2083	0	2148	50	0
1	B	2039	0	2100	39	0
2	A	138	0	0	6	0
2	B	132	0	0	7	0
All	All	4392	0	4248	89	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (89) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:358:LYS:HB3	1:B:359:PRO:CD	1.50	1.33
1:B:358:LYS:HB3	1:B:359:PRO:HD2	1.17	1.10
1:B:358:LYS:HB3	1:B:359:PRO:HD3	1.38	1.01
1:B:358:LYS:CB	1:B:359:PRO:CD	2.40	0.98
1:B:358:LYS:HE3	1:B:459:THR:HG23	1.45	0.97
1:A:393:LEU:HD21	1:A:409:ILE:CG2	1.96	0.94
1:A:393:LEU:CD2	1:A:409:ILE:CG2	2.48	0.92
1:A:393:LEU:CD2	1:A:409:ILE:HB	2.00	0.91
1:A:393:LEU:HD22	1:A:409:ILE:HG22	1.59	0.84
1:A:393:LEU:HD21	1:A:409:ILE:HG21	1.61	0.82
1:B:358:LYS:CB	1:B:359:PRO:HD3	2.08	0.80
1:A:393:LEU:HD13	1:A:413:LEU:CD1	2.13	0.79
1:A:393:LEU:CD2	1:A:409:ILE:CB	2.60	0.79
1:B:325:ILE:HD12	1:B:388:ILE:HG23	1.66	0.78
1:B:358:LYS:CE	1:B:459:THR:HG23	2.16	0.75
1:A:212:ARG:O	1:A:216:LYS:HG2	1.88	0.73
1:B:327:TYR:CE2	1:B:367:LYS:HE3	2.23	0.73
1:A:285:CYS:SG	2:A:628:HOH:O	2.47	0.73
1:A:393:LEU:HD13	1:A:413:LEU:HD11	1.69	0.72
1:A:438:LYS:NZ	2:A:501:HOH:O	2.21	0.72
1:B:441:ASP:O	1:B:445:ILE:HG13	1.88	0.71
1:A:393:LEU:CD2	1:A:409:ILE:HG22	2.17	0.71
1:A:319:LYS:HZ3	1:A:474:LYS:HG3	1.54	0.71
1:A:393:LEU:HD23	1:A:409:ILE:HB	1.72	0.69
1:B:460:GLU:HG2	1:B:460:GLU:O	1.91	0.69
1:A:239:GLY:HA2	2:A:541:HOH:O	1.93	0.69
1:B:451:GLN:HA	1:B:454:GLN:HG2	1.77	0.67
1:B:262:ILE:HD11	1:B:345:GLN:HB3	1.77	0.66
1:A:325:ILE:HG23	1:A:388:ILE:HD12	1.78	0.65
1:A:393:LEU:HD21	1:A:409:ILE:CB	2.26	0.64
1:A:370:PHE:HB2	1:A:445:ILE:HD11	1.82	0.62
1:B:276:GLU:HG2	1:B:277:VAL:N	2.13	0.62
1:B:370:PHE:HB2	1:B:445:ILE:HD11	1.81	0.61
1:B:358:LYS:CB	1:B:359:PRO:HD2	2.11	0.60
1:A:393:LEU:HD22	1:A:409:ILE:CG2	2.22	0.60
1:A:350:ARG:NH2	1:A:365:GLU:OE2	2.28	0.60
1:B:358:LYS:HE3	1:B:459:THR:CG2	2.28	0.59
1:A:302:SER:HB3	2:A:513:HOH:O	2.02	0.58
1:A:319:LYS:NZ	1:A:474:LYS:CG	2.66	0.58
1:B:285:CYS:SG	2:B:623:HOH:O	2.54	0.58
1:B:358:LYS:CD	1:B:359:PRO:HD3	2.34	0.58
1:B:288:ARG:NH2	2:B:502:HOH:O	2.38	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:319:LYS:NZ	1:A:474:LYS:HG3	2.20	0.56
1:A:437:GLN:O	1:A:440:THR:HG22	2.05	0.56
1:A:463:MET:HG2	2:A:504:HOH:O	2.06	0.56
1:B:275:LYS:O	1:B:280:ARG:NH1	2.38	0.56
1:A:242:THR:O	1:A:242:THR:OG1	2.22	0.55
1:B:466:HIS:N	2:B:509:HOH:O	2.40	0.55
1:B:276:GLU:HG2	1:B:277:VAL:H	1.71	0.54
1:A:279:ILE:O	1:A:283:GLN:HG2	2.07	0.53
1:B:358:LYS:CG	1:B:359:PRO:HD3	2.38	0.52
1:A:276:GLU:OE2	1:A:357:ARG:HD3	2.09	0.52
1:B:275:LYS:HB3	1:B:280:ARG:HD3	1.92	0.51
1:A:207:GLU:HG3	1:A:209:ALA:H	1.75	0.51
1:A:422:LYS:HG3	2:A:594:HOH:O	2.12	0.50
1:A:235:ALA:O	1:A:238:THR:OG1	2.28	0.50
1:A:471:GLU:HA	1:A:474:LYS:HE2	1.94	0.50
1:B:313:ASP:OD2	2:B:501:HOH:O	2.20	0.49
1:A:320:TYR:CZ	1:A:398:PRO:HG2	2.47	0.49
1:B:446:VAL:O	1:B:450:VAL:HG13	2.12	0.49
1:A:465:LEU:HD23	1:A:470:GLN:HG2	1.96	0.47
1:A:333:LEU:HB3	1:A:340:LEU:HB2	1.96	0.47
1:A:323:HIS:CE1	1:A:473:TYR:CE1	3.03	0.46
1:A:436:LEU:O	1:A:439:MET:HB2	2.16	0.46
1:B:334:MET:HG2	1:B:339:VAL:HB	1.98	0.46
1:B:234:ARG:HD3	1:B:234:ARG:HA	1.81	0.45
1:A:357:ARG:O	1:A:357:ARG:HG3	2.16	0.45
1:A:282:PHE:O	1:A:286:GLN:HG2	2.17	0.45
1:B:234:ARG:NH2	2:B:517:HOH:O	2.49	0.45
1:A:319:LYS:HZ1	1:A:474:LYS:CG	2.30	0.44
1:A:370:PHE:CB	1:A:445:ILE:HD11	2.46	0.44
1:A:443:ARG:O	1:A:447:THR:HG23	2.17	0.44
1:A:319:LYS:HZ1	1:A:474:LYS:HG2	1.83	0.44
1:B:267:ILE:N	2:B:518:HOH:O	2.50	0.44
1:A:319:LYS:HZ3	1:A:474:LYS:CG	2.24	0.44
1:A:433:ALA:O	1:A:437:GLN:HG3	2.19	0.43
1:B:307:VAL:HG22	2:B:552:HOH:O	2.16	0.43
1:B:370:PHE:HB2	1:B:445:ILE:CD1	2.48	0.43
1:B:259:GLU:HG2	1:B:264:PHE:HD2	1.84	0.42
1:A:241:THR:O	1:A:241:THR:OG1	2.33	0.42
1:B:264:PHE:HD1	1:B:264:PHE:HA	1.72	0.42
1:B:394:SER:O	1:B:397:ARG:HG2	2.20	0.42
1:A:393:LEU:HD21	1:A:409:ILE:HB	1.86	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:300:ALA:HA	1:B:303:ILE:HD12	2.02	0.41
1:B:327:TYR:CZ	1:B:367:LYS:HE3	2.56	0.41
1:A:327:TYR:CE1	1:A:367:LYS:HE3	2.55	0.41
1:A:260:ASP:OD1	1:A:260:ASP:N	2.51	0.40
1:A:277:VAL:HG13	1:A:278:ALA:N	2.37	0.40
1:B:356:LEU:HB2	1:B:361:GLY:HA2	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	256/275 (93%)	249 (97%)	7 (3%)	0	100	100
1	B	246/275 (90%)	236 (96%)	8 (3%)	2 (1%)	19	22
All	All	502/550 (91%)	485 (97%)	15 (3%)	2 (0%)	34	40

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	357	ARG
1	B	358	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	233/248 (94%)	226 (97%)	7 (3%)	41	54
1	B	227/248 (92%)	221 (97%)	6 (3%)	46	60
All	All	460/496 (93%)	447 (97%)	13 (3%)	43	57

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	238	THR
1	A	257	MET
1	A	285	CYS
1	A	357	ARG
1	A	393	LEU
1	A	402	ASN
1	A	439	MET
1	B	234	ARG
1	B	262	ILE
1	B	267	ILE
1	B	275	LYS
1	B	322	VAL
1	B	357	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	260/275 (94%)	0.36	22 (8%) 10 14	7, 26, 61, 92	0
1	B	254/275 (92%)	0.37	14 (5%) 25 30	7, 27, 62, 74	0
All	All	514/550 (93%)	0.36	36 (7%) 16 20	7, 27, 62, 92	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	476	LEU	9.3
1	B	263	LYS	6.2
1	B	456	ILE	5.9
1	A	475	ASP	5.7
1	A	261	LYS	5.6
1	B	461	THR	4.5
1	B	475	ASP	4.1
1	A	243	ASP	3.9
1	A	241	THR	3.4
1	A	264	PHE	3.4
1	B	459	THR	3.4
1	A	240	LYS	3.2
1	A	262	ILE	3.2
1	A	238	THR	3.1
1	B	260	ASP	3.1
1	A	244	LYS	3.0
1	A	256	MET	2.9
1	B	245	SER	2.9
1	A	474	LYS	2.8
1	A	263	LYS	2.8
1	A	257	MET	2.7
1	A	239	GLY	2.6
1	A	260	ASP	2.5
1	B	363	PHE	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	267	ILE	2.4
1	A	393	LEU	2.4
1	A	242	THR	2.4
1	A	275	LYS	2.4
1	B	262	ILE	2.4
1	A	280	ARG	2.4
1	B	256	MET	2.3
1	A	258	GLY	2.3
1	A	253	ASN	2.2
1	B	473	TYR	2.2
1	B	454	GLN	2.0
1	B	261	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.