



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 29, 2020 – 06:43 PM BST

PDB ID : 6W0I
Title : Closed-gate KcsA soaked in 10mM KCl/5mM BaCl₂
Authors : Rohaim, A.; Gong, L.; Li, J.
Deposited on : 2020-02-29
Resolution : 2.33 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.13
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.33 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5974 (2.34-2.30)
Clashscore	141614	6604 (2.34-2.30)
Ramachandran outliers	138981	6523 (2.34-2.30)
Sidechain outliers	138945	6523 (2.34-2.30)
RSRZ outliers	127900	5855 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	219	<div> <div>3%</div> <div> <div></div> <div>82%</div> <div>18%</div> </div> </div>
2	B	212	<div> <div>2%</div> <div> <div></div> <div>82%</div> <div>18%</div> </div> </div>
3	C	103	<div> <div></div> <div> <div>90%</div> <div>10%</div> </div> </div>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4090 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fab Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	219	Total	C	N	O	S	0	0	0
			1605	1020	267	312	6			

- Molecule 2 is a protein called Fab Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	212	Total	C	N	O	S	0	0	0
			1646	1022	282	337	5			

- Molecule 3 is a protein called pH-gated potassium channel KcsA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	103	Total	C	N	O	S	0	0	0
			769	504	132	131	2			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	90	CYS	LEU	engineered mutation	UNP P0A334

- Molecule 4 is POTASSIUM ION (three-letter code: K) (formula: K) (labeled as "Ligand of Interest" by author).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	6	Total	K	0	0
			6	6		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	10	Total	O	0	0
			10	10		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	33	Total	O	0	0
			33	33		
5	C	21	Total	O	0	0
			21	21		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Fab Heavy Chain



- Molecule 2: Fab Light Chain



- Molecule 3: pH-gated potassium channel KcsA



4 Data and refinement statistics

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, α , β , γ	154.24Å 154.24Å 75.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	67.97 – 2.33 77.12 – 2.33	Depositor EDS
% Data completeness (in resolution range)	98.2 (67.97-2.33) 98.3 (77.12-2.33)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.40 (at 2.32Å)	Xtriage
Refinement program	PHENIX 1.14_3260	Depositor
R, R_{free}	0.200 , 0.232 0.201 , 0.232	Depositor DCC
R_{free} test set	1862 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	50.2	Xtriage
Anisotropy	0.342	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 52.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.026 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4090	wwPDB-VP
Average B, all atoms (Å ²)	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.38	0/1649	0.59	0/2261
2	B	0.41	0/1683	0.60	0/2283
3	C	0.49	0/789	0.58	0/1085
All	All	0.42	0/4121	0.59	0/5629

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1605	0	1530	21	0
2	B	1646	0	1572	22	0
3	C	769	0	775	6	0
4	C	6	0	0	0	0
5	A	10	0	0	0	0
5	B	33	0	0	0	0
5	C	21	0	0	1	1
All	All	4090	0	3877	45	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:83:LEU:HB3	1:A:86:LEU:HD21	1.76	0.67
2:B:37:GLN:HB2	2:B:47:LEU:HD11	1.83	0.61
1:A:124:PRO:HB3	1:A:150:TYR:HB3	1.84	0.59
2:B:11:LEU:HD23	2:B:19:VAL:CG2	2.33	0.59
3:C:44:SER:OG	3:C:66:LEU:HA	2.02	0.58
2:B:163:TRP:CD1	2:B:175:MET:HG3	2.38	0.58
1:A:38:LYS:HE2	1:A:40:ARG:HD2	1.86	0.58
2:B:13:VAL:HG22	2:B:17:GLU:HB3	1.85	0.57
1:A:218:ARG:HH22	2:B:119:PRO:HD2	1.70	0.56
1:A:31:SER:HB2	3:C:62:TYR:CE1	2.41	0.56
3:C:118:GLU:OE1	5:C:301:HOH:O	2.18	0.55
2:B:183:LYS:O	2:B:187:GLU:HG2	2.07	0.54
1:A:149:GLY:HA2	1:A:179:LEU:HB3	1.88	0.53
1:A:123:PRO:HB3	1:A:209:THR:HG21	1.91	0.53
2:B:155:ARG:HE	2:B:157:ASN:HB2	1.73	0.53
1:A:204:HIS:HB3	1:A:209:THR:OG1	2.10	0.52
2:B:193:THR:HG22	2:B:208:SER:HB3	1.92	0.52
2:B:138:ASN:ND2	2:B:170:ASP:OD2	2.43	0.52
2:B:107:LYS:HA	2:B:140:TYR:OH	2.10	0.51
1:A:193:TRP:CD1	1:A:194:PRO:HA	2.46	0.50
2:B:79:GLU:HG3	2:B:81:GLU:HG2	1.93	0.49
1:A:185:SER:HB2	2:B:135:PHE:CE2	2.49	0.48
3:C:115:VAL:O	3:C:119:GLN:HG2	2.16	0.46
1:A:32:ASP:OD2	1:A:100:ARG:NH1	2.49	0.46
1:A:12:VAL:HG11	1:A:86:LEU:HD12	1.98	0.45
2:B:186:TYR:CE2	2:B:211:ARG:HD3	2.53	0.44
2:B:185:GLU:HA	2:B:188:ARG:HE	1.82	0.44
1:A:11:LEU:HB2	1:A:152:PRO:HG3	2.00	0.44
1:A:195:SER:OG	1:A:196:GLU:N	2.50	0.44
1:A:194:PRO:HG3	1:A:217:PRO:HG3	1.99	0.44
2:B:11:LEU:HD23	2:B:19:VAL:HG22	1.97	0.44
1:A:12:VAL:O	1:A:116:VAL:HA	2.19	0.43
3:C:51:GLU:HG3	3:C:59:LEU:HB3	2.00	0.43
3:C:66:LEU:O	3:C:70:VAL:HG23	2.18	0.43
2:B:94:TRP:CG	2:B:95:PRO:HA	2.54	0.43
2:B:61:ARG:NH1	2:B:82:ASP:OD1	2.51	0.42
2:B:198:HIS:CE1	2:B:200:THR:HG23	2.55	0.42
1:A:35:HIS:CG	1:A:105:PHE:HE2	2.38	0.42
1:A:17:SER:HB2	1:A:84:SER:HA	2.01	0.42
2:B:189:HIS:O	2:B:211:ARG:NH2	2.53	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:76:ASN:O	2:B:77:SER:HB3	2.20	0.41
2:B:11:LEU:HD23	2:B:19:VAL:HG21	2.03	0.41
1:A:102:ASP:O	2:B:50:TYR:OH	2.29	0.40
1:A:143:LEU:HB3	1:A:215:ILE:HG21	2.04	0.40
1:A:17:SER:CB	1:A:84:SER:HA	2.52	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:C:311:HOH:O	5:C:313:HOH:O[3_755]	1.94	0.26

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	217/219 (99%)	203 (94%)	13 (6%)	1 (0%)	29	35
2	B	210/212 (99%)	202 (96%)	8 (4%)	0	100	100
3	C	101/103 (98%)	100 (99%)	1 (1%)	0	100	100
All	All	528/534 (99%)	505 (96%)	22 (4%)	1 (0%)	47	58

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	191	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	170/185 (92%)	164 (96%)	6 (4%)	36	49
2	B	189/190 (100%)	186 (98%)	3 (2%)	62	77
3	C	73/75 (97%)	72 (99%)	1 (1%)	67	80
All	All	432/450 (96%)	422 (98%)	10 (2%)	50	66

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	13	LYS
1	A	43	HIS
1	A	133	SER
1	A	148	LYS
1	A	200	CYS
1	A	201	ASN
2	B	70	ASP
2	B	153	SER
2	B	165	ASP
3	C	27	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	219/219 (100%)	0.22	7 (3%)	47	55	41, 81, 112, 130	0
2	B	212/212 (100%)	0.32	5 (2%)	59	66	37, 72, 120, 148	1 (0%)
3	C	103/103 (100%)	0.61	0	100	100	35, 45, 64, 83	0
All	All	534/534 (100%)	0.33	12 (2%)	62	69	35, 69, 115, 148	1 (0%)

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	179	LEU	3.9
2	B	209	PHE	3.3
1	A	63	LYS	3.1
1	A	55	TYR	2.9
1	A	67	LYS	2.8
2	B	118	PHE	2.8
1	A	129	LEU	2.6
1	A	158	THR	2.5
2	B	181	LEU	2.4
2	B	157	ASN	2.1
1	A	126	VAL	2.1
1	A	83	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	K	C	201	1/1	0.90	0.31	87,87,87,87	1
4	K	C	206	1/1	0.90	0.44	92,92,92,92	1
4	K	C	202	1/1	0.97	0.24	36,36,36,36	1
4	K	C	205	1/1	1.00	0.20	36,36,36,36	1
4	K	C	204	1/1	1.00	0.23	39,39,39,39	1
4	K	C	203	1/1	1.00	0.25	36,36,36,36	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around K C 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




Electron density around K C 206:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around K C 202:




$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around K C 204:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.