



Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 04:22 pm BST

PDB ID : 2W19
Title : Non-covalent complex between dahp synthase and chorismate mutase from Mycobacterium tuberculosis
Authors : Okvist, M.; Sasso, S.; Roderer, K.; Gamper, M.; Codoni, G.; Krenkel, U.; Kast, P.
Deposited on : 2008-10-16
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	472	<div> <div>3%</div> <div> <div></div> <div>83%</div> <div>6%</div> <div>11%</div> </div> </div>
1	B	472	<div> <div>%</div> <div> <div></div> <div>80%</div> <div>8%</div> <div>12%</div> </div> </div>
2	C	90	<div> <div>44%</div> <div> <div></div> <div>78%</div> <div>20%</div> </div> </div>
2	D	90	<div> <div>32%</div> <div> <div></div> <div>81%</div> <div>17%</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	1463	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7951 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 3-DEOXY-D-ARABINO-HEPTULOSONATE 7-PHOSPHATE SYNTHASE AROG.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	419	Total	C	N	O	S	0	0	1
			3211	2017	587	592	15			
1	B	417	Total	C	N	O	S	0	0	1
			3189	2001	583	590	15			

- Molecule 2 is a protein called CHORISMATE MUTASE.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	72	Total	C	N	O	S	0	0	0
			543	338	108	95	2			
2	D	75	Total	C	N	O	S	0	0	0
			576	357	114	103	2			

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	158	Total	O		0	0
			158	158			
5	B	189	Total	O		0	0
			189	189			
5	C	1	Total	O		0	0
			1	1			
5	D	3	Total	O		0	0
			3	3			


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 3-DEOXY-D-ARABINO-HEPTULOSONATE 7-PHOSPHATE SYNTHASE AROG




- Molecule 1: 3-DEOXY-D-ARABINO-HEPTULOSONATE 7-PHOSPHATE SYNTHASE AROG



- Molecule 2: CHORISMATE MUTASE



- Molecule 2: CHORISMATE MUTASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	204.02Å 204.02Å 66.51Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	66.82 – 2.15 66.78 – 2.15	Depositor EDS
% Data completeness (in resolution range)	96.6 (66.82-2.15) 92.5 (66.78-2.15)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.03 (at 2.14Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.178 , 0.211 0.182 , 0.213	Depositor DCC
R_{free} test set	4007 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	25.2	Xtriage
Anisotropy	0.288	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 53.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.028 for -h,-k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7951	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	0/3276	0.64	0/4459
1	B	0.52	0/3253	0.64	0/4427
2	C	0.39	0/546	0.61	0/726
2	D	0.39	0/579	0.56	0/770
All	All	0.49	0/7654	0.63	0/10382

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	C	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	C	20	GLU	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3211	0	3191	20	0
1	B	3189	0	3159	29	0
2	C	543	0	577	0	0
2	D	576	0	614	0	0
3	A	30	0	40	4	0
3	B	36	0	48	3	0
4	A	5	0	0	0	0
4	B	10	0	0	0	0
5	A	158	0	0	2	0
5	B	189	0	0	2	0
5	C	1	0	0	0	0
5	D	3	0	0	0	0
All	All	7951	0	7629	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:239:GLN:HB3	1:A:240:THR:C	1.90	0.90
1:B:197:VAL:HA	1:B:200:TRP:CE3	2.25	0.70
1:A:239:GLN:HB3	1:A:240:THR:CA	2.25	0.66
1:A:197:VAL:HA	1:A:200:TRP:CE3	2.32	0.64
1:B:96:GLU:HG3	1:B:97:PRO:HD3	1.83	0.61
1:B:197:VAL:HA	1:B:200:TRP:CZ3	2.36	0.60
1:A:122:VAL:HA	3:A:1463:GOL:H11	1.83	0.60
1:A:281:ILE:HD11	1:A:296:ALA:HB2	1.85	0.57
1:B:9:ILE:HD13	1:B:51:VAL:CG2	2.37	0.55
1:B:297:GLN:NE2	5:B:2135:HOH:O	2.41	0.52
1:A:261:LEU:HD11	1:A:269:PRO:HB3	1.89	0.52
1:B:9:ILE:HD13	1:B:51:VAL:HG23	1.90	0.51
1:A:12:LEU:HD22	1:B:92:MET:HA	1.93	0.51
1:B:9:ILE:CD1	1:B:51:VAL:HG23	2.40	0.51
1:A:4:THR:HG22	1:B:8:PRO:HA	1.93	0.50
1:B:83:GLN:HA	1:B:124:VAL:O	2.13	0.48
1:A:197:VAL:HA	1:A:200:TRP:CZ3	2.48	0.48
1:B:395:PHE:O	1:B:399:ARG:HG2	2.16	0.46
1:B:104:ARG:HG3	1:B:200:TRP:CE3	2.52	0.45
1:B:115:TYR:OH	1:B:220:GLU:HG2	2.17	0.45
1:B:294:ALA:HA	1:B:297:GLN:HE21	1.81	0.44
1:B:18:LEU:N	3:B:1469:GOL:H2	2.32	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:381:THR:HG22	1:B:386:ARG:HG3	1.99	0.44
1:B:23:ARG:HG2	3:B:1469:GOL:H11	2.00	0.44
1:A:121:VAL:O	3:A:1463:GOL:H11	2.18	0.44
1:B:114:THR:OG1	3:B:1464:GOL:H12	2.17	0.44
1:B:271:LEU:HD12	1:B:298:VAL:CG1	2.48	0.44
1:B:134:PRO:HB2	1:B:284:ARG:HD2	2.00	0.43
1:A:123:LYS:NZ	3:A:1463:GOL:H12	2.33	0.43
1:B:104:ARG:HG3	1:B:200:TRP:CZ3	2.53	0.43
1:A:117:ALA:HB1	1:A:119:MET:HG2	2.01	0.43
1:B:383:HIS:CE1	1:B:387:ILE:HD11	2.53	0.43
1:A:297:GLN:NE2	5:A:2133:HOH:O	2.51	0.43
1:B:52:LEU:O	1:B:55:VAL:HG22	2.19	0.42
1:B:197:VAL:HG22	1:B:200:TRP:HZ3	1.84	0.42
1:B:7:ILE:HA	1:B:8:PRO:HD3	1.92	0.42
1:A:327:ASN:ND2	5:A:2141:HOH:O	2.52	0.42
1:A:239:GLN:CB	1:A:240:THR:CA	2.95	0.42
1:A:83:GLN:HA	1:A:124:VAL:O	2.19	0.42
1:A:4:THR:HG22	1:B:8:PRO:CA	2.50	0.41
1:A:9:ILE:HD11	1:B:5:VAL:HG23	2.03	0.41
1:B:327:ASN:ND2	5:B:2151:HOH:O	2.53	0.41
1:A:123:LYS:HZ3	3:A:1463:GOL:C1	2.34	0.41
1:B:187:THR:CG2	1:B:243:ILE:HD12	2.52	0.40
1:A:347:LEU:N	1:A:348:PRO:CD	2.85	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	411/472 (87%)	403 (98%)	8 (2%)	0	100	100
1	B	409/472 (87%)	403 (98%)	5 (1%)	1 (0%)	47	46

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	C	70/90 (78%)	69 (99%)	1 (1%)	0	100	100
2	D	73/90 (81%)	71 (97%)	2 (3%)	0	100	100
All	All	963/1124 (86%)	946 (98%)	16 (2%)	1 (0%)	51	53

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	461	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	336/385 (87%)	326 (97%)	10 (3%)	41	40
1	B	332/385 (86%)	325 (98%)	7 (2%)	53	57
2	C	53/75 (71%)	52 (98%)	1 (2%)	57	61
2	D	58/75 (77%)	56 (97%)	2 (3%)	37	35
All	All	779/920 (85%)	759 (97%)	20 (3%)	46	47

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	40	TRP
1	A	191	LEU
1	A	196	LEU
1	A	215	GLU
1	A	226	ARG
1	A	235	ASP
1	A	240	THR
1	A	297	GLN
1	A	327	ASN
1	A	337	ARG
1	B	4	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	40	TRP
1	B	96	GLU
1	B	225	LEU
1	B	309	PRO
1	B	327	ASN
1	B	337	ARG
2	C	85	ARG
2	D	85	ARG
2	D	87	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	297	GLN
1	A	327	ASN
1	A	404	HIS
1	A	447	GLN
1	B	237	ASN
1	B	297	GLN
1	B	327	ASN
1	B	383	HIS
2	C	56	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	A	1462	-	5,5,5	0.33	0	5,5,5	0.62	0
3	GOL	B	1466	-	5,5,5	0.47	0	5,5,5	0.74	0
3	GOL	A	1466	-	5,5,5	0.54	0	5,5,5	0.51	0
4	SO4	A	1467	-	4,4,4	0.19	0	6,6,6	0.25	0
3	GOL	B	1469	-	5,5,5	0.91	0	5,5,5	1.34	0
4	SO4	B	1463	-	4,4,4	0.17	0	6,6,6	0.44	0
3	GOL	A	1465	-	5,5,5	0.40	0	5,5,5	0.19	0
3	GOL	B	1467	-	5,5,5	0.32	0	5,5,5	0.47	0
3	GOL	B	1464	-	5,5,5	0.40	0	5,5,5	0.33	0
4	SO4	B	1462	-	4,4,4	0.19	0	6,6,6	0.44	0
3	GOL	A	1463	-	5,5,5	0.73	0	5,5,5	1.27	1 (20%)
3	GOL	A	1464	-	5,5,5	0.39	0	5,5,5	0.72	0
3	GOL	B	1468	-	5,5,5	0.59	0	5,5,5	0.41	0
3	GOL	B	1465	-	5,5,5	0.31	0	5,5,5	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	1462	-	-	2/4/4/4	-
3	GOL	B	1466	-	-	0/4/4/4	-
3	GOL	A	1466	-	-	3/4/4/4	-
3	GOL	B	1469	-	-	4/4/4/4	-
3	GOL	A	1465	-	-	2/4/4/4	-
3	GOL	B	1467	-	-	2/4/4/4	-
3	GOL	B	1464	-	-	0/4/4/4	-
3	GOL	A	1463	-	-	3/4/4/4	-
3	GOL	A	1464	-	-	2/4/4/4	-
3	GOL	B	1468	-	-	2/4/4/4	-
3	GOL	B	1465	-	-	0/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	A	1463	GOL	O1-C1-C2	2.38	121.59	110.20

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1462	GOL	O1-C1-C2-C3
3	A	1466	GOL	O1-C1-C2-C3
3	B	1469	GOL	O1-C1-C2-C3
3	B	1469	GOL	C1-C2-C3-O3
3	A	1465	GOL	C1-C2-C3-O3
3	A	1465	GOL	O2-C2-C3-O3
3	A	1463	GOL	C1-C2-C3-O3
3	B	1468	GOL	O1-C1-C2-C3
3	B	1467	GOL	O2-C2-C3-O3
3	B	1467	GOL	C1-C2-C3-O3
3	A	1464	GOL	C1-C2-C3-O3
3	B	1469	GOL	O1-C1-C2-O2
3	B	1469	GOL	O2-C2-C3-O3
3	A	1463	GOL	O2-C2-C3-O3
3	B	1468	GOL	O1-C1-C2-O2
3	A	1466	GOL	O1-C1-C2-O2
3	A	1462	GOL	O1-C1-C2-O2
3	A	1464	GOL	O2-C2-C3-O3
3	A	1463	GOL	O1-C1-C2-O2
3	A	1466	GOL	O2-C2-C3-O3

There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1469	GOL	2	0
3	B	1464	GOL	1	0
3	A	1463	GOL	4	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	419/472 (88%)	0.06	12 (2%) 51 61	18, 28, 44, 57	0
1	B	417/472 (88%)	0.01	6 (1%) 75 80	19, 27, 44, 58	0
2	C	72/90 (80%)	2.36	40 (55%) 0 0	23, 32, 35, 36	0
2	D	75/90 (83%)	1.84	29 (38%) 0 0	23, 32, 35, 37	0
All	All	983/1124 (87%)	0.34	87 (8%) 9 14	18, 28, 43, 58	0

All (87) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	48	ALA	6.5
2	C	21	ILE	6.1
2	D	17	LEU	6.0
1	A	381	THR	5.9
1	A	462	ASP	5.7
2	C	43	GLY	5.6
2	D	16	THR	5.6
2	C	47	MET	5.0
2	D	50	GLY	4.9
2	C	49	SER	4.4
2	D	48	ALA	4.4
1	A	240	THR	4.4
2	C	86	GLY	4.3
2	D	41	ALA	4.2
2	D	21	ILE	4.1
1	A	237	ASN	3.9
2	C	24	LEU	3.9
2	C	30	ALA	3.8
2	C	38	VAL	3.8
2	C	42	ILE	3.8
2	C	46	ARG	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	C	54	LEU	3.7
2	C	53	ARG	3.7
2	D	65	ARG	3.7
2	C	31	LEU	3.6
1	B	237	ASN	3.6
2	D	19	GLU	3.5
2	D	87	ARG	3.5
2	D	45	ALA	3.5
2	D	23	ARG	3.5
2	C	41	ALA	3.5
2	C	90	HIS	3.4
2	C	44	LYS	3.3
2	D	84	GLY	3.3
2	C	67	SER	3.3
2	D	38	VAL	3.3
2	C	20	GLU	3.3
2	D	44	LYS	3.3
2	C	65	ARG	3.2
2	C	45	ALA	3.2
2	D	53	ARG	3.2
2	D	34	ARG	3.2
2	D	18	ARG	3.1
2	C	83	LEU	3.1
2	D	49	SER	3.1
2	C	60	MET	3.1
2	C	50	GLY	3.1
2	D	39	SER	3.1
1	A	443	ARG	3.1
2	C	39	SER	3.0
2	D	68	GLU	3.0
2	D	24	LEU	3.0
2	C	37	GLU	2.9
2	C	55	VAL	2.8
2	C	34	ARG	2.8
2	C	32	VAL	2.8
2	C	64	GLU	2.8
1	B	9	ILE	2.8
2	C	28	ILE	2.8
2	D	22	ASP	2.6
2	C	85	ARG	2.6
2	C	58	ARG	2.6
2	C	61	LYS	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	11	GLN	2.6
1	A	191	LEU	2.6
1	A	236	ARG	2.5
2	C	79	LEU	2.6
1	B	8	PRO	2.5
1	A	370	GLY	2.5
2	D	58	ARG	2.4
2	D	52	THR	2.4
2	C	87	ARG	2.4
2	D	30	ALA	2.4
2	D	32	VAL	2.3
1	B	155	PHE	2.3
2	C	36	ALA	2.3
1	A	235	ASP	2.2
1	B	3	TRP	2.2
2	C	71	PRO	2.1
2	D	42	ILE	2.1
1	A	200	TRP	2.1
2	D	20	GLU	2.1
2	C	68	GLU	2.1
1	B	200	TRP	2.1
1	A	3	TRP	2.0
2	D	85	ARG	2.0
2	C	25	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	GOL	A	1466	6/6	0.84	0.20	42,48,49,49	0
3	GOL	A	1465	6/6	0.85	0.18	57,60,62,62	0
3	GOL	A	1463	6/6	0.85	0.12	23,29,32,34	0
3	GOL	B	1468	6/6	0.85	0.24	50,52,52,54	0
3	GOL	B	1466	6/6	0.90	0.13	26,33,35,35	0
3	GOL	A	1464	6/6	0.91	0.10	33,41,44,45	0
3	GOL	B	1469	6/6	0.91	0.22	27,32,32,33	0
3	GOL	B	1464	6/6	0.92	0.13	28,30,31,31	0
3	GOL	A	1462	6/6	0.94	0.09	30,31,31,32	0
3	GOL	B	1465	6/6	0.97	0.07	23,25,27,28	0
3	GOL	B	1467	6/6	0.98	0.10	25,29,30,30	0
4	SO4	A	1467	5/5	0.99	0.08	32,35,36,36	0
4	SO4	B	1462	5/5	0.99	0.04	27,28,30,30	0
4	SO4	B	1463	5/5	0.99	0.09	31,31,32,32	0

6.5 Other polymers [i](#)

There are no such residues in this entry.