



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 12, 2020 – 10:30 AM EDT

PDB ID : 5W26  
Title : INFLUENZA VIRUS NEURAMINIDASE N9 IN COMPLEX WITH 4-DEOXYGENATED 2,3-DIFLUORO-N-ACETYLNEURAMINIC ACID  
Authors : Streltsov, V.A.; Mckimm-Breschkin, J.; Barrett, S.; Pilling, P.; Hader, S.; Watt, A.G.  
Deposited on : 2017-06-05  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.14.6
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.14.6


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	388	<div> <div>2%</div> <div>90%</div> <div>8%</div> </div>
2	B	8	<div>100%</div>
3	C	2	<div>50%</div> <div>50%</div>

## 2 Entry composition [i](#)


There are 9 unique types of molecules in this entry. The entry contains 3718 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Neuraminidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	388	3090	1924	548	595	23	0	7	0

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	8	94	52	2	40	0	0	0

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	2	28	16	2	10	0	0	0

- Molecule 4 is (2R,3R,5R,6R)-5-acetamido-2,3-bis(fluoranyl)-6-[(1R,2R)-1,2,3-tris(oxidanyl)propyl]oxane-2-carboxylic acid (three-letter code: 9SG) (formula: C<sub>11</sub>H<sub>17</sub>F<sub>2</sub>NO<sub>7</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	F	N	O	0	0
			21	11	2	1	7		

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Ca	0	0
			1	1		

- Molecule 7 is (2 {R},3 {R},5 {R})-3-acetamido-5-fluoranyl-2-[(1 {R},2 {R})-1,2,3-tris(oxidanyl)propyl]-2,3,4,5-tetrahydropyran-1-ium-6-carboxylic acid (three-letter code: 9WM) (formula: C<sub>11</sub>H<sub>17</sub>FNO<sub>7</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	A	1	Total	C	F	N	O	0	1
			20	11	1	1	7		

- Molecule 8 is 5-acetamido-2,6-anhydro-3,4,5-trideoxy-3-fluoro-D-erythro-L-glucos-9-nononic acid (three-letter code: 9VP) (formula: C<sub>11</sub>H<sub>18</sub>FNO<sub>7</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
8	A	1	Total	C	F	N	O	0	1
			20	11	1	1	7		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	430	Total	O	0	0
			430	430		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Neuraminidase



- Molecule 2: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 4 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	180.94Å 180.94Å 180.94Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.00 – 1.90 45.23 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.9 (42.00-1.90) 99.9 (45.23-1.90)	Depositor EDS
$R_{merge}$	0.17	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.67 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.8.0151	Depositor
R, $R_{free}$	0.136 , 0.177 0.150 , 0.185	Depositor DCC
$R_{free}$ test set	2004 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.5	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 47.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	3718	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.95% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, 9VP, NAG, CA, 9SG, 9WM, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.97	4/3210 (0.1%)	1.14	16/4370 (0.4%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	128	GLU	CD-OE2	6.68	1.32	1.25
1	A	119	GLU	CD-OE2	5.72	1.31	1.25
1	A	372	SER	CB-OG	-5.66	1.34	1.42
1	A	364	ARG	CD-NE	-5.62	1.36	1.46

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	364	ARG	NE-CZ-NH1	24.66	132.63	120.30
1	A	364	ARG	NE-CZ-NH2	-22.63	108.99	120.30
1	A	173	ARG	NE-CZ-NH1	13.97	127.28	120.30
1	A	173	ARG	NE-CZ-NH2	-10.26	115.17	120.30
1	A	364	ARG	CB-CG-CD	8.77	134.39	111.60
1	A	358[A]	VAL	CG1-CB-CG2	8.29	124.17	110.90
1	A	358[B]	VAL	CG1-CB-CG2	8.29	124.17	110.90
1	A	210	ARG	NE-CZ-NH1	-7.49	116.55	120.30
1	A	447	MET	CG-SD-CE	6.76	111.01	100.20
1	A	364	ARG	CD-NE-CZ	6.71	132.99	123.60
1	A	127	ASP	CB-CG-OD1	6.47	124.12	118.30
1	A	221	ARG	NE-CZ-NH2	-6.26	117.17	120.30
1	A	125	ASP	CB-CG-OD1	5.91	123.62	118.30
1	A	307	ASP	CB-CG-OD1	5.46	123.22	118.30
1	A	358[A]	VAL	CA-CB-CG1	5.17	118.66	110.90
1	A	358[B]	VAL	CA-CB-CG1	5.17	118.66	110.90

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3090	0	2921	24	0
2	B	94	0	79	0	0
3	C	28	0	25	1	0
4	A	21	0	0	0	0
5	A	14	0	13	0	0
6	A	1	0	0	0	0
7	A	20	0	0	2	0
8	A	20	0	0	1	0
9	A	430	0	0	3	5
All	All	3718	0	3038	24	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:168:THR:H	1:A:171:ASN:HD21	1.28	0.80
1:A:285[B]:ARG:NH2	9:A:601:HOH:O	2.13	0.80
1:A:87:LEU:H	1:A:234:HIS:HD2	1.31	0.77
1:A:364:ARG:HA	9:A:905:HOH:O	1.94	0.65
1:A:98:HIS:HE1	1:A:420:ARG:HH11	1.45	0.64
1:A:87:LEU:H	1:A:234:HIS:CD2	2.15	0.62
1:A:97:TRP:H	1:A:394:GLN:HE22	1.52	0.56
1:A:173:ARG:HD3	1:A:210:ARG:CZ	2.38	0.54
1:A:168:THR:H	1:A:171:ASN:ND2	2.02	0.51
1:A:98:HIS:CE1	1:A:420:ARG:HH11	2.26	0.50
1:A:364:ARG:HD2	1:A:375:GLU:OE2	2.12	0.50
1:A:98:HIS:HD2	1:A:99:ILE:O	1.93	0.50
1:A:285[B]:ARG:NH1	9:A:602:HOH:O	2.19	0.49
1:A:121:TYR:CG	1:A:229:SER:HA	2.50	0.47

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:173:ARG:HD3	1:A:210:ARG:NH1	2.30	0.47
1:A:405:TYR:OH	7:A:514[B]:9WM:C2	2.65	0.45
1:A:168:THR:OG1	1:A:171:ASN:ND2	2.50	0.44
1:A:140:ILE:HD11	1:A:439:TRP:CZ2	2.54	0.43
1:A:423:PHE:CZ	1:A:447:MET:HG3	2.54	0.43
1:A:405:TYR:CE1	8:A:515[A]:9VP:C1	3.03	0.42
1:A:367:SER:HB2	1:A:399:ASN:HD21	1.86	0.41
1:A:277:GLU:OE2	7:A:514[B]:9WM:O9	2.38	0.41
1:A:83:ASP:HB2	3:C:1:NAG:H83	2.03	0.41
1:A:279:CYS:HB3	1:A:290:CYS:HB3	2.04	0.40

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
9:A:642:HOH:O	9:A:994:HOH:O[15_555]	1.81	0.39
9:A:806:HOH:O	9:A:957:HOH:O[15_555]	1.94	0.26
9:A:975:HOH:O	9:A:977:HOH:O[9_555]	1.98	0.22
9:A:659:HOH:O	9:A:763:HOH:O[16_555]	2.11	0.09
9:A:994:HOH:O	9:A:998:HOH:O[16_555]	2.13	0.07

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	392/388 (101%)	377 (96%)	15 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	348/341 (102%)	341 (98%)	7 (2%)	55 51

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	171	ASN
1	A	222	ASN
1	A	358[A]	VAL
1	A	358[B]	VAL
1	A	364	ARG
1	A	372	SER
1	A	415	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	95	ASN
1	A	98	HIS
1	A	171	ASN
1	A	222	ASN
1	A	234	HIS
1	A	345	ASN
1	A	346	ASN
1	A	392	GLN
1	A	394	GLN
1	A	399	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

10 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	B	1	1,2	14,14,15	0.68	0	17,19,21	1.99	4 (23%)
2	NAG	B	2	2	14,14,15	1.20	1 (7%)	17,19,21	1.50	5 (29%)
2	BMA	B	3	2	11,11,12	0.55	0	15,15,17	1.30	1 (6%)
2	MAN	B	4	2	11,11,12	0.83	0	15,15,17	1.93	5 (33%)
2	MAN	B	5	2	11,11,12	0.78	0	15,15,17	1.42	2 (13%)
2	MAN	B	6	2	11,11,12	0.69	0	15,15,17	1.14	2 (13%)
2	MAN	B	7	2	11,11,12	0.73	0	15,15,17	2.19	5 (33%)
2	MAN	B	8	2	11,11,12	1.10	0	15,15,17	1.64	4 (26%)
3	NAG	C	1	1,3	14,14,15	0.94	0	17,19,21	1.49	3 (17%)
3	NAG	C	2	3	14,14,15	1.22	1 (7%)	17,19,21	1.73	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1
2	BMA	B	3	2	-	0/2/19/22	0/1/1/1
2	MAN	B	4	2	-	0/2/19/22	0/1/1/1
2	MAN	B	5	2	-	0/2/19/22	0/1/1/1
2	MAN	B	6	2	-	0/2/19/22	0/1/1/1
2	MAN	B	7	2	-	1/2/19/22	0/1/1/1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MAN	B	8	2	-	2/2/19/22	0/1/1/1
3	NAG	C	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	C	2	3	-	2/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2	NAG	O5-C1	-3.47	1.38	1.43
3	C	2	NAG	C1-C2	2.23	1.55	1.52

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	7	MAN	C1-O5-C5	4.44	118.21	112.19
3	C	2	NAG	C2-N2-C7	4.39	129.15	122.90
2	B	4	MAN	O2-C2-C1	-4.33	100.30	109.15
2	B	5	MAN	O5-C5-C6	4.23	113.84	107.20
2	B	7	MAN	O5-C5-C6	4.13	113.67	107.20
2	B	1	NAG	C1-C2-N2	-3.99	103.67	110.49
2	B	7	MAN	O4-C4-C3	-3.90	101.34	110.35
3	C	1	NAG	C8-C7-N2	3.72	122.40	116.10
2	B	1	NAG	O5-C1-C2	-3.66	105.51	111.29
2	B	1	NAG	C8-C7-N2	-3.62	109.97	116.10
2	B	8	MAN	O5-C1-C2	-3.50	105.36	110.77
2	B	4	MAN	C1-O5-C5	3.44	116.85	112.19
2	B	7	MAN	O6-C6-C5	2.72	120.63	111.29
2	B	7	MAN	C3-C4-C5	2.71	115.08	110.24
2	B	2	NAG	O5-C1-C2	-2.68	107.06	111.29
3	C	2	NAG	O5-C1-C2	-2.65	107.10	111.29
2	B	8	MAN	O5-C5-C6	2.63	111.33	107.20
2	B	4	MAN	O6-C6-C5	-2.56	102.50	111.29
2	B	8	MAN	C1-O5-C5	2.52	115.61	112.19
2	B	8	MAN	C1-C2-C3	2.52	112.77	109.67
2	B	3	BMA	O5-C1-C2	2.49	114.61	110.77
2	B	1	NAG	C4-C3-C2	-2.42	107.47	111.02
2	B	2	NAG	C1-O5-C5	-2.40	108.94	112.19
2	B	2	NAG	O3-C3-C4	2.39	115.88	110.35
2	B	4	MAN	C6-C5-C4	-2.39	107.42	113.00
3	C	2	NAG	O5-C5-C6	2.36	110.90	107.20
3	C	1	NAG	O3-C3-C4	2.32	115.71	110.35
2	B	6	MAN	O3-C3-C4	-2.28	105.07	110.35
2	B	4	MAN	O2-C2-C3	2.17	114.48	110.14

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	NAG	C8-C7-N2	2.14	119.72	116.10
2	B	6	MAN	O5-C5-C6	-2.13	103.86	107.20
3	C	1	NAG	O7-C7-N2	-2.13	118.03	121.95
2	B	2	NAG	O6-C6-C5	-2.13	103.99	111.29
2	B	5	MAN	C6-C5-C4	-2.12	108.04	113.00

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	8	MAN	C4-C5-C6-O6
2	B	8	MAN	O5-C5-C6-O6
3	C	2	NAG	C4-C5-C6-O6
2	B	7	MAN	C4-C5-C6-O6
3	C	2	NAG	C3-C2-N2-C7

There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	NAG	A	512	1	14,14,15	0.85	0	17,19,21	1.48	3 (17%)
7	9WM	A	514[B]	-	14,20,20	0.94	0	18,28,28	1.97	3 (16%)
8	9VP	A	515[A]	1	17,20,20	0.67	0	16,28,28	1.51	2 (12%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	9SG	A	501	-	15,21,21	1.45	1 (6%)	17,31,31	1.74	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	A	512	1	-	0/6/23/26	0/1/1/1
7	9WM	A	514[B]	-	-	0/14/34/34	0/0/1/1
8	9VP	A	515[A]	1	-	2/14/34/34	0/1/1/1
4	9SG	A	501	-	-	4/14/39/39	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	501	9SG	C4-C3	4.85	1.56	1.51

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	514[B]	9WM	C4-C5-N5	-4.74	103.54	110.86
7	A	514[B]	9WM	F1-C3-C4	-4.26	105.24	108.78
7	A	514[B]	9WM	C4-C3-C2	4.16	118.42	111.80
4	A	501	9SG	O6-C6-C7	-4.04	101.05	107.29
8	A	515[A]	9VP	C4-C5-N5	-3.79	105.00	110.86
8	A	515[A]	9VP	C6-C5-N5	-3.24	104.57	110.58
5	A	512	NAG	C1-C2-N2	-3.11	105.17	110.49
5	A	512	NAG	C3-C4-C5	-3.05	104.80	110.24
4	A	501	9SG	O10-C10-C11	-2.85	116.76	122.06
4	A	501	9SG	O6-C6-C5	2.14	111.93	109.62
4	A	501	9SG	O8-C8-C7	2.06	114.11	109.10
5	A	512	NAG	C4-C3-C2	2.04	114.01	111.02

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	501	9SG	O7-C7-C8-O8
4	A	501	9SG	C6-C7-C8-O8

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
8	A	515[A]	9VP	C7-C8-C9-O9
4	A	501	9SG	O7-C7-C8-C9
8	A	515[A]	9VP	O8-C8-C9-O9
4	A	501	9SG	C6-C7-C8-C9

There are no ring outliers.


2 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	514[B]	9WM	2	0
8	A	515[A]	9VP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## Ligand 9VP A 515 (A)





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	388/388 (100%)	-0.21	6 (1%) 73 76	15, 21, 33, 53	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	438	TRP	2.9
1	A	296	TRP	2.9
1	A	248	THR	2.8
1	A	83	ASP	2.7
1	A	82[A]	ARG	2.6
1	A	345	ASN	2.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	NAG	C	2	14/15	0.76	0.36	51,60,70,75	0
2	MAN	B	8	11/12	0.82	0.35	55,62,66,73	0
3	NAG	C	1	14/15	0.93	0.21	29,34,40,51	0
2	NAG	B	1	14/15	0.95	0.10	19,24,34,37	0
2	MAN	B	7	11/12	0.95	0.18	30,33,41,41	0
2	MAN	B	5	11/12	0.95	0.12	24,26,29,30	0

*Continued on next page...*


*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NAG	B	2	14/15	0.96	0.07	19,22,30,31	0
2	MAN	B	4	11/12	0.97	0.07	23,26,34,39	0
2	MAN	B	6	11/12	0.98	0.09	20,24,27,28	0
2	BMA	B	3	11/12	0.98	0.09	22,24,26,27	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.







## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	9SG	A	501	21/21	0.85	0.30	42,55,62,69	0
5	NAG	A	512	14/15	0.87	0.31	48,58,66,70	0
7	9WM	A	514[B]	20/20	0.95	0.12	19,25,29,45	20
8	9VP	A	515[A]	20/20	0.96	0.12	13,13,14,15	20
6	CA	A	513	1/1	0.99	0.05	26,26,26,26	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around 9SG A 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around 9WM A 514 (B):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around 9VP A 515 (A):**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.