



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 8, 2020 – 09:23 PM BST

PDB ID : 5W6T  
Title : Crystal structure of the A/Puerto Rico/8/1934 (H1N1) influenza virus hemagglutinin in complex with cyclic peptide CP151070 (P7)  
Authors : Wilson, I.A.; Kadam, R.U.  
Deposited on : 2017-06-16  
Resolution : 2.59 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	326	<div> <div style="width: 100%; height: 10px; background-color: red;"></div> <div style="width: 83%; height: 10px; background-color: green;"></div> <div style="width: 15%; height: 10px; background-color: yellow;"></div> <div style="width: 5%; height: 10px; background-color: orange;"></div> <div style="width: 5%; height: 10px; background-color: grey;"></div> </div>
2	B	176	<div> <div style="width: 7%; height: 10px; background-color: red;"></div> <div style="width: 87%; height: 10px; background-color: green;"></div> <div style="width: 10%; height: 10px; background-color: yellow;"></div> <div style="width: 5%; height: 10px; background-color: orange;"></div> <div style="width: 5%; height: 10px; background-color: grey;"></div> </div>
3	F	12	<div> <div style="width: 25%; height: 10px; background-color: green;"></div> <div style="width: 42%; height: 10px; background-color: yellow;"></div> <div style="width: 33%; height: 10px; background-color: orange;"></div> </div>
4	C	2	<div> <div style="width: 100%; height: 10px; background-color: green;"></div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	GOL	A	406	-	-	X	-

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 4174 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemagglutinin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	322	Total	C	N	O	S	0	0	0
			2542	1603	443	483	13			


- Molecule 2 is a protein called Hemagglutinin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	171	Total	C	N	O	S	0	0	0
			1380	866	235	272	7			

- Molecule 3 is a protein called ACE-PH8-ORN-MLE-GLU-TYR-ZCL-GLU-TRP-LEU-SER-9WV.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	F	12	Total	C	Cl	N	O	0	0	0
			115	79	2	15	19			

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	C	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	1	Total 1	Cl 1	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	48	Total 48	O 48	0	0
8	B	19	Total 19	O 19	0	0
8	F	1	Total 1	O 1	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Hemagglutinin




- Molecule 2: Hemagglutinin



- Molecule 3: ACE-PH8-ORN-MLE-GLU-TYR-ZCL-GLU-TRP-LEU-SER-9WV



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 21 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	164.33Å 164.33Å 164.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.08 – 2.59 41.08 – 2.59	Depositor EDS
% Data completeness (in resolution range)	92.2 (41.08-2.59) 92.2 (41.08-2.59)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.12	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.50 (at 2.58Å)	Xtriage
Refinement program	PHENIX (1.10_2155: ???)	Depositor
R, $R_{free}$	0.219 , 0.256 0.220 , 0.257	Depositor DCC
$R_{free}$ test set	1072 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	55.4	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 39.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.026 for -l,-k,-h	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	4174	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.02% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NAG, ACE, 9WV, ZCL, CL, ORN, PH8, MLE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.52	0/2606	0.60	0/3544
2	B	0.46	0/1407	0.54	0/1891
3	F	2.68	8/59 (13.6%)	1.63	1/78 (1.3%)
All	All	0.59	8/4072 (0.2%)	0.61	1/5513 (0.0%)

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	6	TYR	CB-CG	-7.53	1.40	1.51
3	F	9	TRP	CD2-CE2	-6.52	1.33	1.41
3	F	9	TRP	CG-CD2	-6.10	1.33	1.43
3	F	10	LEU	CA-C	-6.08	1.37	1.52
3	F	11	SER	CA-C	-5.74	1.38	1.52
3	F	5	GLU	CA-C	-5.50	1.38	1.52
3	F	9	TRP	CA-C	-5.42	1.38	1.52
3	F	6	TYR	CA-C	-5.18	1.39	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	9	TRP	CD1-CG-CD2	5.67	110.83	106.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2542	0	2469	36	0
2	B	1380	0	1310	14	0
3	F	115	0	88	3	0
4	C	28	0	25	0	0
5	A	28	0	26	0	0
6	A	12	0	15	7	0
7	B	1	0	0	0	0
8	A	48	0	0	1	0
8	B	19	0	0	0	0
8	F	1	0	0	0	0
All	All	4174	0	3933	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:50:ASN:O	2:B:54:THR:CG2	2.21	0.88
2:B:50:ASN:O	2:B:54:THR:HG23	1.72	0.86
1:A:73:ASN:ND2	6:A:406:GOL:O1	2.07	0.86
1:A:73:ASN:HD21	6:A:406:GOL:HO1	1.24	0.84
1:A:76:CYS:O	1:A:149:ARG:NH2	2.14	0.80
1:A:37:THR:OG1	1:A:38:HIS:ND1	2.15	0.76
1:A:37:THR:HG1	1:A:38:HIS:CE1	2.10	0.68
1:A:73:ASN:ND2	6:A:406:GOL:HO1	1.93	0.64
2:B:50:ASN:O	2:B:54:THR:HG22	1.97	0.64
1:A:38:HIS:CE1	2:B:21:TRP:HE1	2.17	0.62
1:A:288:ILE:HG22	1:A:289:ASN:O	2.01	0.61
1:A:95(A):GLY:H	6:A:406:GOL:H31	1.66	0.60
3:F:11:SER:O	3:F:12:9WV:N	2.36	0.59
1:A:71:LEU:O	1:A:148:TYR:HB3	2.07	0.55
1:A:103:ILE:HG13	1:A:233:TYR:CE2	2.42	0.54
1:A:83:ARG:NH1	8:A:503:HOH:O	2.40	0.53
1:A:43:LEU:HB2	1:A:314:LEU:HB2	1.92	0.52
1:A:92:ASN:OD1	1:A:92:ASN:N	2.42	0.52
1:A:301:THR:HB	1:A:305:CYS:SG	2.51	0.51
1:A:53:ARG:NH1	1:A:55:GLY:O	2.44	0.50
1:A:95:ASN:HD22	6:A:406:GOL:H12	1.75	0.50

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:117:SER:HB3	1:A:261:SER:OG	2.12	0.49
1:A:211:ARG:HG2	1:A:213:PHE:CZ	2.49	0.48
1:A:95:ASN:HA	6:A:406:GOL:H2	1.95	0.48
1:A:74:PRO:HB2	1:A:141:HIS:HB2	1.95	0.47
2:B:17:MET:SD	2:B:23:GLY:HA3	2.54	0.47
3:F:6:TYR:H	3:F:10:LEU:HD12	1.79	0.47
1:A:125(A):LYS:HE2	1:A:152:LEU:HD11	1.96	0.46
1:A:66:ILE:HG12	1:A:89:GLU:OE1	2.15	0.46
2:B:119:TYR:CE1	2:B:136:GLY:HA2	2.51	0.46
1:A:125(B):GLU:O	1:A:125(C):SER:HB2	2.16	0.45
2:B:49:THR:OG1	3:F:4:MLE:HN2	2.17	0.45
2:B:97:GLU:O	2:B:101:LEU:HG	2.16	0.45
1:A:15:ILE:HD11	2:B:122:VAL:HG21	1.99	0.44
1:A:316:MET:HE1	2:B:55:VAL:HG11	2.01	0.43
2:B:163:SER:H	2:B:163:SER:HG	1.54	0.43
1:A:202:VAL:HG13	1:A:247:ALA:HB2	2.00	0.43
1:A:89:GLU:O	1:A:269:THR:HA	2.19	0.43
2:B:170:ARG:O	2:B:171:GLU:HG2	2.19	0.42
1:A:220:ARG:NH1	1:A:228:GLY:HA2	2.34	0.42
1:A:98:TYR:O	1:A:232:TYR:OH	2.30	0.42
1:A:71:LEU:HD22	1:A:151:LEU:HD11	2.01	0.42
1:A:24:ASP:O	1:A:35:THR:HA	2.20	0.41
1:A:97:CYS:HA	6:A:406:GOL:H32	2.03	0.41
1:A:74:PRO:HG3	1:A:147:PHE:O	2.21	0.41
2:B:4:GLY:O	2:B:8:GLY:HA3	2.21	0.41
1:A:28:THR:HG22	2:B:104:ASN:HB3	2.03	0.41
1:A:37:THR:OG1	1:A:38:HIS:CE1	2.67	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	320/326 (98%)	307 (96%)	11 (3%)	2 (1%)	25	47
2	B	169/176 (96%)	163 (96%)	6 (4%)	0	100	100
3	F	5/12 (42%)	5 (100%)	0	0	100	100
All	All	494/514 (96%)	475 (96%)	17 (3%)	2 (0%)	34	57

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	125(C)	SER
1	A	125(A)	LYS

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	285/289 (99%)	279 (98%)	6 (2%)	53	77
2	B	147/151 (97%)	146 (99%)	1 (1%)	84	94
3	F	6/6 (100%)	6 (100%)	0	100	100
All	All	438/446 (98%)	431 (98%)	7 (2%)	62	82

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	SER
1	A	102	PHE
1	A	156	GLU
1	A	261	SER
1	A	290	SER
1	A	302	ILE
2	B	54	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	95	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

5 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	PH8	F	2	3	12,13,14	1.62	2 (16%)	10,15,17	1.24	1 (10%)
3	ZCL	F	7	3	12,13,14	1.46	1 (8%)	14,17,19	0.98	1 (7%)
3	MLE	F	4	3	7,8,9	0.56	0	6,9,11	1.46	2 (33%)
3	ORN	F	3	3	6,7,8	0.43	0	2,7,9	0.59	0
3	9WV	F	12	3	8,10,11	0.78	0	10,11,13	0.97	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PH8	F	2	3	-	3/7/8/10	0/1/1/1
3	ZCL	F	7	3	-	0/5/6/8	0/1/1/1
3	MLE	F	4	3	-	0/5/8/10	-
3	ORN	F	3	3	-	0/5/6/8	-
3	9WV	F	12	3	-	4/10/11/13	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	2	PH8	CJ-CG	-4.97	1.37	1.51
3	F	7	ZCL	CB-CG	-4.88	1.39	1.51
3	F	2	PH8	CB-CA	-2.17	1.50	1.53

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	4	MLE	CN-N-CA	2.42	121.18	113.64
3	F	2	PH8	CI-CJ-CG	-2.34	104.86	113.68
3	F	7	ZCL	CE2-CZ-CLZ	2.01	122.44	118.41
3	F	4	MLE	CG-CB-CA	-2.00	110.34	115.34

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	F	2	PH8	C-CA-CB-CI
3	F	2	PH8	N-CA-CB-CI
3	F	12	9WV	O-C-CA-CB
3	F	12	9WV	CEH-CEG-N-CA
3	F	12	9WV	OEK-CEG-N-CA
3	F	12	9WV	CEG-CEH-CEI-NEJ
3	F	2	PH8	CA-CB-CI-CJ

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	4	MLE	1	0
3	F	12	9WV	1	0

## 5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	NAG	C	1	1,4	14,14,15	0.29	0	17,19,21	0.58	0
4	NAG	C	2	4	14,14,15	0.28	0	17,19,21	0.59	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	C	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	C	2	4	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.6 Ligand geometry

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	GOL	A	406	1	5,5,5	0.36	0	5,5,5	0.78	0
6	GOL	A	405	-	5,5,5	0.31	0	5,5,5	0.61	0
5	NAG	A	404	1	14,14,15	0.52	0	17,19,21	0.99	1 (5%)
5	NAG	A	403	1	14,14,15	0.42	0	17,19,21	0.59	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	A	406	1	-	1/4/4/4	-
6	GOL	A	405	-	-	4/4/4/4	-
5	NAG	A	404	1	-	2/6/23/26	0/1/1/1
5	NAG	A	403	1	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	404	NAG	C1-O5-C5	3.54	116.99	112.19
5	A	403	NAG	C1-O5-C5	2.04	114.95	112.19

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	405	GOL	O1-C1-C2-C3
5	A	404	NAG	C4-C5-C6-O6
5	A	404	NAG	O5-C5-C6-O6
5	A	403	NAG	C4-C5-C6-O6
6	A	406	GOL	O1-C1-C2-C3
6	A	405	GOL	C1-C2-C3-O3
6	A	405	GOL	O1-C1-C2-O2
5	A	403	NAG	O5-C5-C6-O6
6	A	405	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	406	GOL	7	0

## 5.7 Other polymers

There are no such residues in this entry.



## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	322/326 (98%)	-0.17	3 (0%) 84 82	25, 43, 77, 97	0
2	B	171/176 (97%)	0.49	13 (7%) 13 10	25, 62, 104, 118	0
3	F	6/12 (50%)	0.24	0 100 100	64, 73, 83, 87	0
All	All	499/514 (97%)	0.06	16 (3%) 47 40	25, 48, 97, 118	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	168	LEU	4.1
1	A	325	SER	4.1
2	B	141	TYR	3.5
2	B	65	ALA	3.1
2	B	62	GLN	2.8
1	A	92	ASN	2.8
2	B	164	GLU	2.6
2	B	61	ILE	2.6
2	B	162	TYR	2.5
2	B	157	TYR	2.5
2	B	166	SER	2.5
2	B	138	PHE	2.4
2	B	158	ASP	2.4
1	A	196	GLN	2.3
2	B	29	GLU	2.1
2	B	147	GLU	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column

labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	9WV	F	12	11/12	0.85	0.22	93,96,102,102	0
3	ORN	F	3	8/9	0.91	0.17	54,62,80,93	0
3	ZCL	F	7	13/14	0.92	0.22	62,65,71,89	0
3	PH8	F	2	13/14	0.94	0.17	53,58,69,73	0
3	MLE	F	4	9/10	0.94	0.21	45,58,65,71	0

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	NAG	C	1	14/15	0.79	0.29	99,114,127,136	0
4	NAG	C	2	14/15	0.89	0.27	109,138,152,159	0

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	NAG	A	404	14/15	0.70	0.37	114,132,137,140	0
5	NAG	A	403	14/15	0.79	0.34	99,116,122,124	0
6	GOL	A	406	6/6	0.80	0.34	51,57,61,77	0
6	GOL	A	405	6/6	0.90	0.17	41,54,57,66	0
7	CL	B	201	1/1	0.99	0.35	32,32,32,32	1

### 6.5 Other polymers [i](#)

There are no such residues in this entry.