



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 21, 2022 – 12:04 PM JST

PDB ID : 7W7X  
Title : The crystal structure of human abl1 kinase domain in complex with ABL1-A11  
Authors : Zhu, C.; Zhang, Z.M.  
Deposited on : 2021-12-06  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.27  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.27


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	272	<div> <div>4%</div> <div> <div></div> <div>89%</div> <div>8%</div> <div>..</div> </div> </div>
1	B	272	<div> <div>6%</div> <div> <div></div> <div>91%</div> <div>6%</div> <div>.</div> </div> </div>

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 4766 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tyrosine-protein kinase ABL1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	267	Total	C	N	O	P	S	0	0	0
			2152	1385	349	400	1	17			
1	B	264	Total	C	N	O	P	S	0	0	0
			2094	1351	338	388	1	16			

- Molecule 2 is 5-[5-(dimethylcarbamoyl)pyridin-3-yl]-3-(5-fluorosulfonyloxy-2-methoxy-phenyl)-1H-pyrrolo[2,3-b]pyridine (three-letter code: 8DW) (formula: C<sub>22</sub>H<sub>19</sub>FN<sub>4</sub>O<sub>5</sub>S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	S	0	0
			33	22	1	4	5	1		
2	B	1	Total	C	F	N	O	S	0	0
			33	22	1	4	5	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	258	Total 258	O 258	0	0
3	B	196	Total 196	O 196	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosine-protein kinase ABL1



- Molecule 1: Tyrosine-protein kinase ABL1



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 2 <sub>1</sub> 2 <sub>1</sub> 2 <sub>1</sub>	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	56.41Å 104.88Å 132.91Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	27.24 – 2.00 27.24 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.9 (27.24-2.00) 99.9 (27.24-2.00)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.03 (at 1.99Å)	Xtriage
Refinement program	PHENIX 1.9_1692+SVN	Depositor
R, $R_{free}$	0.168 , 0.195 0.170 , 0.194	Depositor DCC
$R_{free}$ test set	1987 reflections (3.68%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	28.7	Xtriage
Anisotropy	0.464	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 58.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4766	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.68% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PTR, 8DW

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.42	0/2189	0.58	0/2964
1	B	0.38	0/2130	0.53	0/2891
All	All	0.40	0/4319	0.56	0/5855

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2152	0	2074	16	0
1	B	2094	0	1993	13	0
2	A	33	0	0	0	0
2	B	33	0	0	0	0
3	A	258	0	0	6	1
3	B	196	0	0	6	1
All	All	4766	0	4067	28	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:351:MET:SD	3:A:896:HOH:O	2.15	1.04
1:B:351:MET:SD	3:B:839:HOH:O	2.26	0.93
1:A:319:THR:OG1	3:A:701:HOH:O	1.95	0.83
1:A:304:VAL:HG13	1:A:306:THR:HG23	1.69	0.72
1:A:257:TYR:OH	3:A:703:HOH:O	2.07	0.69
1:B:393:PTR:O3P	3:B:701:HOH:O	2.14	0.65
1:A:450:GLU:OE1	3:A:704:HOH:O	2.17	0.60
1:B:279:GLU:N	3:B:708:HOH:O	2.35	0.59
1:B:279:GLU:N	3:B:709:HOH:O	2.36	0.57
1:A:351:MET:HE3	1:A:379:VAL:HG21	1.88	0.56
1:B:305:CYS:HB2	1:B:312:TYR:HB2	1.89	0.54
1:B:356:LYS:HG3	3:B:745:HOH:O	2.10	0.51
1:B:389:THR:HG23	1:B:390:GLY:N	2.25	0.51
1:A:237:MET:HE3	1:A:303:GLY:HA3	1.93	0.51
1:B:237:MET:SD	1:B:303:GLY:HA3	2.54	0.48
1:A:278:MET:N	3:A:713:HOH:O	2.47	0.47
1:B:263:LYS:HD3	1:B:264:TYR:CE1	2.51	0.46
1:B:390:GLY:N	3:B:712:HOH:O	2.49	0.45
1:A:386:ARG:HD2	1:B:500:SER:C	2.37	0.45
1:A:404:LYS:HE3	1:A:445:LEU:HD23	1.99	0.44
1:A:305:CYS:HB2	1:A:312:TYR:HB2	1.98	0.44
1:A:275:GLU:HA	3:A:708:HOH:O	2.18	0.44
1:B:241:ASP:N	1:B:241:ASP:OD1	2.50	0.43
1:B:351:MET:HE3	1:B:379:VAL:HG21	2.01	0.43
1:A:356:LYS:HB2	1:A:356:LYS:HE2	1.68	0.43
1:A:361:HIS:O	1:A:362:ARG:HB2	2.20	0.42
1:A:248:LEU:HD11	1:A:317:PHE:HE1	1.84	0.42
1:A:263:LYS:HD2	1:A:264:TYR:CE2	2.55	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:958:HOH:O	3:B:885:HOH:O[4_754]	1.81	0.39



## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	262/272 (96%)	256 (98%)	6 (2%)	0	100	100
1	B	259/272 (95%)	253 (98%)	6 (2%)	0	100	100
All	All	521/544 (96%)	509 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	225/237 (95%)	222 (99%)	3 (1%)	69	74
1	B	214/237 (90%)	212 (99%)	2 (1%)	78	83
All	All	439/474 (93%)	434 (99%)	5 (1%)	73	78

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	248	LEU
1	A	273	LEU
1	A	304	VAL
1	B	273	LEU
1	B	374	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	PTR	A	393	1	15,16,17	1.31	1 (6%)	19,22,24	1.00	3 (15%)
1	PTR	B	393	1	15,16,17	1.36	1 (6%)	19,22,24	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PTR	A	393	1	-	0/10/11/13	0/1/1/1
1	PTR	B	393	1	-	0/10/11/13	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	393	PTR	OH-CZ	-4.34	1.30	1.40
1	A	393	PTR	OH-CZ	-4.30	1.30	1.40

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	393	PTR	P-OH-CZ	2.38	131.38	123.75
1	A	393	PTR	O2P-P-OH	2.07	111.71	105.24
1	A	393	PTR	OH-CZ-CE2	2.00	125.19	119.23

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	393	PTR	1	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	8DW	A	601	-	34,36,36	2.37	7 (20%)	41,53,53	2.41	14 (34%)
2	8DW	B	601	-	34,36,36	2.39	8 (23%)	41,53,53	2.07	11 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	8DW	A	601	-	-	3/22/23/23	0/4/4/4
2	8DW	B	601	-	-	2/22/23/23	0/4/4/4

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	8DW	CAG-NAH	7.63	1.44	1.34
2	B	601	8DW	CAG-NAH	7.41	1.44	1.34

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	8DW	OBG-SBD	6.96	1.52	1.40
2	B	601	8DW	OBG-SBD	6.93	1.52	1.40
2	A	601	8DW	OBF-SBD	5.36	1.49	1.40
2	B	601	8DW	OBF-SBD	5.35	1.49	1.40
2	A	601	8DW	OBA-CAZ	3.91	1.43	1.37
2	B	601	8DW	OBA-CAZ	3.77	1.43	1.37
2	B	601	8DW	CAA-CAG	3.61	1.56	1.50
2	B	601	8DW	CAT-CAU	2.82	1.54	1.50
2	A	601	8DW	CAT-CAU	2.56	1.53	1.50
2	A	601	8DW	CAA-CAG	2.49	1.54	1.50
2	B	601	8DW	CAE-CAL	2.20	1.54	1.49
2	B	601	8DW	CAS-NAR	-2.14	1.32	1.36
2	A	601	8DW	CAQ-CAL	2.02	1.43	1.38

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	8DW	CAL-CAM-NAN	-6.40	119.63	125.55
2	B	601	8DW	CAL-CAM-NAN	-6.08	119.92	125.55
2	B	601	8DW	CAM-NAN-CAO	5.94	122.66	116.69
2	A	601	8DW	CAE-CAF-CAA	-5.29	115.36	121.09
2	A	601	8DW	CAA-CAG-NAH	4.69	124.24	119.00
2	A	601	8DW	CAM-NAN-CAO	4.66	121.38	116.69
2	A	601	8DW	OBF-SBD-OBG	-4.31	108.82	120.92
2	B	601	8DW	OBF-SBD-OBG	-4.21	109.08	120.92
2	A	601	8DW	CAF-CAE-CAD	3.90	120.98	117.11
2	A	601	8DW	CAQ-CAL-CAM	3.81	121.56	116.24
2	A	601	8DW	OBA-CAZ-CAU	3.54	121.55	116.26
2	A	601	8DW	OBA-CAZ-CAY	-3.52	118.34	124.37
2	B	601	8DW	OBA-CAZ-CAY	-3.03	119.18	124.37
2	B	601	8DW	CAA-CAG-NAH	3.01	122.36	119.00
2	B	601	8DW	CAB-NAC-CAD	3.01	121.59	117.48
2	B	601	8DW	OBA-CAZ-CAU	2.99	120.72	116.26
2	B	601	8DW	CAF-CAE-CAD	2.89	119.97	117.11
2	A	601	8DW	CAW-OBC-SBD	-2.87	116.19	121.66
2	B	601	8DW	CAE-CAF-CAA	-2.69	118.18	121.09
2	A	601	8DW	OAK-CAG-CAA	-2.66	115.06	120.23
2	A	601	8DW	CAL-CAQ-CAP	-2.61	116.59	122.30
2	B	601	8DW	CAQ-CAL-CAM	2.44	119.64	116.24
2	A	601	8DW	CBB-OBA-CAZ	-2.29	114.07	117.53
2	A	601	8DW	CAF-CAA-CAB	2.26	120.28	117.95
2	B	601	8DW	CAW-OBC-SBD	-2.23	117.40	121.66

There are no chirality outliers.


All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	601	8DW	CAP-CAT-CAU-CAZ
2	B	601	8DW	CAP-CAT-CAU-CAZ
2	A	601	8DW	CAV-CAW-OBC-SBD
2	B	601	8DW	CAV-CAW-OBC-SBD
2	A	601	8DW	CAY-CAZ-OBA-CBB

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	266/272 (97%)	-0.07	10 (3%) 40 39	18, 30, 59, 85	0
1	B	263/272 (96%)	0.17	16 (6%) 21 20	24, 38, 78, 93	0
All	All	529/544 (97%)	0.05	26 (4%) 29 28	18, 34, 71, 93	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	232	TYR	7.0
1	A	232	TYR	5.1
1	B	311	PHE	4.8
1	A	250	GLY	4.3
1	A	249	GLY	3.6
1	B	264	TYR	3.5
1	B	309	PRO	3.5
1	B	310	PRO	3.5
1	B	398	GLY	3.5
1	B	273	LEU	3.1
1	B	261	TRP	3.0
1	A	275	GLU	3.0
1	B	251	GLY	2.9
1	B	243	THR	2.7
1	B	263	LYS	2.6
1	B	265	SER	2.5
1	B	307	ARG	2.5
1	B	399	ALA	2.4
1	B	382	PHE	2.4
1	A	364	LEU	2.3
1	A	428	LEU	2.2
1	A	424	ALA	2.2
1	A	253	TYR	2.1
1	A	274	LYS	2.1

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	250	GLY	2.1
1	A	427	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	PTR	B	393	16/17	0.94	0.11	25,34,76,80	0
1	PTR	A	393	16/17	0.97	0.09	27,34,65,69	0

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	8DW	A	601	33/33	0.93	0.14	22,32,58,68	0
2	8DW	B	601	33/33	0.96	0.10	29,40,82,85	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around 8DW A 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around 8DW B 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.