



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 14, 2020 – 09:45 am BST

PDB ID : 2W8Z  
Title : Geobacillus stearothermophilus 6-phosphogluconate dehydrogenase with  
bound 6- phosphogluconate  
Authors : Cameron, S.; Hunter, W.N.  
Deposited on : 2009-01-20  
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

## 2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 7886 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 6-PHOSPHOGLUCONATE DEHYDROGENASE, DECARBOXYLATING.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	468	Total	C	N	O	S	2	17	0
			3772	2413	646	699	14			
1	B	467	Total	C	N	O	S	0	11	0
			3704	2368	626	695	15			

- Molecule 2 is 6-PHOSPHOGLUCONIC ACID (three-letter code: 6PG) (formula:  $C_6H_{13}O_{10}P$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	O	P	0	0
			17	6	10	1		
2	B	1	Total	C	O	P	0	0
			17	6	10	1		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula:  $O_4S$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: C<sub>4</sub>H<sub>12</sub>NO<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	1
			16	8	2	6		
4	A	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	168	Total 168	O 168	0	0
5	B	179	Total 179	O 179	0	0

SEQUENCE-PLOTS INFOmissingINFO

### 3 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	67.03Å 119.97Å 142.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	71.43 – 2.30 71.41 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.9 (71.43-2.30) 99.9 (71.41-2.30)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.58 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.182 , 0.234 0.181 , 0.230	Depositor DCC
$R_{free}$ test set	2618 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.2	Xtriage
Anisotropy	0.234	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 39.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	7886	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.11% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 4 Model quality ⓘ

### 4.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, 6PG, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.69	2/3891 (0.1%)	0.69	2/5243 (0.0%)
1	B	0.59	0/3808	0.66	2/5142 (0.0%)
All	All	0.64	2/7699 (0.0%)	0.68	4/10385 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	308[A]	VAL	CB-CG1	-15.19	1.21	1.52
1	A	308[B]	VAL	CB-CG1	-15.19	1.21	1.52

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	308[A]	VAL	CA-CB-CG1	9.06	124.50	110.90
1	A	308[B]	VAL	CA-CB-CG1	9.06	124.50	110.90
1	B	415	ARG	NE-CZ-NH2	-6.58	117.01	120.30
1	B	202	LEU	CA-CB-CG	5.19	127.23	115.30

There are no chirality outliers.

There are no planarity outliers.

### 4.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3772	0	3824	68	0
1	B	3704	0	3718	50	0
2	A	17	0	10	4	0
2	B	17	0	10	3	0
3	A	5	0	0	0	0
4	A	24	0	36	11	0
5	A	168	0	0	12	0
5	B	179	0	0	8	0
All	All	7886	0	7598	114	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 114 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:353[B]:ARG:HD3	5:B:2126:HOH:O	1.31	1.25
1:B:359[B]:MET:SD	1:B:376:LYS:HD2	1.82	1.19
1:B:353[A]:ARG:HD2	5:B:2125:HOH:O	1.48	1.12
4:A:1472[B]:TRS:H22	5:A:2166:HOH:O	1.49	1.09
4:A:1472[B]:TRS:O3	5:A:2167:HOH:O	1.84	0.95

There are no symmetry-related clashes.

## 4.3 Torsion angles [i](#)

### 4.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	484/470 (103%)	468 (97%)	14 (3%)	2 (0%)	34	42
1	B	476/470 (101%)	460 (97%)	15 (3%)	1 (0%)	47	58
All	All	960/940 (102%)	928 (97%)	29 (3%)	3 (0%)	41	50

All (3) Ramachandran outliers are listed below:



Mol	Chain	Res	Type
1	A	371	PHE
1	A	3	LYS
1	B	137	GLY

#### 4.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	393/377 (104%)	371 (94%)	22 (6%)	21	29
1	B	385/377 (102%)	369 (96%)	16 (4%)	30	42
All	All	778/754 (103%)	740 (95%)	38 (5%)	24	35

5 of 38 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	271	LEU
1	A	376	LYS
1	B	374	LYS
1	A	297	VAL
1	B	34	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	317	HIS
1	B	186	ASN
1	A	452	HIS
1	A	145	GLN
1	B	86	GLN

#### 4.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 4.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 4.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 4.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SO4	A	1471	-	4,4,4	0.12	0	6,6,6	0.36	0
4	TRS	A	1472[A]	-	7,7,7	0.30	0	9,9,9	0.60	0
4	TRS	A	1474	-	7,7,7	0.52	0	9,9,9	0.65	0
2	6PG	A	1470	-	13,16,16	1.13	1 (7%)	19,23,23	1.83	3 (15%)
2	6PG	B	1469	-	13,16,16	1.02	0	19,23,23	1.75	3 (15%)
4	TRS	A	1472[B]	-	7,7,7	0.28	0	9,9,9	0.34	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	6PG	A	1470	-	-	11/18/22/22	-
2	6PG	B	1469	-	-	1/18/22/22	-
4	TRS	A	1472[B]	-	-	6/9/9/9	-
4	TRS	A	1472[A]	-	-	3/9/9/9	-
4	TRS	A	1474	-	-	5/9/9/9	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1470	6PG	C4-C3	2.04	1.57	1.53

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1470	6PG	C4-C3-C2	5.75	123.71	113.58
2	B	1469	6PG	C4-C3-C2	5.00	122.39	113.58
2	A	1470	6PG	C5-C4-C3	3.02	117.19	112.47
2	B	1469	6PG	O4-C4-C3	2.73	115.84	109.47
2	A	1470	6PG	O4-C4-C3	2.28	114.78	109.47

There are no chirality outliers.

5 of 26 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	1472[A]	TRS	N-C-C1-O1
4	A	1474	TRS	N-C-C1-O1
4	A	1474	TRS	C1-C-C3-O3
4	A	1474	TRS	C2-C-C3-O3
4	A	1474	TRS	N-C-C3-O3


There are no ring outliers.

5 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1472[A]	TRS	3	0
4	A	1474	TRS	4	0
2	A	1470	6PG	4	0
2	B	1469	6PG	3	0
4	A	1472[B]	TRS	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





#### 4.7 Other polymers [i](#)

There are no such residues in this entry.

#### 4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 5 Fit of model and data [i](#)

### 5.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	468/470 (99%)	-0.17	10 (2%) 63 70	13, 25, 44, 62	4 (0%)
1	B	467/470 (99%)	-0.25	9 (1%) 66 73	10, 23, 42, 65	1 (0%)
All	All	935/940 (99%)	-0.21	19 (2%) 65 71	10, 24, 44, 65	5 (0%)

The worst 5 of 19 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	244	GLU	5.3
1	A	2	ALA	4.5
1	B	463	PHE	4.4
1	B	468	LEU	4.4
1	A	243	GLU	4.3

### 5.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.


### 5.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	TRS	A	1474	8/8	0.80	0.19	36,37,37,37	0
2	6PG	A	1470	17/17	0.84	0.29	73,75,77,77	0
2	6PG	B	1469	17/17	0.87	0.18	40,49,56,57	0
3	SO4	A	1471	5/5	0.89	0.27	72,72,72,73	0
4	TRS	A	1472[A]	8/8	0.92	0.18	19,21,21,21	8
4	TRS	A	1472[B]	8/8	0.92	0.18	21,21,21,21	8

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 5.5 Other polymers [i](#)

There are no such residues in this entry.