



Full wwPDB X-ray Structure Validation Report ⓘ

May 27, 2020 – 06:29 pm BST

PDB ID : 1WAB
Title : PLATELET-ACTIVATING FACTOR ACETYLHYDROLASE
Authors : Ho, Y.S.; Swenson, L.; Derewenda, U.; Serre, L.; Wei, Y.; Dauter, Z.; Hattori, M.; Aoki, J.; Arai, H.; Adachi, T.; Inoue, K.; Derewenda, Z.S.
Deposited on : 1996-10-30
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	232	

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 1925 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PLATELET-ACTIVATING FACTOR ACETYLHYDROLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	212	Total	C	N	O	S	40	0	0
			1689	1066	321	297	5			

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	232	Total	O	0	0
			232	232		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: PLATELET-ACTIVATING FACTOR ACETYLHYDROLASE



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	81.26Å 81.26Å 72.60Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	8.50 – 1.70	Depositor
% Data completeness (in resolution range)	(Not available) (8.50-1.70)	Depositor
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	PROLSQ, REFMAC, X-PLOR	Depositor
R, R_{free}	0.195 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	1925	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.00	2/1734 (0.1%)	1.76	38/2358 (1.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	5

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	57	ILE	C-N	-14.25	1.01	1.34
1	A	61	LEU	C-N	7.37	1.50	1.34

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	153	ARG	CD-NE-CZ	24.10	157.34	123.60
1	A	160	ARG	CD-NE-CZ	15.38	145.14	123.60
1	A	214	ARG	CD-NE-CZ	15.20	144.87	123.60
1	A	83	ARG	NE-CZ-NH1	13.01	126.80	120.30
1	A	101	VAL	CB-CA-C	-11.45	89.65	111.40
1	A	57	ILE	C-N-CA	10.68	148.40	121.70
1	A	166	HIS	CA-CB-CG	10.17	130.89	113.60
1	A	75	ASP	CB-CG-OD2	-9.70	109.57	118.30
1	A	191	TYR	CB-CG-CD1	-9.65	115.21	121.00
1	A	214	ARG	NE-CZ-NH1	9.45	125.03	120.30
1	A	168	ARG	NE-CZ-NH1	-8.69	115.96	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	149	ARG	NE-CZ-NH2	8.66	124.63	120.30
1	A	57	ILE	CA-C-O	8.43	137.79	120.10
1	A	198	ARG	CD-NE-CZ	8.35	135.28	123.60
1	A	198	ARG	CG-CD-NE	7.53	127.61	111.80
1	A	168	ARG	NE-CZ-NH2	7.37	123.98	120.30
1	A	189	ASP	CB-CG-OD2	-7.28	111.75	118.30
1	A	59	ARG	CD-NE-CZ	6.72	133.01	123.60
1	A	61	LEU	C-N-CA	-6.67	105.01	121.70
1	A	83	ARG	NE-CZ-NH2	-6.64	116.98	120.30
1	A	30	PHE	CB-CG-CD2	-6.54	116.22	120.80
1	A	59	ARG	CG-CD-NE	6.35	125.13	111.80
1	A	93	ARG	NE-CZ-NH1	6.19	123.40	120.30
1	A	214	ARG	CG-CD-NE	6.08	124.56	111.80
1	A	33	ASP	CB-CG-OD2	-5.85	113.04	118.30
1	A	42	VAL	CA-CB-CG1	5.63	119.35	110.90
1	A	160	ARG	NE-CZ-NH1	5.63	123.11	120.30
1	A	29	ARG	NE-CZ-NH1	-5.62	117.49	120.30
1	A	101	VAL	CA-CB-CG2	5.61	119.32	110.90
1	A	85	GLU	CG-CD-OE2	-5.57	107.17	118.30
1	A	132	ARG	NE-CZ-NH2	-5.43	117.58	120.30
1	A	198	ARG	NE-CZ-NH1	5.42	123.01	120.30
1	A	171	PHE	CB-CG-CD1	-5.32	117.08	120.80
1	A	154	ARG	NE-CZ-NH1	5.32	122.96	120.30
1	A	206	ARG	CA-CB-CG	5.23	124.90	113.40
1	A	214	ARG	N-CA-CB	5.19	119.94	110.60
1	A	24	MET	CG-SD-CE	5.16	108.45	100.20
1	A	201	TYR	CB-CG-CD1	-5.01	118.00	121.00

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	133	VAL	Mainchain
1	A	169	ALA	Mainchain
1	A	215	LEU	Mainchain
1	A	55	CYS	Peptide
1	A	57	ILE	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1689	0	1668	31	0
2	A	4	0	3	1	0
3	A	232	0	0	11	3
All	All	1925	0	1671	31	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:153:ARG:NH1	3:A:496:HOH:O	1.96	0.98
1:A:150:GLU:OE1	3:A:460:HOH:O	1.80	0.96
1:A:180:HIS:HD2	3:A:515:HOH:O	1.63	0.81
1:A:77:THR:H	1:A:105:ASN:HD21	1.31	0.76
1:A:150:GLU:HG2	3:A:402:HOH:O	1.85	0.76
1:A:214:ARG:HB2	1:A:214:ARG:HH11	1.51	0.74
1:A:47:SER:HB3	2:A:300:ACT:C	2.21	0.70
1:A:5:GLU:O	3:A:511:HOH:O	2.10	0.69
1:A:36:ASP:HB3	3:A:412:HOH:O	1.94	0.67
1:A:168:ARG:HD3	3:A:476:HOH:O	1.98	0.64
1:A:198:ARG:NH1	3:A:507:HOH:O	2.38	0.56
1:A:132:ARG:HH11	1:A:168:ARG:HG2	1.70	0.56
1:A:146:ASN:HD22	1:A:148:LEU:H	1.54	0.55
1:A:63:SER:HB2	1:A:64:PRO:HD3	1.93	0.51
1:A:180:HIS:CD2	3:A:515:HOH:O	2.49	0.50
1:A:123:LEU:HD13	1:A:123:LEU:C	2.32	0.50
1:A:129:PRO:O	1:A:168:ARG:NH2	2.45	0.50
1:A:211:LEU:HA	1:A:214:ARG:HG3	1.94	0.50
1:A:24:MET:HG3	1:A:28:HIS:CE1	2.47	0.50
1:A:150:GLU:HB3	3:A:496:HOH:O	2.13	0.49
1:A:163:LEU:HA	1:A:166:HIS:CD2	2.49	0.48
1:A:214:ARG:CB	1:A:214:ARG:HH11	2.24	0.47
1:A:153:ARG:O	1:A:157:GLU:HG3	2.15	0.46
1:A:6:ASN:ND2	1:A:8:ALA:H	2.14	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:48:LEU:HD12	3:A:397:HOH:O	2.16	0.45
1:A:212:LEU:O	1:A:216:LEU:HB2	2.18	0.44
1:A:168:ARG:HB2	1:A:168:ARG:CZ	2.48	0.43
1:A:6:ASN:HD21	1:A:112:GLN:HB3	1.84	0.43
1:A:6:ASN:HD22	1:A:7:PRO:HD2	1.84	0.42
1:A:43:PHE:HE2	1:A:208:LEU:HD21	1.85	0.42
1:A:202:THR:N	1:A:203:PRO:HD2	2.36	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:370:HOH:O	3:A:430:HOH:O[5_555]	1.89	0.31
3:A:312:HOH:O	3:A:381:HOH:O[5_555]	2.08	0.12
3:A:443:HOH:O	3:A:469:HOH:O[5_555]	2.18	0.02

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	210/232 (90%)	200 (95%)	10 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	183/198 (92%)	172 (94%)	11 (6%)	19 6

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	ASN
1	A	55	CYS
1	A	59	ARG
1	A	61	LEU
1	A	101	VAL
1	A	150	GLU
1	A	163	LEU
1	A	198	ARG
1	A	206	ARG
1	A	212	LEU
1	A	214	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	6	ASN
1	A	15	GLN
1	A	105	ASN
1	A	122	GLN
1	A	143	GLN
1	A	146	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ACT	A	300	-	1,3,3	6.48	1 (100%)	0,3,3	0.00	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	300	ACT	CH3-C	6.48	1.57	1.48

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	300	ACT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	57:ILE	C	58:TRP	N	1.01

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.