



Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 01:38 am BST

PDB ID : 2WD0
Title : CRYSTAL STRUCTURE OF NONSYNDROMIC DEAFNESS (DFNB12)
ASSOCIATED MUTANT D124G OF MOUSE CADHERIN-23 EC1-2
Authors : Sotomayor, M.; Weihofen, W.; Gaudet, R.; Corey, D.P.
Deposited on : 2009-03-18
Resolution : 2.74 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.74 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1271 (2.76-2.72)
Clashscore	141614	1322 (2.76-2.72)
Ramachandran outliers	138981	1297 (2.76-2.72)
Sidechain outliers	138945	1298 (2.76-2.72)
RSRZ outliers	127900	1243 (2.76-2.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	214	<div> <div>%</div> <div> <div></div> <div>75%</div> <div>20%</div> <div>• •</div> </div> </div>
1	C	214	<div> <div>4%</div> <div> <div></div> <div>74%</div> <div>19%</div> <div>• •</div> </div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3339 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CADHERIN-23.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	208	Total	C	N	O	S	0	1	0
			1638	1043	271	323	1			
1	C	208	Total	C	N	O	S	0	1	0
			1638	1043	271	323	1			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	expression tag	UNP Q99PF4
A	102	GLY	ASP	engineered mutation	UNP Q99PF4
A	207	LEU	-	expression tag	UNP Q99PF4
A	208	GLU	-	expression tag	UNP Q99PF4
A	209	HIS	-	expression tag	UNP Q99PF4
A	210	HIS	-	expression tag	UNP Q99PF4
A	211	HIS	-	expression tag	UNP Q99PF4
A	212	HIS	-	expression tag	UNP Q99PF4
A	213	HIS	-	expression tag	UNP Q99PF4
A	214	HIS	-	expression tag	UNP Q99PF4
C	1	MET	-	expression tag	UNP Q99PF4
C	102	GLY	ASP	engineered mutation	UNP Q99PF4
C	207	LEU	-	expression tag	UNP Q99PF4
C	208	GLU	-	expression tag	UNP Q99PF4
C	209	HIS	-	expression tag	UNP Q99PF4
C	210	HIS	-	expression tag	UNP Q99PF4
C	211	HIS	-	expression tag	UNP Q99PF4
C	212	HIS	-	expression tag	UNP Q99PF4
C	213	HIS	-	expression tag	UNP Q99PF4
C	214	HIS	-	expression tag	UNP Q99PF4

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	3	Total Ca 3 3	0	0
2	C	3	Total Ca 3 3	0	0

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Na 1 1	0	0
3	C	1	Total Na 1 1	0	0

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	2	Total Cl 2 2	0	0
4	C	1	Total Cl 1 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	26	Total O 26 26	0	0
5	C	26	Total O 26 26	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: CADHERIN-23



• Molecule 1: CADHERIN-23



4 Data and refinement statistics

Property	Value	Source
Space group	P 42 21 2	Depositor
Cell constants a, b, c, α , β , γ	179.69 Å 179.69 Å 63.98 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.77 – 2.74 29.54 – 2.74	Depositor EDS
% Data completeness (in resolution range)	97.9 (29.77-2.74) 97.9 (29.54-2.74)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.93 (at 2.76 Å)	Xtriage
Refinement program	REFMAC 5.5.0070	Depositor
R, R_{free}	0.219 , 0.249 0.226 , 0.261	Depositor DCC
R_{free} test set	1384 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	42.9	Xtriage
Anisotropy	0.013	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 42.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	3339	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 20.31 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.9306e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, CA, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.55	0/1679	0.56	0/2301
1	C	0.57	0/1679	0.57	0/2301
All	All	0.56	0/3358	0.57	0/4602

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1638	0	1576	33	0
1	C	1638	0	1576	31	0
2	A	3	0	0	0	0
2	C	3	0	0	0	0
3	A	1	0	0	0	0
3	C	1	0	0	0	0
4	A	2	0	0	0	0
4	C	1	0	0	0	0
5	A	26	0	0	1	1
5	C	26	0	0	1	0
All	All	3339	0	3152	60	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:66[A]:TRP:CE3	1:C:66[A]:TRP:CE2	2.04	1.08
1:A:66[A]:TRP:CD2	1:C:66[A]:TRP:CE2	2.50	1.00
1:A:11:ASN:HD21	1:A:34:LEU:H	1.19	0.88
1:A:184:ASN:HD21	1:A:186:THR:HG23	1.45	0.80
1:A:130:PHE:HB3	1:A:165:VAL:CG1	2.15	0.76
1:C:171:LEU:HD13	1:C:204:ILE:HD12	1.69	0.73
1:C:130:PHE:HB3	1:C:165:VAL:HG13	1.70	0.72
1:C:122:ASN:HD22	1:C:122:ASN:C	1.93	0.72
1:A:73:ARG:NH1	1:A:74:GLU:OE1	2.23	0.72
1:C:11:ASN:HD21	1:C:34:LEU:H	1.35	0.72
1:A:66[A]:TRP:CD2	1:C:66[A]:TRP:CD2	2.68	0.71
1:A:7:PRO:O	1:A:92:ILE:HD12	1.93	0.68
1:A:186:THR:HG22	1:A:195:SER:CB	2.23	0.68
1:A:52:ALA:HB1	1:A:81:VAL:HG13	1.79	0.64
1:A:184:ASN:HD21	1:A:186:THR:CG2	2.12	0.62
1:A:186:THR:HG22	1:A:195:SER:HB2	1.83	0.60
1:A:186:THR:HG22	1:A:195:SER:HB3	1.84	0.60
1:A:11:ASN:ND2	1:A:34:LEU:H	1.94	0.60
1:A:66[A]:TRP:CE3	1:C:66[A]:TRP:NE1	2.66	0.57
1:C:73:ARG:NH1	1:C:74:GLU:OE1	2.38	0.57
1:C:77:SER:HA	1:C:100:VAL:HG13	1.87	0.55
1:C:17:TYR:CE1	1:C:99:GLN:HG3	2.44	0.53
1:A:184:ASN:ND2	1:A:186:THR:HG23	2.18	0.53
1:A:110:PHE:CD2	1:A:198:ALA:HB2	2.44	0.52
1:C:73:ARG:NH2	1:C:139:LEU:HD13	2.24	0.52
1:A:11:ASN:HD21	1:A:34:LEU:N	1.99	0.51
1:C:43:LEU:HA	1:C:86:SER:O	2.11	0.51
1:A:130:PHE:HB3	1:A:165:VAL:HG12	1.92	0.51
1:A:171:LEU:HD13	1:A:204:ILE:HD12	1.93	0.49
1:C:19:LEU:HD12	1:C:99:GLN:HB2	1.94	0.49
1:A:73:ARG:NE	1:A:139:LEU:HD22	2.28	0.49
1:C:184:ASN:HD21	1:C:186:THR:HB	1.77	0.49
1:C:110:PHE:CD2	1:C:198:ALA:HB2	2.48	0.49
1:C:122:ASN:ND2	1:C:122:ASN:C	2.62	0.48
1:A:26:VAL:HG13	5:A:2004:HOH:O	2.14	0.48
1:A:130:PHE:HB3	1:A:165:VAL:HG11	1.92	0.47
1:C:156:PHE:CE1	1:C:167:VAL:HG22	2.49	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:5:ARG:NH2	5:C:2001:HOH:O	2.48	0.47
1:A:159:ASP:OD1	1:A:162:ARG:N	2.44	0.46
1:A:20:ILE:HD12	1:A:98:ILE:HG21	1.98	0.46
1:C:122:ASN:HD22	1:C:123:THR:N	2.14	0.46
1:A:77:SER:HA	1:A:100:VAL:HG13	1.97	0.46
1:C:128:PRO:HA	1:C:166:THR:HG22	1.98	0.46
1:A:110:PHE:CE2	1:A:198:ALA:HB2	2.52	0.45
1:A:19:LEU:HD12	1:A:99:GLN:O	2.17	0.45
1:C:37:ASP:OD2	1:C:39:ASP:HB2	2.17	0.45
1:C:130:PHE:HB3	1:C:165:VAL:CG1	2.41	0.44
1:C:11:ASN:ND2	1:C:34:LEU:H	2.09	0.43
1:A:141:ALA:HB1	1:A:188:GLN:O	2.18	0.43
1:A:173:TYR:CD2	1:A:173:TYR:C	2.92	0.42
1:A:125:VAL:HG12	1:A:169:GLN:N	2.35	0.42
1:C:16:THR:OG1	1:C:17:TYR:N	2.53	0.42
1:A:73:ARG:NH2	1:A:139:LEU:HD13	2.34	0.42
1:A:156:PHE:CZ	1:A:202:ILE:HG21	2.54	0.41
1:C:173:TYR:CD1	1:C:206:ASP:HB2	2.56	0.41
1:C:52:ALA:CB	1:C:81:VAL:HG22	2.51	0.41
1:C:67:LEU:HD11	1:C:71:LEU:HG	2.02	0.41
1:C:120:PRO:HA	1:C:205:THR:HB	2.03	0.41
1:C:177:GLN:HE21	1:C:177:GLN:HB3	1.73	0.40
1:C:39:ASP:O	1:C:40:ASN:CB	2.69	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:2022:HOH:O	5:A:2022:HOH:O[8_555]	2.01	0.19

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	207/214 (97%)	202 (98%)	5 (2%)	0	100	100
1	C	207/214 (97%)	199 (96%)	7 (3%)	1 (0%)	29	48
All	All	414/428 (97%)	401 (97%)	12 (3%)	1 (0%)	47	69

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	208	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	185/190 (97%)	167 (90%)	18 (10%)	8	14
1	C	185/190 (97%)	170 (92%)	15 (8%)	11	21
All	All	370/380 (97%)	337 (91%)	33 (9%)	9	18

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	26	VAL
1	A	38	MET
1	A	73	ARG
1	A	100	VAL
1	A	109	THR
1	A	118	ARG
1	A	122	ASN
1	A	139	LEU
1	A	146	LEU
1	A	166	THR
1	A	176	THR
1	A	177	GLN
1	A	180	GLN
1	A	184	ASN
1	A	191	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	200	LEU
1	A	203	ILE
1	A	208	GLU
1	C	38	MET
1	C	50	GLU
1	C	75	THR
1	C	81	VAL
1	C	100	VAL
1	C	122	ASN
1	C	139	LEU
1	C	165	VAL
1	C	174	GLU
1	C	177	GLN
1	C	180	GLN
1	C	184	ASN
1	C	190	LYS
1	C	200	LEU
1	C	205	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	11	ASN
1	A	97	ASN
1	A	99	GLN
1	A	122	ASN
1	A	180	GLN
1	A	184	ASN
1	C	11	ASN
1	C	122	ASN
1	C	150	GLN
1	C	177	GLN
1	C	180	GLN
1	C	184	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 11 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	208/214 (97%)	-0.12	3 (1%) 75 80	14, 27, 43, 57	0
1	C	208/214 (97%)	0.04	8 (3%) 40 44	9, 27, 43, 68	0
All	All	416/428 (97%)	-0.04	11 (2%) 56 63	9, 27, 43, 68	0

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	208	GLU	3.7
1	A	208	GLU	3.3
1	C	174	GLU	3.2
1	C	175	VAL	3.1
1	C	2	GLN	3.1
1	C	209	HIS	2.9
1	A	209	HIS	2.7
1	C	190	LYS	2.4
1	C	91	VAL	2.3
1	C	90	GLY	2.1
1	A	207	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CA	C	504	1/1	0.93	0.04	30,30,30,30	0
3	NA	A	601	1/1	0.93	0.06	29,29,29,29	0
3	NA	C	601	1/1	0.93	0.07	26,26,26,26	0
2	CA	A	504	1/1	0.97	0.08	14,14,14,14	0
4	CL	A	702	1/1	0.97	0.07	52,52,52,52	0
2	CA	C	501	1/1	0.97	0.05	26,26,26,26	0
4	CL	C	701	1/1	0.98	0.07	41,41,41,41	0
4	CL	A	701	1/1	0.98	0.11	33,33,33,33	0
2	CA	A	502	1/1	0.99	0.07	19,19,19,19	0
2	CA	A	501	1/1	0.99	0.04	22,22,22,22	0
2	CA	C	503	1/1	0.99	0.04	21,21,21,21	0

6.5 Other polymers [i](#)

There are no such residues in this entry.