



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 13, 2020 – 11:02 AM BST

PDB ID : 2WEU
Title : Crystal structure of tryptophan 5-halogenase (PyrH) complex with substrate tryptophan
Authors : Leang, K.; Zhu, X.; Naismith, J.H.
Deposited on : 2009-04-01
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.14.4.dev1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.14.4.dev1


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	511	<div> <div>92%</div> <div>5% •</div> </div>
1	B	511	<div> <div>2%</div> <div>90%</div> <div>7% •</div> </div>
1	C	511	<div> <div>2%</div> <div>89%</div> <div>8% •</div> </div>
1	D	511	<div> <div>2%</div> <div>92%</div> <div>5% •</div> </div>

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 17519 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRYPTOPHAN 5-HALOGENASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	494	Total	C	N	O	S	0	2	0
			3987	2537	697	733	20			
1	B	499	Total	C	N	O	S	0	0	0
			4012	2549	706	738	19			
1	C	499	Total	C	N	O	S	0	1	0
			4013	2549	705	740	19			
1	D	498	Total	C	N	O	S	0	0	0
			3995	2540	699	737	19			

- Molecule 2 is TRYPTOPHAN (three-letter code: TRP) (formula: $C_{11}H_{12}N_2O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			15	11	2	2		
2	B	1	Total	C	N	O	0	0
			15	11	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	C	1	Total	C	N	O	0	0
			15	11	2	2		
2	D	1	Total	C	N	O	0	0
			15	11	2	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	393	Total	O	0	0
			393	393		
3	B	368	Total	O	0	0
			368	368		
3	C	365	Total	O	0	0
			365	365		
3	D	326	Total	O	0	0
			326	326		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: TRYPTOPHAN 5-HALOGENASE




• Molecule 1: TRYPTOPHAN 5-HALOGENASE




• Molecule 1: TRYPTOPHAN 5-HALOGENASE



• Molecule 1: TRYPTOPHAN 5-HALOGENASE





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	142.70Å 141.08Å 122.95Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	100.50 – 1.70 29.39 – 1.70	Depositor EDS
% Data completeness (in resolution range)	88.5 (100.50-1.70) 88.5 (29.39-1.70)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.94 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.5.0070	Depositor
R, R_{free}	0.215 , 0.248 0.224 , 0.258	Depositor DCC
R_{free} test set	12099 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	17.0	Xtriage
Anisotropy	0.081	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 24.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	0.240 for k,h,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	17519	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.94% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.61	0/4093	0.70	2/5548 (0.0%)
1	B	0.62	0/4119	0.69	2/5583 (0.0%)
1	C	0.62	0/4119	0.71	2/5583 (0.0%)
1	D	0.61	1/4101 (0.0%)	0.69	1/5559 (0.0%)
All	All	0.62	1/16432 (0.0%)	0.70	7/22273 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	367	GLN	CD-NE2	-5.43	1.19	1.32

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	471	ARG	NE-CZ-NH2	-7.38	116.61	120.30
1	A	471	ARG	NE-CZ-NH2	-7.14	116.73	120.30
1	C	260	ARG	NE-CZ-NH2	-6.92	116.84	120.30
1	B	260	ARG	NE-CZ-NH2	-6.14	117.23	120.30
1	C	260	ARG	NE-CZ-NH1	5.42	123.01	120.30
1	A	44	VAL	CG1-CB-CG2	5.38	119.50	110.90
1	B	309	LEU	CD1-CG-CD2	-5.20	94.89	110.50

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	113	GLY	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3987	0	3824	16	0
1	B	4012	0	3851	25	0
1	C	4013	0	3845	29	0
1	D	3995	0	3829	19	0
2	A	15	0	9	0	0
2	B	15	0	9	0	0
2	C	15	0	9	0	0
2	D	15	0	9	0	0
3	A	393	0	0	2	0
3	B	368	0	0	6	0
3	C	365	0	0	10	0
3	D	326	0	0	8	0
All	All	17519	0	15385	87	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (87) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:125:LEU:O	1:D:129:LEU:HD13	1.67	0.95
1:B:423:MET:HE2	1:B:427:LEU:HD23	1.56	0.87
1:A:125:LEU:O	1:A:129:LEU:HD13	1.76	0.84
1:C:423:MET:HE3	1:C:427:LEU:HD23	1.59	0.83
1:C:44:VAL:HG23	3:C:2007:HOH:O	1.78	0.82
1:A:405:LYS:HE3	1:A:423[A]:MET:HE3	1.68	0.75
1:D:402:LEU:HG	3:D:2232:HOH:O	1.87	0.74
1:D:325:ILE:HD13	3:D:2184:HOH:O	1.90	0.70
1:C:419:LYS:HE2	3:C:2295:HOH:O	1.90	0.70
1:B:423:MET:CE	1:B:427:LEU:HD23	2.21	0.70
1:C:267:ARG:NH2	1:C:282:THR:HG21	2.11	0.66

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:260:ARG:HD2	3:B:2213:HOH:O	1.95	0.66
1:B:12:THR:HG21	1:B:358:ILE:CD1	2.29	0.62
1:D:12:THR:HG21	1:D:358:ILE:CD1	2.29	0.62
1:C:423:MET:CE	1:C:427:LEU:HD23	2.29	0.62
1:B:128:ARG:NH2	1:B:494:ASP:OD1	2.27	0.60
1:B:370:LYS:HG3	3:B:2264:HOH:O	2.01	0.59
1:D:475:LEU:HG	3:D:2298:HOH:O	2.02	0.59
1:C:128:ARG:NH2	1:C:494:ASP:OD1	2.25	0.58
1:B:267:ARG:NH2	1:B:282:THR:HG21	2.19	0.57
1:A:12:THR:HG21	1:A:358:ILE:CD1	2.34	0.57
1:C:12:THR:HG21	1:C:358:ILE:CD1	2.34	0.57
1:A:480:ALA:HB3	1:A:481:PRO:HD3	1.88	0.56
1:C:44:VAL:CG2	3:C:2007:HOH:O	2.46	0.55
1:A:336:TRP:CG	1:A:341:VAL:HG22	2.42	0.54
1:B:260:ARG:CD	3:B:2213:HOH:O	2.52	0.54
1:D:242:GLN:NE2	3:D:2134:HOH:O	2.19	0.53
1:D:336:TRP:CG	1:D:341:VAL:HG22	2.43	0.53
1:B:298:GLU:HB3	3:B:2215:HOH:O	2.08	0.53
1:C:336:TRP:CG	1:C:341:VAL:HG22	2.43	0.53
1:D:480:ALA:HB3	1:D:481:PRO:HD3	1.91	0.52
1:D:405:LYS:HE3	1:D:423:MET:CE	2.39	0.52
1:C:334:ARG:NH1	3:C:2241:HOH:O	2.32	0.51
1:A:267:ARG:NH1	3:A:2194:HOH:O	2.44	0.50
1:C:402:LEU:HD11	1:C:460:ASN:OD1	2.12	0.49
1:C:39:VAL:HG11	3:C:2018:HOH:O	2.11	0.49
1:D:336:TRP:CD2	1:D:341:VAL:HG22	2.48	0.48
1:D:334:ARG:NH1	3:D:2191:HOH:O	2.45	0.48
1:C:336:TRP:CD2	1:C:341:VAL:HG22	2.48	0.48
1:B:249:PRO:HG3	1:B:421:ARG:CZ	2.44	0.47
1:A:440:ASP:CG	1:B:388:ARG:HH21	2.18	0.47
1:D:471:ARG:NH2	3:D:2299:HOH:O	2.43	0.47
1:C:81:GLU:HB3	1:C:89:TYR:HB3	1.96	0.47
1:A:267:ARG:NH2	1:A:282:THR:HG21	2.30	0.47
1:B:286:PHE:O	1:B:287:LYS:HB2	2.15	0.47
1:C:28:ARG:NH2	3:C:2013:HOH:O	2.48	0.47
1:A:88[A]:GLU:HA	1:A:88[A]:GLU:OE1	2.15	0.47
1:C:260:ARG:HD2	3:C:2211:HOH:O	2.15	0.46
1:B:325:ILE:HD13	3:B:2232:HOH:O	2.16	0.46
1:C:260:ARG:CD	3:C:2211:HOH:O	2.64	0.46
1:A:471:ARG:NH2	3:A:2364:HOH:O	2.48	0.46
1:C:249:PRO:HG3	1:C:421:ARG:CZ	2.44	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:125:LEU:O	1:B:129:LEU:HD13	2.16	0.45
1:D:397:LYS:CE	3:D:2229:HOH:O	2.65	0.44
1:A:367:GLN:HE22	1:A:370:LYS:HD2	1.83	0.44
1:B:402:LEU:HD11	1:B:460:ASN:OD1	2.18	0.44
1:B:81:GLU:HB3	1:B:89:TYR:HB3	2.00	0.44
1:C:36:SER:OG	1:C:194:ASP:OD1	2.26	0.44
1:D:267:ARG:NH2	1:D:282:THR:HG21	2.33	0.44
1:D:397:LYS:HE2	3:D:2229:HOH:O	2.17	0.44
1:C:286:PHE:O	1:C:287:LYS:HB2	2.18	0.43
1:C:423:MET:HE1	1:C:431:LEU:HD11	2.00	0.43
1:B:44:VAL:HG22	1:B:45:GLY:H	1.83	0.43
1:A:249:PRO:HG3	1:A:421:ARG:CZ	2.48	0.43
1:B:336:TRP:CG	1:B:341:VAL:HG22	2.53	0.43
1:A:336:TRP:CD2	1:A:341:VAL:HG22	2.53	0.43
1:B:51:THR:HB	1:B:157:LEU:HD23	2.00	0.43
1:C:45:GLY:N	3:C:2028:HOH:O	2.46	0.43
1:B:480:ALA:HB3	1:B:481:PRO:HD3	2.01	0.43
1:C:51:THR:HB	1:C:157:LEU:HD23	1.99	0.43
1:B:67:LEU:HB3	1:B:68:PRO:HD3	2.01	0.42
1:C:125:LEU:O	1:C:129:LEU:HD13	2.19	0.42
1:D:405:LYS:HE3	1:D:423:MET:HE2	2.00	0.42
1:B:61:LEU:HD13	1:B:178:TYR:HE2	1.84	0.42
1:D:249:PRO:HG3	1:D:421:ARG:CZ	2.49	0.42
1:B:267:ARG:NH1	3:B:2201:HOH:O	2.48	0.42
1:A:102:GLY:O	1:B:370:LYS:NZ	2.46	0.42
1:A:51:THR:HB	1:A:157:LEU:HD23	2.02	0.42
1:B:336:TRP:CD2	1:B:341:VAL:HG22	2.55	0.42
1:D:129:LEU:N	1:D:129:LEU:HD12	2.34	0.41
1:C:67:LEU:HB3	1:C:68:PRO:HD3	2.01	0.41
1:C:88:GLU:OE2	3:C:2063:HOH:O	2.22	0.41
1:C:44:VAL:HG22	1:C:45:GLY:H	1.86	0.41
1:A:12:THR:HG21	1:A:358:ILE:HD13	2.02	0.40
1:C:280:MET:HA	1:C:292:GLY:O	2.22	0.40
1:C:323:ASN:C	1:C:323:ASN:OD1	2.59	0.40
1:D:15:TRP:CE3	1:D:180:SER:HA	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	488/511 (96%)	474 (97%)	14 (3%)	0	100	100
1	B	493/511 (96%)	480 (97%)	13 (3%)	0	100	100
1	C	492/511 (96%)	475 (96%)	17 (4%)	0	100	100
1	D	490/511 (96%)	475 (97%)	15 (3%)	0	100	100
All	All	1963/2044 (96%)	1904 (97%)	59 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	411/424 (97%)	407 (99%)	4 (1%)	76	67
1	B	414/424 (98%)	409 (99%)	5 (1%)	71	59
1	C	414/424 (98%)	409 (99%)	5 (1%)	71	59
1	D	412/424 (97%)	409 (99%)	3 (1%)	84	77
All	All	1651/1696 (97%)	1634 (99%)	17 (1%)	69	67

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	44	VAL
1	A	170	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	293	TYR
1	A	325	ILE
1	B	44	VAL
1	B	53	ARG
1	B	170	PHE
1	B	293	TYR
1	B	471	ARG
1	C	44	VAL
1	C	53	ARG
1	C	170	PHE
1	C	293	TYR
1	C	471	ARG
1	D	170	PHE
1	D	293	TYR
1	D	325	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	197	HIS
1	C	197	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	TRP	B	1512	-	12,16,16	0.74	0	12,22,22	0.97	0
2	TRP	A	1512	-	12,16,16	1.05	1 (8%)	12,22,22	0.82	0
2	TRP	D	1512	-	12,16,16	1.11	1 (8%)	12,22,22	0.84	0
2	TRP	C	1512	-	12,16,16	0.96	1 (8%)	12,22,22	0.83	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TRP	B	1512	-	-	1/3/8/8	0/2/2/2
2	TRP	A	1512	-	-	1/3/8/8	0/2/2/2
2	TRP	D	1512	-	-	1/3/8/8	0/2/2/2
2	TRP	C	1512	-	-	1/3/8/8	0/2/2/2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	1512	TRP	CZ2-CE2	-2.72	1.37	1.41
2	A	1512	TRP	CZ2-CE2	-2.51	1.37	1.41
2	C	1512	TRP	CA-N	2.17	1.51	1.47

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1512	TRP	CA-CB-CG-CD1
2	A	1512	TRP	CA-CB-CG-CD1
2	D	1512	TRP	CA-CB-CG-CD1
2	C	1512	TRP	CA-CB-CG-CD1

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	494/511 (96%)	0.08	1 (0%) 95 95	9, 14, 23, 29	0
1	B	499/511 (97%)	0.19	11 (2%) 62 66	6, 14, 24, 31	0
1	C	499/511 (97%)	0.14	9 (1%) 68 72	7, 14, 24, 30	0
1	D	498/511 (97%)	0.17	8 (1%) 72 76	8, 14, 23, 30	0
All	All	1990/2044 (97%)	0.15	29 (1%) 73 77	6, 14, 23, 31	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	3	ARG	3.3
1	B	152	LEU	3.3
1	B	316	GLY	3.0
1	B	64	ARG	2.9
1	A	3	ARG	2.8
1	B	425	ASP	2.8
1	D	376	ARG	2.6
1	B	488	ARG	2.6
1	D	318	ASP	2.6
1	B	44	VAL	2.5
1	D	229	ARG	2.5
1	B	325	ILE	2.5
1	C	469	ARG	2.5
1	B	490	ARG	2.5
1	D	261	GLU	2.4
1	C	325	ILE	2.4
1	C	468	GLU	2.4
1	C	323	ASN	2.3
1	C	316	GLY	2.3
1	C	64	ARG	2.3
1	D	491	ARG	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	119	PHE	2.3
1	C	491	ARG	2.2
1	B	3	ARG	2.2
1	D	323	ASN	2.2
1	C	39	VAL	2.2
1	B	151	SER	2.1
1	D	425	ASP	2.1
1	B	491	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	TRP	C	1512	15/15	0.95	0.08	9,10,11,11	0
2	TRP	A	1512	15/15	0.96	0.08	9,9,11,11	0
2	TRP	B	1512	15/15	0.96	0.10	9,10,11,12	0
2	TRP	D	1512	15/15	0.97	0.09	9,10,11,12	0

6.5 Other polymers [i](#)

There are no such residues in this entry.