



Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 07:23 am BST

PDB ID : 4WES
Title : Nitrogenase molybdenum-iron protein from *Clostridium pasteurianum* at 1.08 Å resolution
Authors : Zhang, L.M.; Morrison, C.N.; Kaiser, J.T.; Rees, D.C.
Deposited on : 2014-09-10
Resolution : 1.08 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1386 (1.12-1.04)
Clashscore	141614	1021 (1.10-1.06)
Ramachandran outliers	138981	1381 (1.12-1.04)
Sidechain outliers	138945	1379 (1.12-1.04)
RSRZ outliers	127900	1359 (1.12-1.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	534	<div> <div> <div></div> <div>90%</div> <div>7%</div> <div></div> </div> <div> <div></div> <div></div> </div> </div>
1	C	534	<div> <div> <div></div> <div>89%</div> <div>7%</div> <div></div> </div> <div> <div></div> <div></div> </div> </div>
2	B	458	<div> <div> <div></div> <div>94%</div> <div>6%</div> <div></div> </div> <div> <div></div> <div></div> </div> </div>
2	D	458	<div> <div> <div></div> <div>96%</div> <div></div> <div></div> </div> <div> <div></div> <div></div> </div> </div>

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 18065 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Nitrogenase molybdenum-iron protein alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	519	Total	C	N	O	S	0	16	0
			4125	2637	680	778	30			
1	C	518	Total	C	N	O	S	0	13	0
			4102	2614	680	778	30			

- Molecule 2 is a protein called Nitrogenase molybdenum-iron protein beta chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	458	Total	C	N	O	S	0	21	0
			3626	2325	588	688	25			
2	D	458	Total	C	N	O	S	0	13	0
			3588	2288	586	690	24			

- Molecule 3 is 3-HYDROXY-3-CARBOXY-ADIPIC ACID (three-letter code: HCA) (formula: $C_7H_{10}O_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			14	7	7		
3	C	1	Total	C	O	0	0
			14	7	7		

- Molecule 4 is iron-sulfur-molybdenum cluster with interstitial carbon (three-letter code: ICS) (formula: CFe_7MoS_9).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	Fe	Mo	S	0	0
			18	1	7	1	9		
4	C	1	Total	C	Fe	Mo	S	0	0
			18	1	7	1	9		

- Molecule 5 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: $\text{C}_6\text{H}_{14}\text{O}_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			8	6	2		

- Molecule 6 is FE(8)-S(7) CLUSTER (three-letter code: CLF) (formula: Fe_8S_7).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	Fe	S	0	1
			17	10	7		
6	C	1	Total	Fe	S	0	1
			17	10	7		

- Molecule 7 is FE (II) ION (three-letter code: FE2) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	2	Total 2	Fe 2	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	700	Total 716	O 716	0	15
8	B	533	Total 538	O 538	0	4
8	C	641	Total 644	O 644	0	3
8	D	610	Total 618	O 618	0	7


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Nitrogenase molybdenum-iron protein alpha chain




- Molecule 1: Nitrogenase molybdenum-iron protein alpha chain



- Molecule 2: Nitrogenase molybdenum-iron protein beta chain



- Molecule 2: Nitrogenase molybdenum-iron protein beta chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	72.70Å 170.58Å 87.54Å 90.00° 91.63° 90.00°	Depositor
Resolution (Å)	38.93 – 1.08 35.54 – 1.08	Depositor EDS
% Data completeness (in resolution range)	93.0 (38.93-1.08) 93.0 (35.54-1.08)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.69 (at 1.08Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.110 , 0.133 0.112 , 0.135	Depositor DCC
R_{free} test set	16793 reflections (1.99%)	wwPDB-VP
Wilson B-factor (Å ²)	9.9	Xtriage
Anisotropy	0.302	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 61.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.025 for h,-k,-l	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	18065	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ICS, MPD, HCA, FE2, CLF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.59	1/4262 (0.0%)	0.74	3/5756 (0.1%)
1	C	0.54	0/4227	0.71	1/5710 (0.0%)
2	B	0.64	3/3761 (0.1%)	0.73	1/5082 (0.0%)
2	D	0.67	4/3695 (0.1%)	0.75	1/4991 (0.0%)
All	All	0.61	8/15945 (0.1%)	0.73	6/21539 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	323	GLU	CD-OE2	11.07	1.37	1.25
1	A	521	LYS	C-O	10.75	1.43	1.23
2	B	323	GLU	CD-OE2	10.53	1.37	1.25
2	B	130	GLU	CD-OE2	8.42	1.34	1.25
2	B	130	GLU	CD-OE1	7.81	1.34	1.25
2	D	8	GLU	CD-OE1	6.86	1.33	1.25
2	D	124	ASP	CG-OD2	5.83	1.38	1.25
2	D	227	GLU	CD-OE1	5.47	1.31	1.25

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	406	ARG	NE-CZ-NH2	7.12	123.86	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	406	ARG	NE-CZ-NH1	-7.03	116.79	120.30
1	C	406	ARG	NE-CZ-NH2	-5.70	117.45	120.30
2	D	408	ARG	NE-CZ-NH1	5.28	122.94	120.30
2	B	416	ARG	NE-CZ-NH1	5.05	122.83	120.30
1	A	490	TYR	CB-CG-CD1	5.01	124.01	121.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	486	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4125	0	4138	33	0
1	C	4102	0	4083	33	0
2	B	3626	0	3707	20	0
2	D	3588	0	3621	14	0
3	A	14	0	6	1	0
3	C	14	0	6	1	0
4	A	18	0	0	1	0
4	C	18	0	0	1	0
5	A	8	0	14	2	0
6	A	17	0	0	0	0
6	C	17	0	0	0	0
7	B	2	0	0	0	0
8	A	716	0	0	14	0
8	B	538	0	0	5	0
8	C	644	0	0	11	0
8	D	618	0	0	8	0
All	All	18065	0	15575	98	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (98) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:386:ILE:HG23	8:A:1201[B]:HOH:O	1.42	1.20
1:C:20[B]:THR:HG22	8:C:1171:HOH:O	1.50	1.10
1:A:31:GLU:HG2	8:A:1395:HOH:O	1.51	1.09
1:A:20[B]:THR:HG22	8:A:1186:HOH:O	1.65	0.97
1:C:415:GLU:O	1:C:419:LYS:HG2	1.72	0.90
1:A:90[B]:LYS:HE3	8:D:1150:HOH:O	1.73	0.86
1:A:90[B]:LYS:CE	8:D:1150:HOH:O	2.26	0.84
1:A:332[A]:GLU:OE1	8:A:1368:HOH:O	1.97	0.81
2:B:186:GLU:O	2:B:190[A]:LEU:HD23	1.83	0.78
1:A:425:SER:OG	8:A:1235:HOH:O	1.97	0.77
1:C:212:VAL:HG11	1:C:225[B]:MET:HE1	1.64	0.77
1:C:425:SER:OG	8:C:1336:HOH:O	2.03	0.76
1:A:387:ASP:HB3	8:A:1388[A]:HOH:O	1.86	0.75
2:B:145:SER:OG	2:B:147[A]:VAL:HG12	1.88	0.74
2:D:349:LYS:NZ	2:D:353:GLU:HG3	2.02	0.74
1:C:454:GLU:OE1	8:C:1042:HOH:O	2.07	0.72
2:B:360:GLU:OE1	8:B:1130:HOH:O	2.08	0.70
1:A:35:PRO:HD2	1:A:431[B]:MET:HE1	1.73	0.68
2:D:75:ALA:HB2	8:D:922:HOH:O	1.92	0.68
1:A:35:PRO:CD	1:A:431[B]:MET:HE1	2.25	0.66
1:A:35:PRO:HD2	1:A:431[B]:MET:CE	2.26	0.66
1:A:428:GLY:O	1:A:432:LYS:HD3	1.95	0.65
2:B:186:GLU:OE2	2:B:190[A]:LEU:HD21	1.97	0.65
1:A:397[B]:THR:HG23	8:A:1135:HOH:O	1.96	0.64
2:D:349:LYS:HZ2	2:D:353:GLU:HG3	1.59	0.64
1:C:415:GLU:O	1:C:419:LYS:CG	2.46	0.64
1:A:506:GLY:HA2	2:D:307:GLN:HE22	1.63	0.63
1:A:379:GLU:OE2	1:A:432:LYS:NZ	2.31	0.63
2:B:278:MET:HE1	1:C:519:TRP:HB3	1.81	0.63
2:D:317:LEU:HD21	2:D:331[B]:ILE:CD1	2.30	0.62
1:C:44:VAL:O	1:C:47[B]:ILE:HG13	2.00	0.62
2:D:440:VAL:HG13	8:D:1202:HOH:O	2.00	0.60
2:D:8:GLU:HG3	8:D:1035:HOH:O	2.02	0.60
1:C:20[B]:THR:CG2	8:C:1171:HOH:O	2.24	0.60
8:A:1312:HOH:O	2:B:73:GLU:CD	2.41	0.59
1:C:176:GLY:O	8:C:1039[B]:HOH:O	2.17	0.59
2:B:75:ALA:HB2	8:B:787:HOH:O	2.04	0.57
2:B:278:MET:HA	2:B:278:MET:CE	2.35	0.57
2:B:278:MET:HA	2:B:278:MET:HE2	1.87	0.57
1:C:3:GLU:CG	1:C:3:GLU:O	2.52	0.57
1:C:164:GLU:OE2	8:C:1292:HOH:O	2.18	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:317:LEU:HD21	2:D:331[B]:ILE:HD12	1.90	0.53
1:C:231:LYS:HE2	1:C:318:GLU:OE1	2.08	0.53
5:A:603:MPD:O4	5:A:603:MPD:C1	2.56	0.53
1:C:3:GLU:N	1:C:7:ASP:OD2	2.41	0.53
1:C:411:GLU:OE2	1:C:415:GLU:HG2	2.08	0.53
5:A:603:MPD:H12	5:A:603:MPD:O4	2.09	0.52
1:C:293[A]:VAL:HG23	8:C:988:HOH:O	2.10	0.52
2:D:349:LYS:NZ	2:D:353:GLU:CG	2.73	0.50
2:B:440:VAL:HG13	8:B:1108:HOH:O	2.10	0.50
1:C:212:VAL:HG11	1:C:225[B]:MET:CE	2.37	0.50
1:A:44:VAL:O	1:A:47[C]:ILE:HG13	2.12	0.50
1:C:465:ILE:HG21	2:D:54:THR:HG23	1.93	0.49
1:A:386:ILE:CG2	8:A:1201[B]:HOH:O	2.19	0.49
2:B:100:ILE:HG23	8:B:921:HOH:O	2.13	0.49
1:A:465:ILE:HG21	2:B:54:THR:HG23	1.95	0.49
1:C:287:ILE:HD13	1:C:351:TYR:CE1	2.47	0.49
1:A:20[B]:THR:CG2	8:A:1186:HOH:O	2.41	0.48
2:D:106:CYS:HB3	2:D:141[A]:SER:OG	2.13	0.48
1:A:293[A]:VAL:HG23	8:A:952:HOH:O	2.13	0.47
2:D:317:LEU:HD21	2:D:331[B]:ILE:HD11	1.95	0.47
1:C:482:HIS:HB3	3:C:601:HCA:O6	2.15	0.47
2:D:262[A]:LYS:HB3	2:D:262[A]:LYS:HE3	1.45	0.47
1:A:482:HIS:HB3	3:A:601:HCA:O5	2.16	0.46
1:C:3:GLU:O	1:C:3:GLU:HG2	2.14	0.46
2:B:155:GLN:O	2:B:158[A]:VAL:HG22	2.16	0.46
1:A:365:ALA:O	1:A:440:LEU:HA	2.16	0.46
2:B:145:SER:HG	2:B:147[A]:VAL:HG12	1.82	0.45
2:B:251:LYS:HG2	2:B:261:PHE:CZ	2.52	0.45
1:A:90[B]:LYS:NZ	8:A:1380:HOH:O	2.41	0.44
2:B:263[A]:THR:HG22	8:B:865:HOH:O	2.17	0.44
2:B:106:CYS:HB3	2:B:141[A]:SER:OG	2.17	0.44
1:A:281:TRP:CE2	1:A:425:SER:HA	2.52	0.44
1:C:315:VAL:HA	1:C:318:GLU:HG2	1.98	0.44
1:C:365:ALA:O	1:C:440:LEU:HA	2.18	0.44
2:D:100:ILE:HG23	8:D:984:HOH:O	2.17	0.44
1:C:344:GLY:HA3	4:C:602:ICS:S1B	2.58	0.44
1:A:90[B]:LYS:CD	8:D:1150:HOH:O	2.63	0.43
1:A:414:VAL:CG1	1:A:418:LYS:HE3	2.48	0.43
1:A:176:GLY:O	8:A:1044[B]:HOH:O	2.21	0.43
1:A:90[B]:LYS:HD2	8:D:1150:HOH:O	2.18	0.43
1:C:432:LYS:HD2	8:C:945:HOH:O	2.18	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:228:VAL:HG21	8:C:981:HOH:O	2.19	0.42
1:A:47[B]:ILE:HG12	1:A:445:ASN:ND2	2.34	0.42
1:A:42:ARG:NH2	8:A:1297:HOH:O	2.52	0.42
1:C:240:LEU:HA	1:C:241:THR:HA	1.87	0.42
1:C:203:LYS:HE3	1:C:247:GLU:OE1	2.20	0.42
1:A:370:ALA:HB1	1:A:374:ASP:HB2	2.01	0.41
1:C:432:LYS:CD	8:C:945:HOH:O	2.67	0.41
1:C:293[A]:VAL:HG21	8:C:1036:HOH:O	2.20	0.41
1:A:310:LYS:HE3	1:A:314:GLU:OE2	2.20	0.41
1:A:344:GLY:HA3	4:A:602:ICS:S1B	2.61	0.41
1:C:370:ALA:HB1	1:C:374:ASP:HB2	2.01	0.41
2:B:341:VAL:HG11	2:B:354[B]:ILE:HD13	2.02	0.41
2:B:155:GLN:HA	2:B:158[A]:VAL:HG22	2.02	0.41
1:C:225[B]:MET:HB2	1:C:225[B]:MET:HE2	1.87	0.40
1:C:281:TRP:CE2	1:C:425:SER:HA	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	533/534 (100%)	516 (97%)	15 (3%)	2 (0%)	34	10
1	C	529/534 (99%)	513 (97%)	14 (3%)	2 (0%)	34	10
2	B	477/458 (104%)	469 (98%)	8 (2%)	0	100	100
2	D	469/458 (102%)	459 (98%)	10 (2%)	0	100	100
All	All	2008/1984 (101%)	1957 (98%)	47 (2%)	4 (0%)	47	17

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	463	ALA
1	C	463	ALA
1	A	48	ILE
1	C	48	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	452/447 (101%)	448 (99%)	4 (1%)	78	47
1	C	447/447 (100%)	444 (99%)	3 (1%)	84	55
2	B	405/384 (106%)	402 (99%)	3 (1%)	84	55
2	D	397/384 (103%)	397 (100%)	0	100	100
All	All	1701/1662 (102%)	1691 (99%)	10 (1%)	88	60

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	343	VAL
1	A	406	ARG
1	A	432	LYS
1	A	461	PHE
2	B	307	GLN
2	B	422[A]	ASN
2	B	422[B]	ASN
1	C	343	VAL
1	C	419	LYS
1	C	461	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	121	ASN
2	D	307	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 11 ligands modelled in this entry, 2 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	HCA	C	601	-	4,13,13	0.97	0	4,18,18	3.02	2 (50%)
6	CLF	C	603[B]	1,2	0,24,24	0.00	-	-		
4	ICS	A	602	1	18,30,30	2.34	9 (50%)	-		
5	MPD	A	603	-	7,7,7	0.37	0	9,10,10	0.86	0
4	ICS	C	602	1	18,30,30	2.23	7 (38%)	-		
6	CLF	A	604[A]	1,2	0,24,24	0.00	-	-		
6	CLF	C	603[A]	1,2	0,24,24	0.00	-	-		
6	CLF	A	604[B]	1,2	0,24,24	0.00	-	-		
3	HCA	A	601	-	4,13,13	0.79	0	4,18,18	2.83	2 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HCA	C	601	-	-	1/7/17/17	-
6	CLF	C	603[B]	1,2	-	-	0/12/10/10
5	MPD	A	603	-	-	3/5/5/5	-
6	CLF	A	604[A]	1,2	-	-	0/12/10/10
6	CLF	C	603[A]	1,2	-	-	0/12/10/10
6	CLF	A	604[B]	1,2	-	-	0/12/10/10
3	HCA	A	601	-	-	1/7/17/17	-

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	602	ICS	S3B-FE6	-3.95	2.22	2.32
4	A	602	ICS	S4B-FE7	-3.93	2.22	2.32
4	C	602	ICS	S1B-FE6	-3.78	2.23	2.32
4	C	602	ICS	S3B-FE6	-3.73	2.23	2.32
4	A	602	ICS	S1B-FE6	-3.71	2.23	2.32
4	C	602	ICS	S4B-FE7	-3.70	2.23	2.32
4	A	602	ICS	S4B-FE5	-3.25	2.24	2.32
4	A	602	ICS	S3B-FE7	-3.04	2.24	2.32
4	C	602	ICS	S4B-FE5	-3.02	2.24	2.32
4	C	602	ICS	S3B-FE7	-2.75	2.25	2.32
4	C	602	ICS	S2B-FE6	-2.58	2.18	2.24
4	A	602	ICS	S2B-FE6	-2.56	2.18	2.24
4	A	602	ICS	S1B-FE5	-2.53	2.26	2.32
4	C	602	ICS	S1B-FE5	-2.42	2.26	2.32
4	A	602	ICS	S1A-FE2	-2.35	2.26	2.32
4	A	602	ICS	S4A-FE3	-2.10	2.27	2.32

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	601	HCA	C4-C5-C6	5.61	119.89	111.39
3	A	601	HCA	C4-C5-C6	5.20	119.26	111.39
3	A	601	HCA	C4-C3-C7	-2.13	107.77	111.52
3	C	601	HCA	C4-C3-C7	-2.06	107.88	111.52

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	603	MPD	C1-C2-C3-C4
5	A	603	MPD	O2-C2-C3-C4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
5	A	603	MPD	CM-C2-C3-C4
3	A	601	HCA	C1-C2-C3-C4
3	C	601	HCA	C1-C2-C3-C4


There are no ring outliers.

5 monomers are involved in 6 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	601	HCA	1	0
4	A	602	ICS	1	0
5	A	603	MPD	2	0
4	C	602	ICS	1	0
3	A	601	HCA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand ICS A 602




Bond lengths




Bond angles



Torsions



Rings



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	519/534 (97%)	-0.34	7 (1%) 77 71	7, 12, 37, 162	0
1	C	518/534 (97%)	-0.31	11 (2%) 63 56	8, 13, 38, 108	0
2	B	458/458 (100%)	-0.42	2 (0%) 92 89	7, 12, 28, 76	0
2	D	458/458 (100%)	-0.50	1 (0%) 95 91	7, 10, 24, 72	0
All	All	1953/1984 (98%)	-0.39	21 (1%) 80 75	7, 12, 32, 162	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	419	LYS	4.3
1	A	386	ILE	3.7
1	C	386	ILE	3.5
1	A	3	GLU	3.5
1	C	32	THR	3.4
2	B	77	VAL	3.4
2	B	78	PHE	3.1
1	C	412	ASP	2.9
1	C	29	THR	2.9
1	A	419	LYS	2.9
1	C	410	PRO	2.9
1	A	521	LYS	2.8
2	D	78	PHE	2.6
1	C	409	ILE	2.6
1	A	32	THR	2.5
1	A	31	GLU	2.5
1	C	3	GLU	2.4
1	C	415	GLU	2.4
1	A	385	LYS	2.4
1	C	411	GLU	2.3
1	C	30	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	MPD	A	603	8/8	0.91	0.09	26,37,52,63	0
3	HCA	C	601	14/14	0.99	0.06	8,8,10,11	0
3	HCA	A	601	14/14	0.99	0.07	7,8,10,10	0
6	CLF	C	603[B]	15/15	1.00	0.04	7,8,8,9	2
4	ICS	C	602	18/18	1.00	0.04	8,9,10,10	0
7	FE2	B	501	1/1	1.00	0.07	7,7,7,7	1
7	FE2	B	502	1/1	1.00	0.05	7,7,7,7	1
6	CLF	A	604[A]	15/15	1.00	0.05	8,9,10,11	2
6	CLF	C	603[A]	15/15	1.00	0.04	7,8,8,11	2
6	CLF	A	604[B]	15/15	1.00	0.05	8,9,9,9	2
4	ICS	A	602	18/18	1.00	0.05	7,8,8,8	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around ICS C 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ICS A 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.