



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 22, 2020 – 12:14 PM BST

PDB ID : 5WE8
Title : Crystal structure of WNK1 in complex with N-{(3R)-1-[(4-chlorophenyl)methyl]pyrrolidin-3-yl}-2-(3-methoxyphenyl)-N-methylquinoline-4-carboxamide (compound 8)
Authors : Xie, X.; Kohls, D.
Deposited on : 2017-07-07
Resolution : 2.01 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.01 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	279	<div> <div>5%</div> <div> <div></div> <div>87%</div> <div>9%</div> <div></div> </div> </div>
1	B	279	<div> <div>3%</div> <div> <div></div> <div>81%</div> <div>17%</div> <div></div> </div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4748 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein kinase WNK1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	270	Total	C	N	O	S	0	1	0
			2173	1385	370	402	16			
1	B	272	Total	C	N	O	S	0	0	0
			2177	1390	370	401	16			


There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	205	GLY	-	expression tag	UNP Q9H4A3
A	378	ASP	SER	engineered mutation	UNP Q9H4A3
A	396	ALA	GLU	engineered mutation	UNP Q9H4A3
A	397	ALA	GLU	engineered mutation	UNP Q9H4A3
A	398	ALA	LYS	engineered mutation	UNP Q9H4A3
B	205	GLY	-	expression tag	UNP Q9H4A3
B	378	ASP	SER	engineered mutation	UNP Q9H4A3
B	396	ALA	GLU	engineered mutation	UNP Q9H4A3
B	397	ALA	GLU	engineered mutation	UNP Q9H4A3
B	398	ALA	LYS	engineered mutation	UNP Q9H4A3

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mn	0	0
			1	1		
2	A	1	Total	Mn	0	0
			1	1		

- Molecule 3 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula: C₁₀H₁₇N₆O₁₂P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
3	B	1	Total	C	N	O	P	0	0
			31	10	6	12	3		

- Molecule 4 is N-{(3R)-1-[(4-chlorophenyl)methyl]pyrrolidin-3-yl}-2-(3-methoxyphenyl)-N-methylquinoline-4-carboxamide (three-letter code: A7Y) (formula: C₂₉H₂₈ClN₃O₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	Cl	N	O	0	0
			35	29	1	3	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	Cl	N	O	0	0
			35	29	1	3	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	126	Total	O	0	0
			126	126		
5	B	138	Total	O	0	0
			138	138		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Serine/threonine-protein kinase WNK1



- Molecule 1: Serine/threonine-protein kinase WNK1



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	83.10Å 105.23Å 70.11Å 90.00° 117.28° 90.00°	Depositor
Resolution (Å)	41.08 – 2.01 41.08 – 2.01	Depositor EDS
% Data completeness (in resolution range)	96.8 (41.08-2.01) 97.0 (41.08-2.01)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.50 (at 2.00Å)	Xtriage
Refinement program	PHENIX 1.11.1_2575	Depositor
R, R_{free}	0.203 , 0.247 0.203 , 0.246	Depositor DCC
R_{free} test set	1734 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	31.0	Xtriage
Anisotropy	0.522	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 50.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.011 for -1/2*h+1/2*k-l,1/2*h-1/2*k-l,-1/2 *h-1/2*k 0.000 for -1/2*h-1/2*k-l,-1/2*h-1/2*k+l,-1/ 2*h+1/2*k	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4748	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 41.19 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.4494e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MN, ANP, A7Y

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.31	0/2217	0.47	0/2982
1	B	0.26	0/2218	0.45	0/2983
All	All	0.28	0/4435	0.46	0/5965

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2173	0	2197	28	0
1	B	2177	0	2207	30	1
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	31	0	13	0	0
3	B	31	0	13	2	0
4	A	35	0	0	0	0
4	B	35	0	0	1	0
5	A	126	0	0	3	0
5	B	138	0	0	1	0
All	All	4748	0	4430	54	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:276:GLN:CG	1:A:282:ARG:HH11	1.62	1.13
1:A:276:GLN:HG2	1:A:282:ARG:HH11	1.37	0.88
1:A:276:GLN:CG	1:A:282:ARG:NH1	2.44	0.79
1:A:276:GLN:HG3	1:A:282:ARG:HH11	1.47	0.79
1:A:464:LYS:NZ	5:A:601:HOH:O	2.17	0.78
1:B:233:LYS:HB3	1:B:252:LEU:HD13	1.66	0.77
1:B:252:LEU:HB2	1:B:297:ILE:HG23	1.69	0.72
1:A:229[A]:ARG:NH1	5:A:602:HOH:O	2.24	0.71
1:A:276:GLN:HG3	1:A:282:ARG:NH1	2.07	0.66
3:B:502:ANP:H8	3:B:502:ANP:H5'1	1.78	0.65
1:A:283:PHE:HE1	1:A:299:LEU:HD22	1.64	0.63
1:B:321:ILE:HD11	1:B:478:PHE:HE1	1.63	0.63
1:B:351:LYS:NZ	3:B:502:ANP:O2G	2.28	0.60
1:A:432:TYR:O	1:A:435:VAL:HG12	2.01	0.59
1:A:276:GLN:HG2	1:A:282:ARG:NH1	2.15	0.58
1:A:276:GLN:NE2	5:A:603:HOH:O	2.37	0.58
1:A:276:GLN:CB	1:A:282:ARG:HH11	2.19	0.56
1:A:392:PRO:HG3	1:B:435:VAL:HG11	1.88	0.56
1:A:477:ALA:O	1:A:481:GLU:HG2	2.07	0.55
1:B:322:LYS:HD3	5:B:684:HOH:O	2.06	0.54
1:B:321:ILE:HD11	1:B:478:PHE:CE1	2.45	0.52
1:B:343:PRO:HG2	1:B:374:LEU:HD23	1.90	0.52
1:B:390:MET:HB2	1:B:395:TYR:CE2	2.46	0.51
1:A:325:ARG:HD3	1:A:478:PHE:O	2.11	0.50
1:A:282:ARG:HB3	1:A:302:GLU:HB2	1.93	0.50
1:A:226:GLU:OE2	1:A:229[B]:ARG:NH1	2.44	0.50
1:A:255:ARG:NH1	1:B:378:ASP:OD2	2.45	0.50
1:A:390:MET:HG3	1:A:394:MET:HB3	1.94	0.49
1:B:257:LEU:HB3	1:B:262:ARG:HG2	1.95	0.49
1:A:283:PHE:CE1	1:A:299:LEU:HD22	2.46	0.48
1:A:260:SER:HB3	1:B:263:GLN:HG3	1.96	0.47
1:A:329:ARG:NH2	1:A:481:GLU:O	2.48	0.47
1:A:257:LEU:CD1	1:A:261:GLU:HB3	2.45	0.46
1:B:214:MET:HG2	1:B:215:SER:O	2.16	0.46
1:A:392:PRO:HG3	1:B:435:VAL:CG1	2.45	0.46
1:B:449:ILE:HG22	1:B:452:VAL:H	1.81	0.45
1:B:282:ARG:HB3	1:B:302:GLU:HB2	2.00	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:214:MET:SD	1:B:218:GLY:HA2	2.58	0.43
1:B:222:LYS:HB2	1:B:249:TRP:CD2	2.53	0.43
1:B:402:SER:OG	1:B:467:ARG:HD2	2.18	0.43
1:B:372:ALA:HB3	4:B:503:A7Y:C5	2.48	0.43
1:A:464:LYS:HG2	1:A:464:LYS:H	1.62	0.42
1:A:393:GLU:HG3	1:B:399:TYR:HB3	2.02	0.42
1:B:279:ASN:ND2	1:B:330:GLN:HB3	2.35	0.42
1:B:304:MET:HE2	1:B:365:LYS:HB2	2.01	0.42
1:B:398:ALA:O	1:B:464:LYS:HE2	2.19	0.42
1:B:309:LEU:O	1:B:313:LEU:HG	2.20	0.41
1:B:257:LEU:O	1:B:262:ARG:NH1	2.53	0.41
1:A:246:GLU:N	1:A:246:GLU:OE1	2.46	0.41
1:A:257:LEU:HD13	1:A:261:GLU:HB3	2.01	0.41
1:B:311:THR:O	1:B:314:LYS:HG2	2.20	0.41
1:B:258:THR:O	1:B:262:ARG:HG3	2.21	0.41
1:B:222:LYS:HB2	1:B:249:TRP:CE3	2.56	0.41
1:B:223:PHE:HE1	1:B:239:LEU:HB2	1.86	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:305:THR:OG1	1:B:305:THR:OG1[2_756]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	267/279 (96%)	258 (97%)	9 (3%)	0	100	100
1	B	268/279 (96%)	263 (98%)	5 (2%)	0	100	100
All	All	535/558 (96%)	521 (97%)	14 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	238/244 (98%)	235 (99%)	3 (1%)	69	74
1	B	238/244 (98%)	234 (98%)	4 (2%)	60	65
All	All	476/488 (98%)	469 (98%)	7 (2%)	65	69

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	266	LYS
1	A	282	ARG
1	A	353	ASP
1	B	207	LEU
1	B	256	LYS
1	B	403	VAL
1	B	418	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	276	GLN
1	B	480	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	A7Y	A	503	-	39,39,39	0.93	0	51,55,55	1.51	8 (15%)
3	ANP	B	502	2	29,33,33	1.84	5 (17%)	31,52,52	1.79	7 (22%)
3	ANP	A	502	2	29,33,33	1.89	7 (24%)	31,52,52	1.98	8 (25%)
4	A7Y	B	503	-	39,39,39	0.92	0	51,55,55	1.47	10 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	A7Y	A	503	-	-	0/22/31/31	0/5/5/5
3	ANP	B	502	2	-	8/14/38/38	0/3/3/3
3	ANP	A	502	2	-	4/14/38/38	0/3/3/3
4	A7Y	B	503	-	-	2/22/31/31	0/5/5/5

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	502	ANP	PB-N3B	4.54	1.75	1.63
3	B	502	ANP	PG-N3B	4.54	1.75	1.63
3	A	502	ANP	PB-N3B	4.52	1.75	1.63
3	A	502	ANP	PG-N3B	4.50	1.75	1.63
3	B	502	ANP	PB-O1B	3.43	1.51	1.46
3	A	502	ANP	PG-O1G	3.41	1.51	1.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	502	ANP	PG-O1G	3.31	1.51	1.46
3	A	502	ANP	PB-O1B	3.22	1.51	1.46
3	A	502	ANP	PB-O3A	2.81	1.62	1.59
3	B	502	ANP	C5-C4	2.50	1.47	1.40
3	A	502	ANP	C5-C4	2.49	1.47	1.40
3	A	502	ANP	O4'-C1'	2.05	1.43	1.41

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	ANP	O1G-PG-N3B	-6.29	102.51	111.77
3	B	502	ANP	O1G-PG-N3B	-4.93	104.52	111.77
4	B	503	A7Y	C38-N37-C36	4.57	121.46	118.06
4	A	503	A7Y	C38-N37-C36	4.30	121.26	118.06
3	A	502	ANP	O2B-PB-O1B	4.20	118.73	109.92
3	A	502	ANP	O1B-PB-N3B	-3.75	106.24	111.77
3	B	502	ANP	O2B-PB-O1B	3.74	117.75	109.92
4	A	503	A7Y	C24-N15-C16	3.40	107.34	104.02
4	B	503	A7Y	C28-N27-C19	3.31	121.70	117.89
3	A	502	ANP	N3-C2-N1	-3.21	123.66	128.68
4	A	503	A7Y	C28-N27-C19	3.20	121.57	117.89
3	B	502	ANP	N3-C2-N1	-3.13	123.78	128.68
4	A	503	A7Y	C12-N15-C24	3.09	117.68	113.18
3	B	502	ANP	C3'-C2'-C1'	3.03	105.53	100.98
3	B	502	ANP	C4-C5-N7	-2.88	106.40	109.40
3	B	502	ANP	PA-O3A-PB	-2.86	122.54	132.62
4	A	503	A7Y	C34-C32-N27	2.86	122.47	118.80
4	B	503	A7Y	C12-N15-C24	2.83	117.31	113.18
4	B	503	A7Y	C34-C32-N27	2.80	122.39	118.80
4	A	503	A7Y	C35-C34-C32	2.73	124.07	120.41
4	B	503	A7Y	C12-N15-C16	2.73	117.90	113.27
4	B	503	A7Y	C35-C34-C32	2.68	124.00	120.41
3	A	502	ANP	C3'-C2'-C1'	2.67	105.00	100.98
4	B	503	A7Y	O33-C32-C34	-2.63	115.12	120.62
4	A	503	A7Y	O33-C32-C34	-2.55	115.29	120.62
4	B	503	A7Y	C52-O51-C44	2.55	123.05	117.51
3	A	502	ANP	C4-C5-N7	-2.51	106.78	109.40
3	A	502	ANP	O3G-PG-O2G	2.43	114.12	107.64
3	B	502	ANP	O3G-PG-O2G	2.33	113.84	107.64
4	B	503	A7Y	C39-C38-N37	-2.30	120.14	122.23
4	A	503	A7Y	C42-C41-C38	-2.22	117.38	120.59
3	A	502	ANP	PA-O3A-PB	-2.17	124.96	132.62

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	503	A7Y	C41-C38-N37	2.08	119.97	117.09

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	502	ANP	PB-N3B-PG-O1G
3	B	502	ANP	C5'-O5'-PA-O1A
3	B	502	ANP	C5'-O5'-PA-O2A
3	A	502	ANP	PB-N3B-PG-O1G
3	A	502	ANP	PA-O3A-PB-O1B
3	A	502	ANP	PA-O3A-PB-O2B
4	B	503	A7Y	C7-C12-N15-C16
4	B	503	A7Y	C7-C12-N15-C24
3	B	502	ANP	O4'-C4'-C5'-O5'
3	B	502	ANP	C4'-C5'-O5'-PA
3	B	502	ANP	PB-O3A-PA-O2A
3	A	502	ANP	PB-O3A-PA-O2A
3	B	502	ANP	C5'-O5'-PA-O3A
3	B	502	ANP	C3'-C4'-C5'-O5'


There are no ring outliers.

2 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	502	ANP	2	0
4	B	503	A7Y	1	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand A7Y A 503



Ligand ANP B 502





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	270/279 (96%)	0.09	14 (5%) 27 26	21, 38, 61, 76	0
1	B	272/279 (97%)	0.01	9 (3%) 46 45	20, 39, 59, 72	0
All	All	542/558 (97%)	0.05	23 (4%) 36 35	20, 39, 61, 76	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	209	THR	5.0
1	A	483	THR	4.7
1	A	439	VAL	4.1
1	A	256	LYS	4.1
1	A	437	SER	3.9
1	A	381	LYS	3.8
1	B	254	ASP	3.5
1	A	382	SER	3.1
1	B	291	VAL	2.9
1	B	256	LYS	2.8
1	A	254	ASP	2.6
1	A	431	ILE	2.6
1	A	438	GLY	2.6
1	B	290	THR	2.4
1	B	207	LEU	2.4
1	A	255	ARG	2.3
1	B	359	GLY	2.3
1	A	253	GLN	2.2
1	A	384	ILE	2.1
1	B	255	ARG	2.1
1	A	435	VAL	2.1
1	B	480	GLN	2.1
1	B	396	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	ANP	B	502	31/31	0.85	0.16	43,60,80,92	0
3	ANP	A	502	31/31	0.91	0.11	28,36,59,68	0
4	A7Y	B	503	35/35	0.91	0.15	30,42,50,70	0
4	A7Y	A	503	35/35	0.93	0.13	28,37,47,58	0
2	MN	B	501	1/1	0.97	0.05	58,58,58,58	0
2	MN	A	501	1/1	0.98	0.04	41,41,41,41	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ANP B 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around ANP A 502:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around A7Y B 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around A7Y A 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.