



Full wwPDB X-ray Structure Validation Report ⓘ

May 3, 2021 – 10:11 pm BST

PDB ID : 2WG2
Title : NON-AGED CONJUGATE OF TORPEDO CALIFORNICA ACETYL-
CHOLINESTERASE WITH SOMAN (ALTERNATIVE REFINEMENT)
Authors : Sanson, B.; Nachon, F.; Colletier, J.P.; Froment, M.T.; Toker, L.; Greenblatt,
H.M.; Sussman, J.L.; Ashani, Y.; Masson, P.; Silman, I.; Weik, M.
Deposited on : 2009-04-15
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.18
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.18


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	537	<div> <div>3%</div> <div> <div></div> <div>91%</div> <div>7%</div> <div></div> </div> </div>
2	B	2	<div> <div>100%</div> </div>
3	C	2	<div> <div>50%</div> <div>50%</div> </div>

2 Entry composition [i](#)

There are 9 unique types of molecules in this entry. The entry contains 5206 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ACETYLCHOLINESTERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	532	Total	C	N	O	S	0	27	0
			4395	2828	737	805	25			

- Molecule 2 is an oligosaccharide called alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	2	Total	C	N	O	0	0	0
			24	14	1	9			

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 4 is (1R)-1,2,2-TRIMETHYLPROPYL (S)-METHYLPHOSPHINATE (three-letter code: GD8) (formula: C₇H₁₇O₂P).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	P	0	0
			10	7	2	1		

- Molecule 5 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: $C_4H_{12}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			8	4	1	3		
5	A	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 6 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			13	8	5		
6	A	1	Total	C	O	0	0
			13	8	5		
6	A	1	Total	C	O	0	0
			13	8	5		
6	A	1	Total	C	O	0	0
			13	8	5		
6	A	1	Total	C	O	0	0
			13	8	5		

- Molecule 7 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 8 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	1	Total	Cl	0	0
			1	1		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	660	Total	O	0	0
			660	660		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ACETYLCHOLINESTERASE



- Molecule 2: alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	111.41Å 111.41Å 137.38Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	45.79 – 1.95 43.27 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.3 (45.79-1.95) 98.0 (43.27-1.95)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.02 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.166 , 0.196 0.166 , 0.196	Depositor DCC
R_{free} test set	3487 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	29.2	Xtriage
Anisotropy	0.025	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 55.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.026 for -h,-k,l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5206	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, PG4, NAG, CL, TRS, GD8, FUC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.59	0/4600	0.60	1/6240 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	494	LEU	CA-CB-CG	5.92	128.91	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4395	0	4311	40	0
2	B	24	0	22	0	0
3	C	28	0	25	0	0
4	A	10	0	16	2	0
5	A	16	0	24	0	0
6	A	65	0	90	1	0
7	A	7	0	10	0	0
8	A	1	0	0	0	0
9	A	660	0	0	20	0
All	All	5206	0	4498	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:2540:GD8:O2	4:A:2540:GD8:C2	1.66	1.39
1:A:353[A]:MET:CE	9:A:1198:HOH:O	1.69	1.36
1:A:353[A]:MET:CE	9:A:1449:HOH:O	1.69	1.35
1:A:353[A]:MET:SD	9:A:1198:HOH:O	1.95	1.19
1:A:43[B]:MET:HE1	9:A:1066:HOH:O	1.40	1.18
1:A:353[A]:MET:HE1	9:A:1449:HOH:O	1.34	0.95
1:A:350[B]:GLU:HB3	9:A:1448:HOH:O	1.67	0.94
1:A:350[A]:GLU:HB2	9:A:1448:HOH:O	1.78	0.83
1:A:353[A]:MET:HE2	9:A:1449:HOH:O	1.54	0.77
1:A:321:LEU:HD11	1:A:408[A]:VAL:HG12	1.68	0.76
1:A:43[B]:MET:CE	9:A:1066:HOH:O	2.13	0.70
1:A:250[B]:ARG:HG2	1:A:256:LEU:HD11	1.77	0.67
1:A:350[A]:GLU:CG	9:A:1448:HOH:O	2.45	0.64
1:A:43[B]:MET:SD	9:A:1066:HOH:O	2.53	0.62
1:A:409:ASN:HB3	9:A:1506:HOH:O	2.01	0.61
1:A:504[B]:ASP:OD2	1:A:513:HIS:NE2	2.30	0.61
1:A:350[A]:GLU:CB	9:A:1448:HOH:O	2.45	0.59
1:A:440:HIS:HD2	4:A:2540:GD8:H6B	1.67	0.58
1:A:353[A]:MET:HE1	9:A:1198:HOH:O	1.65	0.56
1:A:350[A]:GLU:OE1	9:A:1448:HOH:O	2.17	0.56
1:A:316:LYS:HE3	1:A:414:PHE:HB3	1.90	0.52
1:A:350[B]:GLU:CB	9:A:1448:HOH:O	2.39	0.51
1:A:451:PRO:HA	1:A:458:TYR:CD2	2.46	0.51
1:A:485:PRO:HD3	9:A:1261:HOH:O	2.10	0.50
1:A:349:ARG:O	1:A:353[B]:MET:HG2	2.11	0.50
1:A:238:VAL:HG12	1:A:295[B]:VAL:HG11	1.94	0.49
1:A:223:ILE:HA	1:A:320:LEU:O	2.12	0.49
1:A:238:VAL:HG12	1:A:295[B]:VAL:CG1	2.43	0.48
1:A:419:TYR:CZ	1:A:494:LEU:HD13	2.49	0.48
1:A:197:PHE:CB	1:A:223:ILE:HB	2.45	0.46
1:A:185:GLN:HA	1:A:189:GLY:O	2.15	0.46
1:A:353[A]:MET:HE3	9:A:1198:HOH:O	1.68	0.45
1:A:405:MET:HA	1:A:408[A]:VAL:HG22	1.97	0.45
1:A:250[B]:ARG:CG	1:A:256:LEU:HD11	2.47	0.44
1:A:502:PHE:CZ	1:A:513:HIS:HB2	2.54	0.42
1:A:216:ARG:HG2	1:A:315:LYS:HB2	2.02	0.41
1:A:207:GLY:HA3	1:A:229:PRO:HD3	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:520:MET:HE2	9:A:1612:HOH:O	2.20	0.40
1:A:197:PHE:HB3	1:A:223:ILE:HB	2.02	0.40
1:A:199:GLU:HG3	1:A:443:GLU:OE2	2.22	0.40
6:A:2545:PG4:H21	9:A:1048:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	557/537 (104%)	536 (96%)	21 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	492/469 (105%)	485 (99%)	7 (1%)	67	62

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	197	PHE
1	A	288	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	294	PRO
1	A	330	PHE
1	A	430	LEU
1	A	473	TRP
1	A	487	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	68	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	B	1	2,1	14,14,15	0.50	0	17,19,21	1.03	0
2	FUC	B	2	2	10,10,11	0.68	0	14,14,16	0.62	0
3	NAG	C	1	3,1	14,14,15	0.54	0	17,19,21	0.77	0
3	NAG	C	2	3	14,14,15	0.56	0	17,19,21	0.89	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2,1	-	2/6/23/26	0/1/1/1
2	FUC	B	2	2	-	-	0/1/1/1
3	NAG	C	1	3,1	-	2/6/23/26	0/1/1/1
3	NAG	C	2	3	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	2	NAG	C1-O5-C5	2.62	115.74	112.19

There are no chirality outliers.



All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1	NAG	C4-C5-C6-O6
3	C	1	NAG	C4-C5-C6-O6
2	B	1	NAG	O5-C5-C6-O6
3	C	1	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry ⓘ

Of 10 ligands modelled in this entry, 1 is monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	PG4	A	2544	-	12,12,12	0.60	0	11,11,11	0.55	0
6	PG4	A	2543	-	12,12,12	0.57	0	11,11,11	0.67	0
6	PG4	A	2547	-	12,12,12	0.60	0	11,11,11	0.68	0
6	PG4	A	2545	-	12,12,12	0.53	0	11,11,11	0.42	0
5	TRS	A	2541	-	7,7,7	0.45	0	9,9,9	0.67	0
6	PG4	A	2546	-	12,12,12	0.49	0	11,11,11	0.51	0
7	PEG	A	2548	-	6,6,6	0.75	0	5,5,5	0.30	0
5	TRS	A	2542	-	7,7,7	0.44	0	9,9,9	0.46	0
4	GD8	A	2540	1	6,9,9	6.14	1 (16%)	7,13,13	1.65	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PG4	A	2544	-	-	1/10/10/10	-
6	PG4	A	2543	-	-	4/10/10/10	-
6	PG4	A	2547	-	-	5/10/10/10	-
6	PG4	A	2545	-	-	6/10/10/10	-
5	TRS	A	2541	-	-	2/9/9/9	-
6	PG4	A	2546	-	-	4/10/10/10	-
7	PEG	A	2548	-	-	2/4/4/4	-
5	TRS	A	2542	-	-	3/9/9/9	-
4	GD8	A	2540	1	-	5/6/10/10	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	2540	GD8	O2-C2	15.00	1.66	1.45

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	2540	GD8	O2-C2-C4	4.18	116.68	108.41

There are no chirality outliers.

All (32) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	2541	TRS	N-C-C2-O2
6	A	2543	PG4	O2-C3-C4-O3
6	A	2546	PG4	O4-C7-C8-O5
6	A	2547	PG4	O3-C5-C6-O4
6	A	2545	PG4	O4-C7-C8-O5
7	A	2548	PEG	O2-C3-C4-O4
6	A	2547	PG4	O4-C7-C8-O5
4	A	2540	GD8	O2-C2-C4-C7
4	A	2540	GD8	O2-C2-C4-C6
4	A	2540	GD8	O2-C2-C4-C5
6	A	2547	PG4	O2-C3-C4-O3
7	A	2548	PEG	O1-C1-C2-O2
6	A	2545	PG4	C1-C2-O2-C3
5	A	2542	TRS	C2-C-C3-O3
6	A	2543	PG4	C4-C3-O2-C2
6	A	2545	PG4	O1-C1-C2-O2
4	A	2540	GD8	C3-C2-C4-C6
6	A	2546	PG4	C8-C7-O4-C6
6	A	2547	PG4	C5-C6-O4-C7
6	A	2545	PG4	C5-C6-O4-C7
5	A	2541	TRS	C1-C-C2-O2
5	A	2542	TRS	C1-C-C3-O3
6	A	2546	PG4	C3-C4-O3-C5
6	A	2546	PG4	C1-C2-O2-C3
6	A	2547	PG4	C6-C5-O3-C4
6	A	2545	PG4	O3-C5-C6-O4
6	A	2543	PG4	C8-C7-O4-C6
6	A	2543	PG4	C1-C2-O2-C3
6	A	2545	PG4	C4-C3-O2-C2
6	A	2544	PG4	C3-C4-O3-C5
5	A	2542	TRS	N-C-C3-O3
4	A	2540	GD8	C3-C2-C4-C7

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	2545	PG4	1	0
4	A	2540	GD8	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	532/537 (99%)	-0.10	16 (3%) 50 59	18, 27, 40, 55	7 (1%)

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	486[A]	HIS	6.4
1	A	535	THR	6.2
1	A	488[A]	GLN	5.0
1	A	485	PRO	3.5
1	A	490[A]	SER	2.7
1	A	196	ILE	2.5
1	A	257	ASN	2.5
1	A	177	LEU	2.4
1	A	180	VAL	2.3
1	A	55	SER	2.2
1	A	457	ASN	2.1
1	A	42[A]	ASN	2.1
1	A	114	TRP	2.1
1	A	487	SER	2.0
1	A	218	LEU	2.0
1	A	264	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates ⓘ


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	C	2	14/15	0.79	0.40	59,63,64,64	0
2	FUC	B	2	10/11	0.80	0.44	78,81,82,82	0
3	NAG	C	1	14/15	0.91	0.16	38,42,50,55	0
2	NAG	B	1	14/15	0.91	0.24	55,60,64,70	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	PG4	A	2544	13/13	0.78	0.24	51,54,66,66	0
6	PG4	A	2546	13/13	0.82	0.25	62,64,69,70	0
7	PEG	A	2548	7/7	0.86	0.27	58,62,66,67	0
6	PG4	A	2545	13/13	0.87	0.25	52,54,66,67	0
6	PG4	A	2543	13/13	0.88	0.34	51,58,66,66	0
8	CL	A	2549	1/1	0.88	0.12	77,77,77,77	0
5	TRS	A	2541	8/8	0.90	0.21	33,35,36,37	0
6	PG4	A	2547	13/13	0.90	0.37	52,58,63,64	0
5	TRS	A	2542	8/8	0.92	0.37	59,59,59,60	0
4	GD8	A	2540	10/10	0.97	0.20	26,32,35,36	0

6.5 Other polymers [i](#)

There are no such residues in this entry.