



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 14, 2020 – 06:04 pm BST

PDB ID : 2WHS  
Title : Fluorescent Protein mKeima at pH 3.8  
Authors : Violot, S.; Carpentier, P.; Blanchoin, L.; Bourgeois, D.  
Deposited on : 2009-05-06  
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	240	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 3%, orange 3%, yellow 13%, green 75%, grey 10%);"></div> <div style="display: flex; justify-content: space-between; font-size: 8px;"> <span>3%</span> <span>75%</span> <span>13%</span> <span>• 10%</span> </div> </div>
1	B	240	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 3%, orange 3%, yellow 12%, green 81%, grey 5%);"></div> <div style="display: flex; justify-content: space-between; font-size: 8px;"> <span>3%</span> <span>81%</span> <span>12%</span> <span>• 5%</span> </div> </div>
1	C	240	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, yellow 11%, green 78%, grey 10%);"></div> <div style="display: flex; justify-content: space-between; font-size: 8px;"> <span></span> <span>78%</span> <span>11%</span> <span>• 10%</span> </div> </div>
1	D	240	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 3%, orange 3%, yellow 14%, green 80%, grey 5%);"></div> <div style="display: flex; justify-content: space-between; font-size: 8px;"> <span>3%</span> <span>80%</span> <span>14%</span> <span>• 5%</span> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	C	1221	-	-	X	-

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7919 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LARGE STOKES SHIFT FLUORESCENT PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	217	Total	C	N	O	S	0	0	0
			1740	1107	291	331	11			
1	B	227	Total	C	N	O	S	0	0	0
			1813	1153	306	342	12			
1	C	216	Total	C	N	O	S	0	0	0
			1733	1102	290	330	11			
1	D	227	Total	C	N	O	S	0	1	0
			1817	1155	306	344	12			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	62	CRQ	GLN	chromophore	UNP Q1JV70
A	62	CRQ	TYR	chromophore	UNP Q1JV70
A	62	CRQ	GLY	chromophore	UNP Q1JV70
B	62	CRQ	GLN	chromophore	UNP Q1JV70
B	62	CRQ	TYR	chromophore	UNP Q1JV70
B	62	CRQ	GLY	chromophore	UNP Q1JV70
C	62	CRQ	GLN	chromophore	UNP Q1JV70
C	62	CRQ	TYR	chromophore	UNP Q1JV70
C	62	CRQ	GLY	chromophore	UNP Q1JV70
D	62	CRQ	GLN	chromophore	UNP Q1JV70
D	62	CRQ	TYR	chromophore	UNP Q1JV70
D	62	CRQ	GLY	chromophore	UNP Q1JV70

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	159	Total	O	0	0
			159	159		

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	222	Total 222	O 222	0	0
3	C	166	Total 166	O 166	0	0
3	D	214	Total 214	O 214	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: LARGE STOKES SHIFT FLUORESCENT PROTEIN

Chain A: 





#### • Molecule 1: LARGE STOKES SHIFT FLUORESCENT PROTEIN

Chain B: 





#### • Molecule 1: LARGE STOKES SHIFT FLUORESCENT PROTEIN

Chain C: 



#### • Molecule 1: LARGE STOKES SHIFT FLUORESCENT PROTEIN

Chain D: 





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	92.57Å 98.10Å 123.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	76.70 – 2.10 46.28 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.2 (76.70-2.10) 99.2 (46.28-2.10)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.73 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.190 , 0.248 0.193 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	20.6	Xtriage
Anisotropy	1.084	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 49.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7919	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 46.90 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.0715e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: CRQ, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.79	0/1761	0.78	1/2379 (0.0%)
1	B	0.79	1/1836 (0.1%)	0.80	1/2480 (0.0%)
1	C	0.79	0/1754	0.81	0/2369
1	D	0.78	0/1843	0.82	0/2488
All	All	0.79	1/7194 (0.0%)	0.81	2/9716 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
All	All	0	3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	101	CYS	CB-SG	-8.73	1.67	1.82

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	219	LEU	CA-CB-CG	5.89	128.85	115.30
1	B	193	ARG	NE-CZ-NH2	5.60	123.10	120.30

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1	VAL	Peptide
1	B	61	LEU	Mainchain
1	C	61	LEU	Mainchain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1740	0	1676	45	0
1	B	1813	0	1754	25	0
1	C	1733	0	1664	23	0
1	D	1817	0	1759	22	0
2	A	15	0	0	0	0
2	B	15	0	0	0	0
2	C	10	0	0	3	0
2	D	15	0	0	0	0
3	A	159	0	0	2	0
3	B	222	0	0	3	0
3	C	166	0	0	2	0
3	D	214	0	0	5	0
All	All	7919	0	6853	105	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 105 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:212:ILE:HD11	1:A:214:ILE:CD1	1.65	1.25
1:A:212:ILE:HD11	1:A:214:ILE:HD12	1.03	1.02
1:A:212:ILE:CD1	1:A:214:ILE:HD12	1.95	0.95
1:C:38:GLN:HE22	1:C:65:SER:HB3	1.29	0.94
1:D:7:GLN:HB2	3:D:2017:HOH:O	1.70	0.92

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	214/240 (89%)	209 (98%)	4 (2%)	1 (0%)	29	26
1	B	224/240 (93%)	220 (98%)	4 (2%)	0	100	100
1	C	213/240 (89%)	206 (97%)	6 (3%)	1 (0%)	29	26
1	D	225/240 (94%)	221 (98%)	4 (2%)	0	100	100
All	All	876/960 (91%)	856 (98%)	18 (2%)	2 (0%)	47	49

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	2	SER
1	C	3	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	188/207 (91%)	181 (96%)	7 (4%)	34	35
1	B	196/207 (95%)	191 (97%)	5 (3%)	46	50
1	C	187/207 (90%)	180 (96%)	7 (4%)	34	35
1	D	197/207 (95%)	187 (95%)	10 (5%)	24	22
All	All	768/828 (93%)	739 (96%)	29 (4%)	33	34

5 of 29 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	105	ASN
1	C	153	LEU
1	D	199	SER
1	C	110	GLN
1	C	209	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 27 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	209	GLN
1	C	81	GLN
1	D	200	HIS
1	B	217	HIS
1	A	200	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	CRQ	D	62	1	24,25,26	3.88	6 (25%)	27,34,36	4.44	9 (33%)
1	CRQ	C	62	1	24,25,26	4.01	6 (25%)	27,34,36	5.14	7 (25%)
1	CRQ	B	62	1	24,25,26	3.72	6 (25%)	27,34,36	3.91	7 (25%)
1	CRQ	A	62	1	24,25,26	3.76	6 (25%)	27,34,36	4.44	6 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CRQ	D	62	1	-	1/10/32/33	0/2/2/2
1	CRQ	C	62	1	-	1/10/32/33	0/2/2/2
1	CRQ	B	62	1	-	1/10/32/33	0/2/2/2
1	CRQ	A	62	1	-	2/10/32/33	0/2/2/2

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	62	CRQ	CB2-CA2	17.65	1.49	1.35
1	D	62	CRQ	CB2-CA2	17.05	1.49	1.35
1	A	62	CRQ	CB2-CA2	16.58	1.49	1.35
1	B	62	CRQ	CB2-CA2	15.94	1.48	1.35
1	B	62	CRQ	CA2-C2	-4.91	1.43	1.48

The worst 5 of 29 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	62	CRQ	CA2-C2-N3	20.87	113.24	103.37
1	A	62	CRQ	CA2-C2-N3	16.84	111.33	103.37
1	D	62	CRQ	O2-C2-CA2	-16.63	121.62	130.96
1	D	62	CRQ	CA2-C2-N3	14.00	109.99	103.37
1	C	62	CRQ	O2-C2-CA2	-13.88	123.17	130.96

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	D	62	CRQ	C1-CA1-CB1-CG1
1	B	62	CRQ	C1-CA1-CB1-CG1
1	A	62	CRQ	CA1-CB1-CG1-CD3
1	C	62	CRQ	C-CA3-N3-C1
1	A	62	CRQ	C-CA3-N3-C1

There are no ring outliers.

4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	D	62	CRQ	2	0
1	C	62	CRQ	2	0
1	B	62	CRQ	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	62	CRQ	1	0

## 5.5 Carbohydrates

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	A	1220	-	4,4,4	0.14	0	6,6,6	0.15	0
2	SO4	C	1220	-	4,4,4	0.20	0	6,6,6	0.09	0
2	SO4	B	1222	-	4,4,4	0.29	0	6,6,6	0.42	0
2	SO4	C	1221	-	4,4,4	0.17	0	6,6,6	0.27	0
2	SO4	D	1222	-	4,4,4	0.32	0	6,6,6	0.72	0
2	SO4	A	1221	-	4,4,4	0.11	0	6,6,6	0.22	0
2	SO4	A	1222	-	4,4,4	0.18	0	6,6,6	0.34	0
2	SO4	D	1224	-	4,4,4	0.17	0	6,6,6	0.16	0
2	SO4	D	1223	-	4,4,4	0.13	0	6,6,6	0.16	0
2	SO4	B	1223	-	4,4,4	0.18	0	6,6,6	0.21	0
2	SO4	B	1224	-	4,4,4	0.15	0	6,6,6	0.16	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1221	SO4	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	216/240 (90%)	-0.08	3 (1%) 75 78	13, 21, 32, 39	1 (0%)
1	B	226/240 (94%)	-0.07	8 (3%) 44 50	12, 20, 35, 52	2 (0%)
1	C	215/240 (89%)	-0.17	1 (0%) 91 92	11, 20, 30, 44	1 (0%)
1	D	226/240 (94%)	-0.04	8 (3%) 44 50	14, 21, 37, 49	2 (0%)
All	All	883/960 (91%)	-0.09	20 (2%) 60 65	11, 20, 34, 52	6 (0%)

The worst 5 of 20 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	VAL	5.0
1	D	165	GLY	4.8
1	D	-1	HIS	4.6
1	A	165	GLY	4.4
1	B	165	GLY	4.2

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	CRQ	C	62	24/25	0.90	0.16	21,25,32,34	0
1	CRQ	D	62	24/25	0.92	0.14	20,23,32,33	0
1	CRQ	B	62	24/25	0.92	0.15	19,21,29,33	0
1	CRQ	A	62	24/25	0.92	0.15	18,22,29,31	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SO4	C	1221	5/5	0.81	0.58	93,94,94,95	0
2	SO4	C	1220	5/5	0.82	0.19	88,88,88,89	0
2	SO4	B	1223	5/5	0.87	0.30	90,91,91,91	0
2	SO4	D	1223	5/5	0.88	0.21	70,71,72,73	0
2	SO4	A	1221	5/5	0.88	0.19	86,86,87,87	0
2	SO4	A	1222	5/5	0.89	0.23	80,80,80,81	0
2	SO4	D	1224	5/5	0.92	0.16	90,90,90,90	0
2	SO4	A	1220	5/5	0.94	0.20	82,82,83,83	0
2	SO4	B	1224	5/5	0.96	0.18	79,79,79,79	0
2	SO4	D	1222	5/5	0.97	0.15	36,36,38,41	0
2	SO4	B	1222	5/5	0.99	0.11	34,36,40,41	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.