



# Full wwPDB X-ray Structure Validation Report ⓘ

May 25, 2022 – 08:13 PM JST

PDB ID : 7WHH  
Title : Crystal structure of SARS-CoV-2 omicron RBD and human ACE2  
Authors : Wang, X.Q.; Lan, J.  
Deposited on : 2021-12-30  
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.28.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.28.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	598	<div> <div>3%</div> <div>87%</div> <div>12%</div> <div>.</div> </div>
2	E	194	<div> <div>13%</div> <div>85%</div> <div>14%</div> <div>.</div> </div>
3	B	3	<div> <div>33%</div> <div>67%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	FUC	B	2	-	-	-	X

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 6667 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Processed angiotensin-converting enzyme 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	598	Total	C	N	O	S	0	1	0
			4884	3125	809	921	29			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	18	PRO	-	expression tag	UNP Q9BYF1


- Molecule 2 is a protein called Spike glycoprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	194	Total	C	N	O	S	0	0	0
			1556	1004	261	283	8			

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	339	ASP	GLY	variant	UNP P0DTC2
E	371	LEU	SER	variant	UNP P0DTC2
E	373	PRO	SER	variant	UNP P0DTC2
E	375	PHE	SER	variant	UNP P0DTC2
E	417	ASN	LYS	variant	UNP P0DTC2
E	440	LYS	ASN	variant	UNP P0DTC2
E	446	SER	GLY	variant	UNP P0DTC2
E	477	ASN	SER	variant	UNP P0DTC2
E	478	LYS	THR	variant	UNP P0DTC2
E	484	ALA	GLU	variant	UNP P0DTC2
E	493	LYS	GLN	variant	UNP P0DTC2
E	496	SER	GLY	variant	UNP P0DTC2
E	498	ARG	GLN	variant	UNP P0DTC2
E	501	TYR	ASN	variant	UNP P0DTC2
E	505	HIS	TYR	variant	UNP P0DTC2

- Molecule 3 is an oligosaccharide called alpha-L-fucopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	B	3	Total	C	N	O	0	0	0
			38	22	2	14			


- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Zn	0	0
			1	1		

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Cl	0	0
			1	1		

- Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	E	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	109	Total	O	0	0
			109	109		
7	E	22	Total	O	0	0
			22	22		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Processed angiotensin-converting enzyme 2



- Molecule 2: Spike glycoprotein



- Molecule 3: alpha-L-fucopyranose-(1-3)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)]2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	104.71 Å   104.71 Å   227.10 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	36.54 – 2.60 39.82 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.8 (36.54-2.60) 99.9 (39.82-2.60)	Depositor EDS
$R_{merge}$	0.26	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.69 (at 2.61 Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, $R_{free}$	0.192   ,   0.231 0.192   ,   0.230	Depositor DCC
$R_{free}$ test set	1937 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	47.1	Xtriage
Anisotropy	0.117	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 36.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	6667	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.29% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, FUC, NAG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.46	0/5026	0.57	0/6829
2	E	0.43	0/1602	0.58	0/2180
All	All	0.45	0/6628	0.57	0/9009

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4884	0	4654	42	0
2	E	1556	0	1486	15	0
3	B	38	0	34	1	0
4	A	1	0	0	0	0
5	A	1	0	0	0	0
6	A	42	0	39	4	0
6	E	14	0	13	0	0
7	A	109	0	0	9	0
7	E	22	0	0	2	0
All	All	6667	0	6226	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 5.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:904:NAG:O3	7:A:1001:HOH:O	1.85	0.94
6:A:904:NAG:O4	7:A:1002:HOH:O	1.87	0.92
7:E:722:HOH:O	3:B:3:NAG:O3	2.13	0.67
1:A:330:ASN:OD1	7:A:1003:HOH:O	2.13	0.65
1:A:460:ARG:HH21	1:A:506:VAL:HA	1.62	0.64
1:A:121:ASN:O	1:A:125:THR:HG23	2.04	0.58
1:A:477:TRP:CE3	1:A:500:PRO:HG3	2.40	0.57
1:A:402:GLU:HB3	1:A:518:ARG:HD3	1.87	0.56
1:A:215:TYR:CE1	1:A:577:LYS:HE2	2.41	0.55
1:A:474:MET:HE1	1:A:499:ASP:HB2	1.87	0.55
1:A:353:LYS:HD3	2:E:501:TYR:CZ	2.43	0.54
1:A:346:PRO:HB3	1:A:360:MET:HG3	1.89	0.54
2:E:476:GLY:H	2:E:487:ASN:HB3	1.73	0.53
1:A:185:VAL:O	7:A:1005:HOH:O	2.19	0.52
1:A:111:ASP:OD2	7:A:1006:HOH:O	2.19	0.52
1:A:192:ARG:NH2	1:A:197:GLU:O	2.43	0.52
2:E:354:ASN:O	2:E:398:ASP:HA	2.10	0.52
1:A:320:LEU:HD13	1:A:380:GLN:HG2	1.92	0.51
1:A:586:ASN:OD1	7:A:1004:HOH:O	2.18	0.51
2:E:502:GLY:O	2:E:506:GLN:HG3	2.11	0.50
1:A:197:GLU:HG2	1:A:201:ASP:OD2	2.12	0.50
1:A:233:ILE:HD13	1:A:450:LEU:HD13	1.94	0.50
1:A:31:LYS:HE2	1:A:35:GLU:OE2	2.14	0.48
2:E:496:SER:OG	2:E:498:ARG:NH1	2.47	0.47
1:A:594:TRP:CH2	1:A:598:GLN:HG3	2.49	0.47
1:A:185:VAL:O	1:A:189:GLU:HG3	2.14	0.47
1:A:402:GLU:HB3	1:A:518:ARG:CD	2.44	0.47
6:A:904:NAG:O6	7:A:1007:HOH:O	2.20	0.46
1:A:460:ARG:NH2	1:A:506:VAL:HA	2.29	0.46
2:E:368:LEU:HA	2:E:368:LEU:HD13	1.74	0.46
1:A:229:THR:HB	1:A:581:VAL:HG13	1.97	0.46
2:E:383:SER:HB3	2:E:386:LYS:HD2	1.98	0.46
2:E:393:THR:HA	2:E:522:ALA:HA	1.97	0.46
2:E:393:THR:HG21	2:E:518:LEU:H	1.80	0.45
1:A:134:ASN:O	1:A:136:ASP:N	2.48	0.45
2:E:368:LEU:O	2:E:369:TYR:HB2	2.17	0.44
1:A:50:TYR:CE1	1:A:54:ILE:HG23	2.53	0.44
1:A:538:PRO:HD2	1:A:541:LYS:HD3	1.99	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:37:GLU:HG2	1:A:353:LYS:HE3	1.99	0.43
1:A:237:TYR:CZ	1:A:451:PRO:HG2	2.54	0.43
1:A:134:ASN:C	1:A:136:ASP:H	2.20	0.43
1:A:115:ARG:NE	7:A:1018:HOH:O	2.51	0.43
2:E:369:TYR:OH	2:E:384:PRO:HG2	2.19	0.42
2:E:399:SER:HA	2:E:510:VAL:O	2.20	0.42
1:A:39:LEU:HD23	1:A:39:LEU:HA	1.73	0.42
1:A:155:SER:O	1:A:161:ARG:HD2	2.19	0.42
1:A:340:GLN:OE1	6:A:905:NAG:H82	2.19	0.42
1:A:234:LYS:O	1:A:238:GLU:HG3	2.20	0.42
1:A:144:LEU:HA	1:A:148:LEU:HB2	2.02	0.42
1:A:252:TYR:HB2	1:A:256:ILE:HD12	2.02	0.42
1:A:355:ASP:HA	7:A:1026:HOH:O	2.20	0.42
1:A:524:GLN:HG2	1:A:583:PRO:HG2	2.02	0.42
1:A:320:LEU:HD13	1:A:380:GLN:CG	2.50	0.41
1:A:21:ILE:HG21	1:A:84:PRO:HD2	2.01	0.41
2:E:351:TYR:O	7:E:701:HOH:O	2.22	0.41
2:E:393:THR:HG23	2:E:516:GLU:O	2.21	0.41
1:A:215:TYR:CZ	1:A:568:LEU:HD13	2.56	0.41
1:A:455:MET:HE2	1:A:485:VAL:HG21	2.02	0.41
1:A:41:TYR:CD1	1:A:353:LYS:HG3	2.56	0.40
2:E:374:PHE:O	2:E:436:TRP:HA	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	597/598 (100%)	584 (98%)	13 (2%)	0	100	100
2	E	192/194 (99%)	181 (94%)	11 (6%)	0	100	100
All	All	789/792 (100%)	765 (97%)	24 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	529/528 (100%)	512 (97%)	17 (3%)	39	65
2	E	169/169 (100%)	160 (95%)	9 (5%)	22	45
All	All	698/697 (100%)	672 (96%)	26 (4%)	34	60

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47	SER
1	A	125	THR
1	A	139	GLN
1	A	170	SER
1	A	190	MET
1	A	224	GLU
1	A	257	SER
1	A	283	VAL
1	A	340	GLN
1	A	381	TYR
1	A	385	TYR
1	A	394	ASN
1	A	401	HIS
1	A	455	MET
1	A	511	SER
1	A	581	VAL
1	A	607	SER
2	E	333	THR
2	E	335	LEU
2	E	354	ASN
2	E	376	THR
2	E	377	PHE
2	E	388	ASN
2	E	393	THR
2	E	408	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	E	444	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAG	B	1	3,1	14,14,15	0.49	0	17,19,21	0.59	0
3	FUC	B	2	3	10,10,11	1.77	3 (30%)	14,14,16	1.35	2 (14%)
3	NAG	B	3	3	14,14,15	0.54	0	17,19,21	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	B	1	3,1	-	2/6/23/26	0/1/1/1
3	FUC	B	2	3	-	-	0/1/1/1
3	NAG	B	3	3	-	1/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	2	FUC	O5-C5	3.23	1.50	1.43
3	B	2	FUC	C2-C3	2.46	1.56	1.52
3	B	2	FUC	C1-C2	2.17	1.57	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	2	FUC	C1-O5-C5	2.97	119.50	112.78
3	B	2	FUC	O5-C5-C4	2.56	114.12	109.52

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	1	NAG	O5-C5-C6-O6
3	B	1	NAG	C4-C5-C6-O6
3	B	3	NAG	O5-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	3	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	NAG	E	601	2	14,14,15	0.87	1 (7%)	17,19,21	0.68	0
6	NAG	A	905	1	14,14,15	0.34	0	17,19,21	0.65	1 (5%)
6	NAG	A	904	1	14,14,15	0.84	1 (7%)	17,19,21	0.81	1 (5%)
6	NAG	A	903	1	14,14,15	0.42	0	17,19,21	1.03	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	E	601	2	-	2/6/23/26	0/1/1/1
6	NAG	A	905	1	-	2/6/23/26	0/1/1/1
6	NAG	A	904	1	-	1/6/23/26	0/1/1/1
6	NAG	A	903	1	-	0/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	E	601	NAG	C1-C2	2.89	1.56	1.52
6	A	904	NAG	O5-C1	2.65	1.47	1.43

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	903	NAG	C1-O5-C5	3.65	117.13	112.19
6	A	904	NAG	C1-O5-C5	2.85	116.05	112.19
6	A	905	NAG	C1-O5-C5	2.25	115.23	112.19

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	905	NAG	O5-C5-C6-O6
6	A	905	NAG	C4-C5-C6-O6
6	E	601	NAG	C4-C5-C6-O6
6	E	601	NAG	O5-C5-C6-O6
6	A	904	NAG	C4-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	905	NAG	1	0
6	A	904	NAG	3	0



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	598/598 (100%)	-0.14	16 (2%) 54 48	26, 42, 74, 104	0
2	E	194/194 (100%)	0.55	26 (13%) 3 2	38, 61, 112, 126	0
All	All	792/792 (100%)	0.03	42 (5%) 26 20	26, 46, 94, 126	0

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	339	VAL	6.0
2	E	374	PHE	5.8
2	E	367	VAL	5.4
2	E	368	LEU	4.9
1	A	338	ASN	4.4
1	A	615	ASP	4.3
2	E	517	LEU	4.2
1	A	136	ASP	4.1
1	A	105	SER	3.9
2	E	365	TYR	3.9
1	A	86	GLN	3.9
2	E	391	CYS	3.7
2	E	385	THR	3.5
2	E	520	ALA	3.4
2	E	381	GLY	3.3
2	E	371	LEU	3.2
2	E	515	PHE	3.2
2	E	430	THR	3.2
2	E	524	VAL	3.1
2	E	364	ASP	3.0
2	E	373	PRO	2.9
1	A	138	PRO	2.9
2	E	335	LEU	2.9
2	E	390	LEU	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	135	PRO	2.7
1	A	104	GLY	2.6
2	E	384	PRO	2.5
2	E	389	ASP	2.4
1	A	340	GLN	2.4
2	E	366	SER	2.4
2	E	392	PHE	2.4
1	A	139	GLN	2.3
1	A	341	LYS	2.2
2	E	387	LEU	2.1
1	A	50	TYR	2.1
2	E	386	LYS	2.1
1	A	57	GLU	2.1
2	E	426	PRO	2.1
2	E	525	CYS	2.1
1	A	425	SER	2.1
1	A	427	ASP	2.1
2	E	516	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	FUC	B	2	10/11	0.78	0.41	91,96,103,103	0
3	NAG	B	3	14/15	0.79	0.36	86,96,99,100	0
3	NAG	B	1	14/15	0.93	0.14	74,82,89,93	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
6	NAG	E	601	14/15	0.56	0.32	101,113,119,119	0
6	NAG	A	904	14/15	0.62	0.27	77,85,87,88	0
6	NAG	A	903	14/15	0.77	0.35	73,81,85,86	0
6	NAG	A	905	14/15	0.84	0.39	78,83,87,88	0
4	ZN	A	901	1/1	0.94	0.11	66,66,66,66	0
5	CL	A	902	1/1	0.98	0.13	36,36,36,36	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.