



Full wwPDB X-ray Structure Validation Report ⓘ

May 23, 2020 – 03:27 am BST

PDB ID : 5WJ2
Title : Crystal structure of the green fluorescent protein Clover
Authors : Liu, C.; Campbell, B.C.; Petsko, G.A.
Deposited on : 2017-07-21
Resolution : 2.41 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.41 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	258	<div> <div>5%</div> <div> <div></div> <div>83%</div> <div>9%</div> <div>9%</div> </div> </div>
1	B	258	<div> <div>5%</div> <div> <div></div> <div>82%</div> <div>6%</div> <div>12%</div> </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3768 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Green fluorescent protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	235	Total	C	N	O	S	0	0	0
			1875	1191	322	356	6			
1	B	227	Total	C	N	O	S	0	0	0
			1805	1145	310	345	5			

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-21	MET	-	initiating methionine	UNP P42212
A	-20	GLY	-	expression tag	UNP P42212
A	-19	SER	-	expression tag	UNP P42212
A	-18	SER	-	expression tag	UNP P42212
A	-17	HIS	-	expression tag	UNP P42212
A	-16	HIS	-	expression tag	UNP P42212
A	-15	HIS	-	expression tag	UNP P42212
A	-14	HIS	-	expression tag	UNP P42212
A	-13	HIS	-	expression tag	UNP P42212
A	-12	HIS	-	expression tag	UNP P42212
A	-11	SER	-	expression tag	UNP P42212
A	-10	SER	-	expression tag	UNP P42212
A	-9	GLY	-	expression tag	UNP P42212
A	-8	GLU	-	expression tag	UNP P42212
A	-7	ASN	-	expression tag	UNP P42212
A	-6	LEU	-	expression tag	UNP P42212
A	-5	TYR	-	expression tag	UNP P42212
A	-4	PHE	-	expression tag	UNP P42212
A	-3	GLN	-	expression tag	UNP P42212
A	-2	GLY	-	expression tag	UNP P42212
A	-1	HIS	-	expression tag	UNP P42212
A	0	MET	-	expression tag	UNP P42212
A	1	VAL	-	expression tag	UNP P42212
A	30	ARG	SER	engineered mutation	UNP P42212
A	39	ASN	TYR	engineered mutation	UNP P42212

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	CR2	SER	chromophore	UNP P42212
A	?	CR2	TYR	chromophore	UNP P42212
A	66	CR2	GLY	chromophore	UNP P42212
A	69	ALA	GLN	engineered mutation	UNP P42212
A	99	SER	PHE	engineered mutation	UNP P42212
A	105	THR	ASN	engineered mutation	UNP P42212
A	145	PHE	TYR	engineered mutation	UNP P42212
A	153	THR	MET	engineered mutation	UNP P42212
A	163	ALA	VAL	engineered mutation	UNP P42212
A	171	VAL	ILE	engineered mutation	UNP P42212
A	203	HIS	THR	engineered mutation	UNP P42212
B	-21	MET	-	initiating methionine	UNP P42212
B	-20	GLY	-	expression tag	UNP P42212
B	-19	SER	-	expression tag	UNP P42212
B	-18	SER	-	expression tag	UNP P42212
B	-17	HIS	-	expression tag	UNP P42212
B	-16	HIS	-	expression tag	UNP P42212
B	-15	HIS	-	expression tag	UNP P42212
B	-14	HIS	-	expression tag	UNP P42212
B	-13	HIS	-	expression tag	UNP P42212
B	-12	HIS	-	expression tag	UNP P42212
B	-11	SER	-	expression tag	UNP P42212
B	-10	SER	-	expression tag	UNP P42212
B	-9	GLY	-	expression tag	UNP P42212
B	-8	GLU	-	expression tag	UNP P42212
B	-7	ASN	-	expression tag	UNP P42212
B	-6	LEU	-	expression tag	UNP P42212
B	-5	TYR	-	expression tag	UNP P42212
B	-4	PHE	-	expression tag	UNP P42212
B	-3	GLN	-	expression tag	UNP P42212
B	-2	GLY	-	expression tag	UNP P42212
B	-1	HIS	-	expression tag	UNP P42212
B	0	MET	-	expression tag	UNP P42212
B	1	VAL	-	expression tag	UNP P42212
B	30	ARG	SER	engineered mutation	UNP P42212
B	39	ASN	TYR	engineered mutation	UNP P42212
B	?	CR2	SER	chromophore	UNP P42212
B	?	CR2	TYR	chromophore	UNP P42212
B	66	CR2	GLY	chromophore	UNP P42212
B	69	ALA	GLN	engineered mutation	UNP P42212
B	99	SER	PHE	engineered mutation	UNP P42212
B	105	THR	ASN	engineered mutation	UNP P42212

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	145	PHE	TYR	engineered mutation	UNP P42212
B	153	THR	MET	engineered mutation	UNP P42212
B	163	ALA	VAL	engineered mutation	UNP P42212
B	171	VAL	ILE	engineered mutation	UNP P42212
B	203	HIS	THR	engineered mutation	UNP P42212

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	60	Total O 60 60	0	0
2	B	28	Total O 28 28	0	0

- Molecule 1: Green fluorescent protein



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	53.06Å 135.18Å 168.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.15 – 2.41 42.15 – 2.41	Depositor EDS
% Data completeness (in resolution range)	99.6 (42.15-2.41) 99.6 (42.15-2.41)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.57 (at 2.42Å)	Xtriage
Refinement program	PHENIX (1.11.1 _2575: ???)	Depositor
R, R_{free}	0.193 , 0.217 0.193 , 0.217	Depositor DCC
R_{free} test set	1161 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å ²)	41.5	Xtriage
Anisotropy	0.431	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 50.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3768	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.02% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CR2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/1899	0.49	0/2565
1	B	0.26	0/1826	0.49	0/2467
All	All	0.26	0/3725	0.49	0/5032

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1875	0	1815	13	0
1	B	1805	0	1753	9	0
2	A	60	0	0	2	1
2	B	28	0	0	1	0
All	All	3768	0	3568	21	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:162:LYS:NZ	2:A:301:HOH:O	2.21	0.69
1:A:45:LYS:HD2	1:A:219:VAL:HG22	1.77	0.65
1:B:158:LYS:NZ	1:B:184:GLN:OE1	2.36	0.58
1:B:98:ILE:HB	1:B:106:TYR:HB2	1.87	0.57
1:A:161:ILE:HD11	1:A:196:PRO:CD	2.38	0.54
1:A:166:LYS:HD3	1:A:178:LEU:HD21	1.91	0.53
1:A:214:LYS:HE3	2:A:345:HOH:O	2.10	0.52
1:B:184:GLN:NE2	2:B:305:HOH:O	2.43	0.52
1:A:161:ILE:HD11	1:A:196:PRO:HD2	1.92	0.51
1:B:161:ILE:HD11	1:B:196:PRO:HD2	1.93	0.51
1:A:66:CR2:HD2	1:A:66:CR2:N2	2.27	0.49
1:B:161:ILE:HD11	1:B:196:PRO:CD	2.42	0.49
1:B:46:PHE:CZ	1:B:64:PHE:HB3	2.48	0.49
1:B:66:CR2:N2	1:B:66:CR2:HD1	2.29	0.48
1:A:98:ILE:HB	1:A:106:TYR:HB2	1.99	0.45
1:A:46:PHE:CZ	1:A:64:PHE:HB3	2.52	0.44
1:A:223:PHE:HB3	1:B:204:GLN:OE1	2.17	0.43
1:A:63:THR:CG2	1:A:123:ILE:HG21	2.48	0.43
1:B:121:ASN:ND2	1:B:123:ILE:HD11	2.36	0.41
1:A:167:ILE:HB	1:A:179:ALA:HB3	2.03	0.40
1:A:83:PHE:CG	1:A:161:ILE:HD13	2.56	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:314:HOH:O	2:A:357:HOH:O[8_555]	2.15	0.05

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	230/258 (89%)	227 (99%)	3 (1%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	222/258 (86%)	217 (98%)	4 (2%)	1 (0%)	29	41
All	All	452/516 (88%)	444 (98%)	7 (2%)	1 (0%)	47	62

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	2	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	203/223 (91%)	200 (98%)	3 (2%)	65	80
1	B	196/223 (88%)	194 (99%)	2 (1%)	76	88
All	All	399/446 (90%)	394 (99%)	5 (1%)	69	84

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	GLU
1	A	157	GLN
1	A	222	GLU
1	B	6	GLU
1	B	178	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CR2	B	66	1	20,20,21	4.05	4 (20%)	25,27,29	4.58	10 (40%)
1	CR2	A	66	1	20,20,21	3.81	5 (25%)	25,27,29	4.17	11 (44%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CR2	B	66	1	-	2/6/25/26	0/2/2/2
1	CR2	A	66	1	-	1/6/25/26	0/2/2/2

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	66	CR2	CB2-CA2	17.06	1.49	1.35
1	A	66	CR2	CB2-CA2	15.78	1.48	1.35
1	A	66	CR2	C1-N2	3.41	1.38	1.32
1	B	66	CR2	C1-N2	3.30	1.38	1.32
1	B	66	CR2	C2-N3	-3.00	1.32	1.39
1	A	66	CR2	C2-N3	-2.81	1.33	1.39
1	B	66	CR2	O2-C2	2.76	1.28	1.23
1	A	66	CR2	O2-C2	2.33	1.28	1.23
1	A	66	CR2	CA2-C2	-2.20	1.46	1.48

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	66	CR2	CA2-C2-N3	14.98	110.45	103.37
1	A	66	CR2	CA2-C2-N3	12.70	109.38	103.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	66	CR2	O2-C2-CA2	-12.61	123.88	130.96
1	B	66	CR2	O2-C2-CA2	-11.04	124.76	130.96
1	B	66	CR2	CB2-CA2-C2	7.16	130.82	122.28
1	B	66	CR2	C2-N3-C1	-6.77	104.68	107.99
1	B	66	CR2	C2-CA2-N2	-5.08	105.37	108.93
1	A	66	CR2	C2-N3-C1	-4.70	105.69	107.99
1	A	66	CR2	CB2-CA2-C2	4.69	127.87	122.28
1	A	66	CR2	O3-C3-CA3	-4.24	113.59	126.39
1	B	66	CR2	CB2-CA2-N2	-3.63	123.80	128.83
1	A	66	CR2	C2-CA2-N2	-3.47	106.50	108.93
1	B	66	CR2	CG2-CB2-CA2	-3.29	125.92	129.94
1	A	66	CR2	CG2-CB2-CA2	-3.08	126.17	129.94
1	B	66	CR2	C1-CA1-N1	-3.07	106.05	112.85
1	B	66	CR2	O3-C3-CA3	-2.63	118.45	126.39
1	B	66	CR2	CA3-N3-C2	2.50	129.54	123.80
1	A	66	CR2	C1-CA1-N1	-2.48	107.37	112.85
1	A	66	CR2	CA3-N3-C2	2.43	129.36	123.80
1	A	66	CR2	CB2-CA2-N2	-2.31	125.62	128.83
1	A	66	CR2	CA3-N3-C1	-2.13	124.92	127.86

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	66	CR2	C3-CA3-N3-C1
1	B	66	CR2	C3-CA3-N3-C2
1	A	66	CR2	C3-CA3-N3-C2

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	66	CR2	1	0
1	A	66	CR2	1	0

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	234/258 (90%)	0.22	12 (5%) 28 26	25, 43, 75, 98	0
1	B	226/258 (87%)	0.21	14 (6%) 20 19	31, 58, 91, 106	0
All	All	460/516 (89%)	0.21	26 (5%) 23 22	25, 51, 84, 106	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	155	ASP	5.2
1	A	156	LYS	4.5
1	B	133	ASP	4.4
1	A	157	GLN	4.2
1	A	133	ASP	4.0
1	B	117	ASP	3.9
1	B	229	ILE	3.8
1	B	228	GLY	3.8
1	A	158	LYS	3.6
1	A	182	TYR	3.6
1	B	132	GLU	3.5
1	B	3	LYS	3.3
1	B	80	GLN	3.1
1	A	229	ILE	2.9
1	B	79	LYS	2.9
1	B	4	GLY	2.8
1	B	156	LYS	2.7
1	B	6	GLU	2.4
1	B	30	ARG	2.3
1	A	102	ASP	2.2
1	A	173	ASP	2.2
1	A	68	VAL	2.1
1	A	42	LEU	2.1
1	A	174	GLY	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	51	GLY	2.1
1	B	2	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CR2	B	66	19/20	0.95	0.20	35,44,69,74	0
1	CR2	A	66	19/20	0.96	0.20	31,38,61,72	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.