



# Full wwPDB X-ray Structure Validation Report ⓘ

May 24, 2020 – 10:40 pm BST

PDB ID : 5WJ4  
Title : Crystal structure of redox-sensitive green fluorescent protein Clover mutant roClover1  
Authors : Liu, C.; Campbell, B.C.; Petsko, G.A.  
Deposited on : 2017-07-21  
Resolution : 1.63 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.63 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3122 (1.66-1.62)
Clashscore	141614	3268 (1.66-1.62)
Ramachandran outliers	138981	3215 (1.66-1.62)
Sidechain outliers	138945	3215 (1.66-1.62)
RSRZ outliers	127900	3079 (1.66-1.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	239	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%;"></div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 10px;"> <div style="width: 1%; height: 10px; background-color: red;"></div> <div style="width: 87%, height: 10px; background-color: green;"></div> <div style="width: 8%, height: 10px; background-color: yellow;"></div> <div style="width: 5%, height: 10px; background-color: grey;"></div> </div> </div> </div>
1	B	239	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0; width: 100%;"></div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 10px;"> <div style="width: 2%, height: 10px; background-color: red;"></div> <div style="width: 86%, height: 10px; background-color: green;"></div> <div style="width: 10%, height: 10px; background-color: yellow;"></div> <div style="width: 5%, height: 10px; background-color: grey;"></div> </div> </div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4194 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Green fluorescent protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	228	Total	C	N	O	S	0	4	0
			1831	1160	312	352	7			
1	B	228	Total	C	N	O	S	0	2	0
			1816	1153	309	347	7			

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP P42212
A	-1	HIS	-	expression tag	UNP P42212
A	0	MET	-	expression tag	UNP P42212
A	1	VAL	-	expression tag	UNP P42212
A	30	ARG	SER	engineered mutation	UNP P42212
A	39	ASN	TYR	engineered mutation	UNP P42212
A	65	CR2	SER	chromophore	UNP P42212
A	65	CR2	TYR	chromophore	UNP P42212
A	65	CR2	GLY	chromophore	UNP P42212
A	67	ALA	GLN	engineered mutation	UNP P42212
A	97	SER	PHE	engineered mutation	UNP P42212
A	103	THR	ASN	engineered mutation	UNP P42212
A	143	PHE	TYR	engineered mutation	UNP P42212
A	145	CYS	SER	engineered mutation	UNP P42212
A	146	ASP	HIS	engineered mutation	UNP P42212
A	151	THR	MET	engineered mutation	UNP P42212
A	161	ALA	VAL	engineered mutation	UNP P42212
A	169	VAL	ILE	engineered mutation	UNP P42212
A	201	VAL	THR	engineered mutation	UNP P42212
A	202	CYS	GLN	engineered mutation	UNP P42212
A	220	GLN	GLU	engineered mutation	UNP P42212
B	-2	GLY	-	expression tag	UNP P42212
B	-1	HIS	-	expression tag	UNP P42212
B	0	MET	-	expression tag	UNP P42212
B	1	VAL	-	expression tag	UNP P42212

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	30	ARG	SER	engineered mutation	UNP P42212
B	39	ASN	TYR	engineered mutation	UNP P42212
B	65	CR2	SER	chromophore	UNP P42212
B	65	CR2	TYR	chromophore	UNP P42212
B	65	CR2	GLY	chromophore	UNP P42212
B	67	ALA	GLN	engineered mutation	UNP P42212
B	97	SER	PHE	engineered mutation	UNP P42212
B	103	THR	ASN	engineered mutation	UNP P42212
B	143	PHE	TYR	engineered mutation	UNP P42212
B	145	CYS	SER	engineered mutation	UNP P42212
B	146	ASP	HIS	engineered mutation	UNP P42212
B	151	THR	MET	engineered mutation	UNP P42212
B	161	ALA	VAL	engineered mutation	UNP P42212
B	169	VAL	ILE	engineered mutation	UNP P42212
B	201	VAL	THR	engineered mutation	UNP P42212
B	202	CYS	GLN	engineered mutation	UNP P42212
B	220	GLN	GLU	engineered mutation	UNP P42212



- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	293	Total O 293 293	0	0
2	B	254	Total O 254 254	0	0



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Green fluorescent protein



- Molecule 1: Green fluorescent protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	134.57Å 134.57Å 69.09Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.19 – 1.63 39.19 – 1.63	Depositor EDS
% Data completeness (in resolution range)	100.0 (39.19-1.63) 96.8 (39.19-1.63)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.51 (at 1.63Å)	Xtriage
Refinement program	PHENIX 1.10.1 _2155	Depositor
R, $R_{free}$	0.164 , 0.193 0.164 , 0.194	Depositor DCC
$R_{free}$ test set	3973 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.4	Xtriage
Anisotropy	0.153	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 45.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	4194	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.42% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CR2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.74	0/1863	0.83	2/2517 (0.1%)
1	B	0.71	1/1842 (0.1%)	0.80	2/2490 (0.1%)
All	All	0.73	1/3705 (0.0%)	0.82	4/5007 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	32	GLU	CD-OE2	-6.02	1.19	1.25

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	30	ARG	NE-CZ-NH2	-6.86	116.87	120.30
1	A	86	MET	CG-SD-CE	6.34	110.34	100.20
1	A	195	ASP	CB-CG-OD1	5.31	123.08	118.30
1	B	42	LEU	CB-CG-CD1	-5.25	102.07	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1831	0	1779	17	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1816	0	1770	15	0
2	A	293	0	0	4	3
2	B	254	0	0	3	4
All	All	4194	0	3549	32	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:156:LYS:HE2	1:B:184:THR:HG21	1.68	0.75
1:B:166:ARG:NH2	2:B:301:HOH:O	2.06	0.75
1:A:5:GLU:HG2	2:A:301:HOH:O	1.94	0.66
1:A:140:GLU:HG2	1:A:168:ASN:HB2	1.76	0.66
1:B:103[B]:THR:HG21	2:B:342:HOH:O	1.98	0.64
1:A:86:MET:HE3	1:A:112:PHE:CD1	2.37	0.60
1:B:43:THR:HG22	1:B:219[B]:LEU:HD22	1.83	0.58
1:B:72:TYR:OH	1:B:197:HIS:HE1	1.86	0.58
1:A:86:MET:CE	1:A:112:PHE:CD1	2.87	0.57
1:B:164:LYS:HG2	1:B:178:ASP:OD1	2.05	0.57
1:A:86:MET:HE3	1:A:112:PHE:HD1	1.72	0.54
1:B:155:GLN:NE2	1:B:156:LYS:HG3	2.22	0.54
1:A:2:SER:O	2:A:301:HOH:O	2.18	0.53
1:B:195:ASP:O	1:B:197:HIS:HD2	1.93	0.51
1:B:103[B]:THR:HG23	2:B:331:HOH:O	2.11	0.51
1:B:94:ARG:HE	1:B:181:GLN:NE2	2.10	0.50
1:B:97:SER:HA	1:B:103[B]:THR:HG22	1.94	0.48
1:A:164:LYS:HG2	1:A:178[B]:ASP:OD1	2.14	0.47
1:A:86:MET:HE2	1:A:112:PHE:CE1	2.50	0.47
1:A:82:PHE:CE1	1:A:159:ILE:HD11	2.50	0.47
1:A:65:CR2:HD1	1:A:65:CR2:N2	2.33	0.44
1:B:43:THR:HG22	1:B:219[B]:LEU:CD2	2.48	0.43
1:A:166[A]:ARG:HG2	1:A:176:LEU:CD2	2.49	0.43
1:B:153:ASP:OD1	1:B:155:GLN:OE1	2.36	0.43
1:A:67:ALA:HA	2:A:427:HOH:O	2.17	0.43
1:B:65:CR2:N2	1:B:65:CR2:HD1	2.34	0.43
1:A:160:LYS:HE2	1:A:182:GLN:OE1	2.19	0.42
1:A:140:GLU:CG	1:A:168:ASN:HB2	2.48	0.42
1:A:86:MET:HE2	1:A:112:PHE:CD1	2.55	0.42
1:A:228:THR:HG23	2:A:303:HOH:O	2.20	0.42

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:166[A]:ARG:HG2	1:A:176:LEU:HD23	2.02	0.41
1:B:45:LYS:HE2	1:B:47:ILE:HD11	2.03	0.41

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:314:HOH:O	2:B:314:HOH:O[8_555]	2.01	0.19
2:B:531:HOH:O	2:B:531:HOH:O[8_554]	2.14	0.06
2:A:529:HOH:O	2:B:411:HOH:O[4_455]	2.15	0.05
2:A:489:HOH:O	2:B:316:HOH:O[8_554]	2.16	0.04
2:A:502:HOH:O	2:A:520:HOH:O[8_554]	2.18	0.02

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	227/239 (95%)	225 (99%)	2 (1%)	0	100	100
1	B	225/239 (94%)	223 (99%)	2 (1%)	0	100	100
All	All	452/478 (95%)	448 (99%)	4 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	201/206 (98%)	199 (99%)	2 (1%)	76	59
1	B	199/206 (97%)	198 (100%)	1 (0%)	88	80
All	All	400/412 (97%)	397 (99%)	3 (1%)	81	68

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	142	ASN
1	A	143	PHE
1	B	143	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	181	GLN
1	B	196	ASN
1	B	197	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
1	CR2	B	65	1	20,20,21	2.12	7 (35%)	25,27,29	1.99	6 (24%)
1	CR2	A	65	1	20,20,21	1.75	6 (30%)	25,27,29	1.98	9 (36%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CR2	B	65	1	-	2/6/25/26	0/2/2/2
1	CR2	A	65	1	-	2/6/25/26	0/2/2/2

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	65	CR2	C1-N2	4.09	1.39	1.32
1	B	65	CR2	C1-N3	4.01	1.43	1.37
1	B	65	CR2	CG2-CB2	3.57	1.53	1.46
1	B	65	CR2	C2-N3	3.35	1.47	1.39
1	A	65	CR2	CA2-N2	3.09	1.45	1.38
1	A	65	CR2	C1-N3	2.75	1.41	1.37
1	A	65	CR2	C1-N2	2.72	1.37	1.32
1	B	65	CR2	CA2-C2	2.70	1.51	1.48
1	A	65	CR2	CG2-CB2	2.69	1.52	1.46
1	B	65	CR2	CA1-C1	2.63	1.52	1.49
1	A	65	CR2	CA2-C2	2.56	1.51	1.48
1	B	65	CR2	CA2-N2	2.45	1.43	1.38
1	A	65	CR2	CB2-CA2	-2.44	1.33	1.35

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	65	CR2	CA2-C2-N3	4.90	105.69	103.37
1	A	65	CR2	CD2-CG2-CD1	4.47	124.26	117.64
1	B	65	CR2	CG2-CB2-CA2	-3.90	125.17	129.94
1	A	65	CR2	C1-CA1-N1	-3.78	104.49	112.85
1	A	65	CR2	O3-C3-CA3	-3.66	115.32	126.39
1	B	65	CR2	C1-CA1-N1	-3.37	105.39	112.85
1	B	65	CR2	O3-C3-CA3	-3.35	116.27	126.39
1	B	65	CR2	C2-CA2-N2	-3.03	106.81	108.93
1	A	65	CR2	CE1-CD1-CG2	-2.99	117.35	121.25
1	B	65	CR2	CD2-CG2-CD1	2.68	121.61	117.64
1	A	65	CR2	CG2-CB2-CA2	-2.64	126.71	129.94
1	A	65	CR2	CE2-CD2-CG2	-2.49	118.00	121.25
1	A	65	CR2	CA2-C2-N3	2.17	104.40	103.37
1	A	65	CR2	CD2-CG2-CB2	-2.16	113.86	121.22
1	A	65	CR2	CE2-CZ-CE1	2.11	123.32	119.77

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	65	CR2	C2-CA2-CB2-CG2
1	B	65	CR2	C2-CA2-CB2-CG2
1	B	65	CR2	N2-CA2-CB2-CG2
1	A	65	CR2	N2-CA2-CB2-CG2

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	65	CR2	1	0
1	A	65	CR2	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	227/239 (94%)	-0.45	3 (1%) 77 78	17, 25, 49, 106	0
1	B	227/239 (94%)	-0.22	5 (2%) 62 61	19, 30, 51, 150	0
All	All	454/478 (94%)	-0.34	8 (1%) 68 69	17, 27, 50, 150	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	2	SER	9.3
1	A	2	SER	8.3
1	B	155	GLN	3.8
1	A	3	LYS	3.6
1	A	154	LYS	3.1
1	B	188	ASP	2.7
1	B	229	HIS	2.6
1	B	3	LYS	2.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	CR2	B	65	19/20	0.96	0.09	18,22,26,27	0
1	CR2	A	65	19/20	0.98	0.08	17,19,24,24	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.