



Full wwPDB X-ray Structure Validation Report ⓘ

May 25, 2020 – 05:13 am BST

PDB ID : 1WKR
Title : Crystal structure of aspartic proteinase from *Irpex lacteus*
Authors : Fujimoto, Z.; Fujii, Y.; Kaneko, S.; Kobayashi, H.; Mizuno, H.
Deposited on : 2004-06-02
Resolution : 1.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1058 (1.30-1.30)
Clashscore	141614	1101 (1.30-1.30)
Ramachandran outliers	138981	1058 (1.30-1.30)
Sidechain outliers	138945	1058 (1.30-1.30)
RSRZ outliers	127900	1029 (1.30-1.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	340	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: -10px; left: 0;">%</div> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 100%; background: linear-gradient(to right, red, orange, yellow, green);"></div> <div style="position: absolute; bottom: -10px; left: 0; width: 100%; text-align: center;"> 87% 11% • </div> </div> </div>
2	I	6	<div> <div style="width: 100%; height: 10px; position: relative;"> <div style="position: absolute; top: 0; left: 0; width: 100%; height: 100%; background: linear-gradient(to right, green, yellow);"></div> <div style="position: absolute; bottom: -10px; left: 0; width: 100%; text-align: center;"> 83% 17% </div> </div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3257 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Polyporopepsin.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	340	Total	C	N	O	0	0	0
			2472	1545	396	531			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	194	SER	THR	SEE REMARK 999	UNP P17576

- Molecule 2 is a protein called pepstatin.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	I	6	Total	C	N	O	0	0	0
			48	34	5	9			

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	693	Total O 693 693	0	0
4	I	9	Total O 9 9	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Polyporopepsin



- Molecule 2: pepstatin



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	37.27Å 78.90Å 54.07Å 90.00° 96.71° 90.00°	Depositor
Resolution (Å)	26.99 – 1.30 44.39 – 1.30	Depositor EDS
% Data completeness (in resolution range)	98.8 (26.99-1.30) 99.0 (44.39-1.30)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.46 (at 1.30Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.147 , 0.171 0.143 , 0.168	Depositor DCC
R_{free} test set	3791 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	7.0	Xtriage
Anisotropy	0.063	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 43.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3257	wwPDB-VP
Average B, all atoms (Å ²)	10.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, IVA, STA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.66	46/2516 (1.8%)	1.27	18/3443 (0.5%)
2	I	0.99	0/17	0.89	0/21
All	All	1.66	46/2533 (1.8%)	1.27	18/3464 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
2	I	0	2
All	All	0	6

All (46) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	285	SER	CB-OG	17.48	1.65	1.42
1	A	167	SER	CB-OG	-11.59	1.27	1.42
1	A	94	ARG	NE-CZ	11.35	1.47	1.33
1	A	285	SER	CA-CB	-11.21	1.36	1.52
1	A	94	ARG	CZ-NH1	10.64	1.46	1.33
1	A	196	ARG	CB-CG	-9.54	1.26	1.52
1	A	167	SER	CA-CB	9.21	1.66	1.52
1	A	199	SER	CB-OG	9.08	1.54	1.42
1	A	229	LYS	CB-CG	-8.75	1.28	1.52
1	A	86	LYS	CE-NZ	8.73	1.70	1.49
1	A	153	GLU	CG-CD	-8.41	1.39	1.51
1	A	196	ARG	CG-CD	8.40	1.73	1.51
1	A	176	TYR	CD2-CE2	7.99	1.51	1.39
1	A	302	GLU	CG-CD	7.78	1.63	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	80	GLY	C-O	7.32	1.35	1.23
1	A	153	GLU	CD-OE2	6.92	1.33	1.25
1	A	122	SER	CA-CB	6.78	1.63	1.52
1	A	302	GLU	CD-OE1	-6.64	1.18	1.25
1	A	58	LYS	CD-CE	6.61	1.67	1.51
1	A	121	THR	CB-OG1	-6.52	1.30	1.43
1	A	67	SER	CB-OG	-6.50	1.33	1.42
1	A	54	ALA	C-N	6.48	1.49	1.34
1	A	183	SER	CB-OG	6.23	1.50	1.42
1	A	72	GLU	CD-OE2	6.20	1.32	1.25
1	A	283	GLY	C-O	-6.18	1.13	1.23
1	A	284	GLY	CA-C	6.17	1.61	1.51
1	A	244	ARG	CD-NE	6.09	1.56	1.46
1	A	69	SER	CB-OG	-6.03	1.34	1.42
1	A	154	SER	CA-CB	5.96	1.61	1.52
1	A	290	TYR	CZ-OH	5.93	1.48	1.37
1	A	176	TYR	CE1-CZ	-5.81	1.30	1.38
1	A	60	SER	CA-CB	5.76	1.61	1.52
1	A	60	SER	CB-OG	-5.74	1.34	1.42
1	A	236	ALA	CA-CB	5.61	1.64	1.52
1	A	85	PRO	C-N	-5.58	1.21	1.34
1	A	5	VAL	CB-CG1	-5.46	1.41	1.52
1	A	290	TYR	CE2-CZ	-5.42	1.31	1.38
1	A	280	THR	CB-OG1	-5.42	1.32	1.43
1	A	263	GLN	CD-OE1	5.41	1.35	1.24
1	A	188	TYR	CG-CD1	5.36	1.46	1.39
1	A	85	PRO	CA-C	5.28	1.63	1.52
1	A	266	GLU	CD-OE1	5.25	1.31	1.25
1	A	238	ASN	C-O	5.19	1.33	1.23
1	A	17	ASN	CG-ND2	-5.08	1.20	1.32
1	A	5	VAL	CB-CG2	-5.05	1.42	1.52
1	A	56	SER	CB-OG	-5.05	1.35	1.42

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	196	ARG	NE-CZ-NH2	17.21	128.91	120.30
1	A	244	ARG	NE-CZ-NH1	10.95	125.78	120.30
1	A	315	ARG	NE-CZ-NH1	9.26	124.93	120.30
1	A	153	GLU	OE1-CD-OE2	-8.36	113.27	123.30
1	A	290	TYR	CD1-CE1-CZ	-7.80	112.78	119.80
1	A	284	GLY	O-C-N	7.37	134.50	122.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	238	ASN	O-C-N	-6.52	112.27	122.70
1	A	302	GLU	OE1-CD-OE2	6.08	130.60	123.30
1	A	290	TYR	CE1-CZ-CE2	5.94	129.30	119.80
1	A	196	ARG	NE-CZ-NH1	-5.89	117.36	120.30
1	A	95	ASP	CB-CG-OD1	5.42	123.18	118.30
1	A	315	ARG	NE-CZ-NH2	-5.37	117.62	120.30
1	A	152	SER	CA-C-O	5.35	131.33	120.10
1	A	196	ARG	NH1-CZ-NH2	-5.15	113.74	119.40
1	A	27	TYR	CZ-CE2-CD2	-5.13	115.19	119.80
1	A	111	ASP	CB-CG-OD1	5.11	122.90	118.30
1	A	94	ARG	CD-NE-CZ	-5.02	116.57	123.60
1	A	237	ASP	CB-CG-OD2	5.00	122.80	118.30

There are no chirality outliers.

All (6) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	154	SER	Mainchain
1	A	238	ASN	Mainchain
1	A	54	ALA	Mainchain
1	A	94	ARG	Sidechain
2	I	804	STA	Mainchain,Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2472	0	2407	10	0
2	I	48	0	60	0	0
3	A	35	0	0	0	0
4	A	693	0	0	5	5
4	I	9	0	0	0	0
All	All	3257	0	2467	10	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:86:LYS:NZ	1:A:86:LYS:CE	1.70	1.53
1:A:285:SER:OG	1:A:285:SER:CB	1.65	1.44
1:A:44:LYS:NZ	4:A:1597:HOH:O	1.67	1.20
1:A:78:THR:HG23	4:A:1595:HOH:O	1.79	0.83
1:A:176:TYR:OH	4:A:1546:HOH:O	2.01	0.76
1:A:293:VAL:HG23	4:A:1513:HOH:O	1.91	0.71
1:A:78:THR:CG2	4:A:1595:HOH:O	2.45	0.59
1:A:86:LYS:NZ	1:A:86:LYS:CD	2.62	0.57
1:A:285:SER:HG	1:A:285:SER:CB	2.08	0.55
1:A:152:SER:HB2	1:A:155:SER:OG	2.12	0.49

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:1431:HOH:O	4:A:1574:HOH:O[1_455]	2.02	0.18
4:A:1033:HOH:O	4:A:1504:HOH:O[1_455]	2.12	0.08
4:A:1082:HOH:O	4:A:1504:HOH:O[1_455]	2.15	0.05
4:A:1445:HOH:O	4:A:1576:HOH:O[2_546]	2.19	0.01
4:A:1189:HOH:O	4:A:1579:HOH:O[1_554]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	338/340 (99%)	332 (98%)	6 (2%)	0	100	100
2	I	3/6 (50%)	2 (67%)	1 (33%)	0	100	100
All	All	341/346 (99%)	334 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	278/278 (100%)	278 (100%)	0	100	100
2	I	2/2 (100%)	2 (100%)	0	100	100
All	All	280/280 (100%)	280 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	17	ASN
1	A	192	ASN
1	A	263	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	STA	I	804	2	10,10,11	1.11	1 (10%)	9,12,14	1.11	1 (11%)
2	STA	I	806	2	8,11,11	1.20	1 (12%)	7,14,14	0.82	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	STA	I	804	2	-	1/11/11/12	-
2	STA	I	806	2	-	2/10/12/12	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	I	806	STA	OH-CH	-2.51	1.38	1.43
2	I	804	STA	CH-CA	-2.46	1.50	1.53

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	I	804	STA	CH-CM-C	-2.55	108.63	113.11

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	I	804	STA	O-C-CM-CH
2	I	806	STA	CA-CB-CG-CD1
2	I	806	STA	CA-CB-CG-CD2

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	A	901	-	4,4,4	0.69	0	6,6,6	0.62	0
3	SO4	A	905	-	4,4,4	3.06	3 (75%)	6,6,6	1.12	0
3	SO4	A	902	-	4,4,4	1.99	1 (25%)	6,6,6	0.59	0
3	SO4	A	906	-	4,4,4	1.80	2 (50%)	6,6,6	1.26	1 (16%)
3	SO4	A	907	-	4,4,4	2.81	2 (50%)	6,6,6	0.93	0
3	SO4	A	904	-	4,4,4	1.62	1 (25%)	6,6,6	0.75	0
3	SO4	A	903	-	4,4,4	0.42	0	6,6,6	0.68	0

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	905	SO4	O1-S	4.14	1.68	1.46
3	A	907	SO4	O2-S	4.00	1.67	1.46
3	A	905	SO4	O2-S	3.94	1.67	1.46
3	A	902	SO4	O1-S	3.75	1.66	1.46
3	A	907	SO4	O1-S	3.63	1.65	1.46
3	A	906	SO4	O3-S	2.72	1.70	1.47
3	A	904	SO4	O1-S	2.42	1.59	1.46
3	A	906	SO4	O4-S	2.20	1.66	1.47
3	A	905	SO4	O4-S	2.15	1.65	1.47

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	906	SO4	O4-S-O3	-2.73	97.39	109.06

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	340/340 (100%)	-0.36	3 (0%) 84 85	3, 6, 13, 23	10 (2%)
2	I	3/6 (50%)	-0.47	0 100 100	5, 5, 6, 6	0
All	All	343/346 (99%)	-0.36	3 (0%) 84 85	3, 6, 13, 23	10 (2%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	99	ASP	2.5
1	A	329	LEU	2.1
1	A	238	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	STA	I	806	12/12	0.91	0.08	7,11,14,14	0
2	STA	I	804	11/12	0.97	0.07	3,4,5,6	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	SO4	A	907	5/5	0.85	0.17	27,28,31,35	0
3	SO4	A	905	5/5	0.88	0.21	32,32,35,36	0
3	SO4	A	901	5/5	0.90	0.20	9,13,15,17	0
3	SO4	A	906	5/5	0.93	0.19	24,33,37,43	0
3	SO4	A	902	5/5	0.95	0.21	15,19,24,27	0
3	SO4	A	904	5/5	0.97	0.21	27,27,29,32	0
3	SO4	A	903	5/5	0.98	0.10	11,12,19,20	0

6.5 Other polymers [i](#)

There are no such residues in this entry.