



Full wwPDB X-ray Structure Validation Report ⓘ

May 17, 2020 – 01:33 am BST

PDB ID : 5WKA
Title : Crystal structure of a GH1 beta-glucosidase retrieved from microbial metagenome of Poraque Amazon lake
Authors : Morais, M.A.B.; Toyama, D.; Ramos, F.C.; Zanthorlin, L.M.; Tonoli, C.C.C.; Miranda, F.P.; Ruller, R.; Henrique-Silva, F.; Murakami, M.T.
Deposited on : 2017-07-24
Resolution : 2.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1235 (2.78-2.74)
Clashscore	141614	1277 (2.78-2.74)
Ramachandran outliers	138981	1257 (2.78-2.74)
Sidechain outliers	138945	1257 (2.78-2.74)
RSRZ outliers	127900	1207 (2.78-2.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	473	<div> <div>2%</div> <div> <div></div> <div>89%</div> <div>9%</div> </div> </div>
1	B	473	<div> <div>4%</div> <div> <div></div> <div>89%</div> <div>8%</div> </div> </div>
1	C	473	<div> <div>2%</div> <div> <div></div> <div>92%</div> <div>7%</div> </div> </div>
1	D	473	<div> <div>3%</div> <div> <div></div> <div>87%</div> <div>12%</div> </div> </div>

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 14816 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-glucosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	467	Total	C	N	O	S	0	0	0
			3699	2353	644	694	8			
1	B	458	Total	C	N	O	S	0	0	0
			3636	2315	632	681	8			
1	C	466	Total	C	N	O	S	0	0	0
			3697	2351	643	695	8			
1	D	468	Total	C	N	O	S	0	0	0
			3710	2359	648	695	8			

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	C	1	Total	C	O	0	0
			6	3	3		
2	C	1	Total	C	O	0	0
			6	3	3		
2	D	1	Total	C	O	0	0
			6	3	3		
2	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	4	3		
3	B	1	Total	C	O	0	0
			7	4	3		
3	C	1	Total	C	O	0	0
			7	4	3		
3	D	1	Total	C	O	0	0
			7	4	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	4	Total	O	0	0
			4	4		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	O	0	0
			1	1		
4	C	3	Total	O	0	0
			3	3		
4	D	2	Total	O	0	0
			2	2		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Beta-glucosidase




• Molecule 1: Beta-glucosidase




• Molecule 1: Beta-glucosidase



• Molecule 1: Beta-glucosidase





4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	165.47Å 56.65Å 209.55Å 90.00° 96.66° 90.00°	Depositor
Resolution (Å)	46.46 – 2.75 48.37 – 2.75	Depositor EDS
% Data completeness (in resolution range)	88.2 (46.46-2.75) 80.1 (48.37-2.75)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.10 (at 2.77Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.232 , 0.270 0.235 , 0.273	Depositor DCC
R_{free} test set	1917 reflections (4.70%)	wwPDB-VP
Wilson B-factor (Å ²)	58.6	Xtriage
Anisotropy	0.296	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 23.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	14816	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.20	0/3808	0.39	0/5201
1	B	0.21	0/3743	0.39	0/5111
1	C	0.22	0/3805	0.40	0/5196
1	D	0.24	0/3819	0.41	0/5215
All	All	0.22	0/15175	0.40	0/20723

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1
1	D	0	1
All	All	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	42	VAL	Peptide
1	D	107	ARG	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3699	0	3494	27	0
1	B	3636	0	3437	20	0
1	C	3697	0	3490	24	0
1	D	3710	0	3507	33	1
2	A	6	0	8	0	0
2	B	6	0	8	0	0
2	C	12	0	16	0	0
2	D	12	0	16	0	0
3	A	7	0	10	0	0
3	B	7	0	10	0	0
3	C	7	0	10	0	0
3	D	7	0	10	0	0
4	A	4	0	0	0	0
4	B	1	0	0	0	0
4	C	3	0	0	0	0
4	D	2	0	0	0	0
All	All	14816	0	14016	102	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (102) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:43:ASN:HD22	1:C:43:ASN:N	1.66	0.91
1:C:42:VAL:HG12	1:C:43:ASN:ND2	2.03	0.72
1:B:105:ARG:NH2	1:B:154:ASP:OD2	2.23	0.71
1:C:164:THR:HG23	1:C:168:LEU:HD12	1.72	0.71
1:D:108:ASN:HD22	1:D:108:ASN:N	1.89	0.69
1:D:303:GLU:HA	1:D:306:ALA:H	1.57	0.68
1:D:304:ASP:OD1	1:D:316:ARG:NH2	2.27	0.68
1:D:303:GLU:HB2	1:D:313:VAL:HG21	1.76	0.67
1:C:43:ASN:N	1:C:43:ASN:ND2	2.38	0.66
1:D:108:ASN:ND2	1:D:108:ASN:N	2.44	0.66
1:D:108:ASN:ND2	1:D:108:ASN:H	1.92	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:95:ASP:O	1:D:99:ARG:HG3	1.98	0.63
1:D:328:ILE:HD13	1:D:330:VAL:HG13	1.80	0.62
1:A:227:GLU:OE1	1:A:302:ARG:NH2	2.25	0.62
1:C:98:GLN:NE2	1:C:151:ASP:OD1	2.32	0.62
1:A:372:ALA:O	1:A:389:ARG:NH1	2.32	0.62
1:C:372:ALA:O	1:C:389:ARG:NH1	2.34	0.60
1:D:298:MET:HB2	1:D:328:ILE:HD11	1.84	0.60
1:D:239:GLY:HA2	1:D:243:ARG:HB2	1.85	0.59
1:B:372:ALA:O	1:B:389:ARG:NH1	2.36	0.59
1:B:239:GLY:HA2	1:B:243:ARG:HB2	1.87	0.57
1:A:164:THR:HG23	1:A:168:LEU:HD12	1.87	0.57
1:D:62:ASP:O	1:D:66:GLU:HG3	2.06	0.56
1:A:28:PRO:HA	1:A:32:ASP:OD2	2.06	0.56
1:D:322:ASP:N	1:D:322:ASP:OD1	2.38	0.56
1:A:230:ASP:OD2	1:A:302:ARG:NH1	2.39	0.56
1:C:42:VAL:HG12	1:C:43:ASN:HD22	1.72	0.55
1:C:151:ASP:OD1	1:C:151:ASP:N	2.32	0.53
1:D:386:ASP:OD2	1:D:389:ARG:NH1	2.42	0.52
1:A:307:ARG:HB3	1:B:307:ARG:HB3	1.91	0.52
1:B:209:HIS:HB2	1:B:210:VAL:HG13	1.92	0.51
1:D:18:GLU:HA	1:D:53:HIS:HB3	1.93	0.51
1:D:107:ARG:O	1:D:109:ILE:HG13	2.11	0.50
1:C:417:SER:OG	1:C:418:LEU:N	2.45	0.50
1:D:142:TYR:O	1:D:146:VAL:HG23	2.11	0.50
1:C:142:TYR:O	1:C:146:VAL:HG23	2.11	0.50
1:A:239:GLY:HA2	1:A:243:ARG:HB2	1.93	0.50
1:D:419:LEU:HB3	1:D:437:ILE:HD11	1.94	0.50
1:D:161:GLU:HG2	1:D:217:ASN:HB3	1.95	0.49
1:D:84:ASP:O	1:D:86:THR:N	2.43	0.49
1:C:239:GLY:HA2	1:C:243:ARG:HB2	1.95	0.49
1:A:142:TYR:O	1:A:146:VAL:HG23	2.14	0.48
1:D:186:ALA:HB3	1:D:268:GLY:HA2	1.94	0.48
1:B:417:SER:OG	1:B:418:LEU:N	2.46	0.48
1:C:42:VAL:C	1:C:43:ASN:HD22	2.18	0.47
1:C:70:GLN:N	1:C:70:GLN:OE1	2.48	0.46
1:A:14:ALA:N	1:A:75:SER:HB3	2.30	0.46
1:B:251:LYS:HA	1:B:251:LYS:HD2	1.65	0.46
1:B:394:ASN:O	1:B:398:GLU:HG3	2.16	0.46
1:B:18:GLU:HA	1:B:53:HIS:HB3	1.96	0.46
1:B:322:ASP:OD1	1:B:323:ASN:N	2.49	0.45
1:B:27:THR:HG23	1:B:83:PRO:HB3	1.98	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:14:ALA:H	1:A:75:SER:HB3	1.80	0.45
1:C:376:ASP:OD2	1:C:389:ARG:NH2	2.50	0.45
1:D:304:ASP:N	1:D:316:ARG:HH22	2.15	0.45
1:C:27:THR:HG23	1:C:83:PRO:HB3	1.98	0.45
1:D:290:ASN:CG	1:D:369:GLU:HB2	2.38	0.45
1:C:18:GLU:HA	1:C:53:HIS:HB3	1.99	0.44
1:D:14:ALA:H	1:D:75:SER:HB3	1.82	0.44
1:A:417:SER:OG	1:A:418:LEU:N	2.50	0.44
1:C:16:GLN:HG2	1:C:422:PHE:O	2.17	0.44
1:C:16:GLN:HB3	1:C:421:ASN:HB2	2.00	0.44
1:B:120:LEU:HD12	1:B:121:PRO:HD2	1.99	0.44
1:A:299:ASN:OD1	1:A:300:VAL:N	2.51	0.44
1:D:164:THR:HG23	1:D:168:LEU:HD12	1.99	0.44
1:A:27:THR:HG23	1:A:83:PRO:HB3	1.99	0.43
1:A:416:TRP:HA	1:A:417:SER:HA	1.58	0.43
1:A:340:MET:HG2	1:A:373:ALA:HB3	2.01	0.43
1:B:217:ASN:HA	1:B:290:ASN:HB2	2.00	0.43
1:D:328:ILE:CD1	1:D:330:VAL:HG13	2.49	0.43
1:A:16:GLN:NE2	1:A:421:ASN:OD1	2.52	0.43
1:C:416:TRP:HA	1:C:417:SER:HA	1.59	0.43
1:D:290:ASN:OD1	1:D:368:THR:OG1	2.30	0.43
1:B:142:TYR:O	1:B:146:VAL:HG23	2.19	0.43
1:B:335:ARG:HA	1:B:336:PRO:HD3	1.89	0.43
1:C:290:ASN:CG	1:C:369:GLU:HB2	2.39	0.42
1:D:238:ASP:OD1	1:D:242:ASN:ND2	2.49	0.42
1:D:391:THR:HG23	1:D:465:LEU:HD21	2.02	0.42
1:A:118:TRP:N	1:A:118:TRP:CD1	2.88	0.42
1:A:416:TRP:CD2	1:A:417:SER:HB2	2.53	0.41
1:A:70:GLN:OE1	1:A:70:GLN:N	2.53	0.41
1:C:155:LYS:NZ	1:C:366:TYR:OH	2.53	0.41
1:B:263:SER:HB2	1:D:225:THR:HA	2.01	0.41
1:A:170:TYR:CD2	1:A:179:ARG:HD2	2.55	0.41
1:A:207:ARG:O	1:A:210:VAL:HG22	2.20	0.41
1:A:290:ASN:CG	1:A:369:GLU:HB2	2.41	0.41
1:B:52:ASP:O	1:B:56:ARG:HG3	2.20	0.41
1:A:190:HIS:CE1	1:A:272:VAL:HG22	2.55	0.41
1:A:386:ASP:OD1	1:A:386:ASP:N	2.54	0.41
1:A:366:TYR:CD2	1:A:412:GLY:HA3	2.55	0.41
1:D:230:ASP:OD2	1:D:302:ARG:NH1	2.53	0.41
1:B:416:TRP:HA	1:B:417:SER:HA	1.56	0.41
1:D:160:ASN:HA	1:D:215:THR:HB	2.03	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:386:ASP:OD1	1:C:386:ASP:N	2.53	0.41
1:C:443:ARG:CZ	1:C:445:ILE:HD11	2.51	0.41
1:D:367:ILE:HG12	1:D:410:LEU:HD11	2.02	0.41
1:B:204:ARG:NH2	1:B:285:ASP:OD2	2.54	0.41
1:A:29:SER:HB3	1:A:79:PRO:HG2	2.03	0.40
1:C:228:ASP:O	1:C:232:GLU:HG3	2.21	0.40
1:D:335:ARG:HA	1:D:336:PRO:HD3	1.88	0.40
1:A:335:ARG:HA	1:A:336:PRO:HD3	1.87	0.40
1:B:96:PHE:HD1	1:B:99:ARG:NH2	2.20	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:141:ASP:OD2	1:D:264:ARG:NH1[4_649]	2.16	0.04

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	463/473 (98%)	445 (96%)	18 (4%)	0	100	100
1	B	454/473 (96%)	428 (94%)	26 (6%)	0	100	100
1	C	462/473 (98%)	439 (95%)	22 (5%)	1 (0%)	47	69
1	D	464/473 (98%)	438 (94%)	24 (5%)	2 (0%)	34	53
All	All	1843/1892 (97%)	1750 (95%)	90 (5%)	3 (0%)	47	69

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	43	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	108	ASN
1	D	85	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	376/382 (98%)	374 (100%)	2 (0%)	88	92
1	B	369/382 (97%)	368 (100%)	1 (0%)	92	95
1	C	376/382 (98%)	375 (100%)	1 (0%)	92	95
1	D	377/382 (99%)	374 (99%)	3 (1%)	81	88
All	All	1498/1528 (98%)	1491 (100%)	7 (0%)	88	92

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	ASP
1	A	302	ARG
1	B	304	ASP
1	C	43	ASN
1	D	108	ASN
1	D	257	ASP
1	D	350	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	43	ASN
1	D	108	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	D	502	-	5,5,5	0.37	0	5,5,5	0.26	0
3	PEG	B	502	-	6,6,6	0.62	0	5,5,5	0.29	0
3	PEG	A	502	-	6,6,6	0.60	0	5,5,5	0.29	0
3	PEG	D	503	-	6,6,6	0.61	0	5,5,5	0.29	0
2	GOL	C	502	-	5,5,5	0.36	0	5,5,5	0.24	0
2	GOL	C	501	-	5,5,5	0.37	0	5,5,5	0.27	0
2	GOL	A	501	-	5,5,5	0.36	0	5,5,5	0.24	0
2	GOL	B	501	-	5,5,5	0.36	0	5,5,5	0.31	0
3	PEG	C	503	-	6,6,6	0.61	0	5,5,5	0.26	0
2	GOL	D	501	-	5,5,5	0.38	0	5,5,5	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	D	502	-	-	2/4/4/4	-
3	PEG	B	502	-	-	3/4/4/4	-
3	PEG	A	502	-	-	0/4/4/4	-
3	PEG	D	503	-	-	2/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	C	502	-	-	2/4/4/4	-
2	GOL	C	501	-	-	2/4/4/4	-
2	GOL	A	501	-	-	0/4/4/4	-
2	GOL	B	501	-	-	2/4/4/4	-
3	PEG	C	503	-	-	0/4/4/4	-
2	GOL	D	501	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	502	GOL	O1-C1-C2-C3
2	B	501	GOL	O1-C1-C2-O2
2	C	501	GOL	O1-C1-C2-C3
2	B	501	GOL	O1-C1-C2-C3
2	C	502	GOL	O1-C1-C2-O2
3	B	502	PEG	O2-C3-C4-O4
3	D	503	PEG	O1-C1-C2-O2
2	D	502	GOL	O1-C1-C2-O2
3	D	503	PEG	O2-C3-C4-O4
2	C	501	GOL	O1-C1-C2-O2
3	B	502	PEG	O1-C1-C2-O2
2	D	502	GOL	O1-C1-C2-C3
3	B	502	PEG	C4-C3-O2-C2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	467/473 (98%)	0.42	11 (2%) 59 68	45, 56, 65, 85	3 (0%)
1	B	458/473 (96%)	0.53	20 (4%) 34 41	48, 59, 70, 86	5 (1%)
1	C	466/473 (98%)	0.31	11 (2%) 59 68	45, 57, 69, 83	4 (0%)
1	D	468/473 (98%)	0.30	12 (2%) 56 65	40, 53, 63, 74	13 (2%)
All	All	1859/1892 (98%)	0.39	54 (2%) 51 61	40, 56, 68, 86	25 (1%)

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	316	ARG	6.3
1	B	226	THR	4.5
1	B	323	ASN	3.6
1	D	300	VAL	3.5
1	B	254	TYR	3.1
1	C	473	ASN	3.0
1	A	470	ASP	2.9
1	A	226	THR	2.9
1	B	464	ASP	2.9
1	A	209	HIS	2.8
1	D	329	GLY	2.8
1	B	259	ILE	2.7
1	B	465	LEU	2.7
1	D	22	ALA	2.7
1	B	405	ASP	2.6
1	D	301	GLY	2.6
1	D	86	THR	2.6
1	A	205	ALA	2.6
1	C	14	ALA	2.6
1	C	84	ASP	2.5
1	C	54	TYR	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	313	VAL	2.5
1	B	251	LYS	2.4
1	A	315	PRO	2.4
1	B	459	THR	2.4
1	C	308	ALA	2.3
1	A	171	GLY	2.3
1	B	225	THR	2.3
1	B	232	GLU	2.3
1	A	203	ALA	2.3
1	B	269	PHE	2.3
1	C	142	TYR	2.3
1	A	462	LEU	2.3
1	B	328	ILE	2.3
1	A	137	LEU	2.2
1	D	180	THR	2.2
1	D	226	THR	2.1
1	B	84	ASP	2.1
1	B	43	ASN	2.1
1	A	273	GLN	2.1
1	B	265	TRP	2.1
1	C	120	LEU	2.1
1	C	256	ALA	2.1
1	C	470	ASP	2.1
1	B	209	HIS	2.1
1	B	410	LEU	2.1
1	D	112	LEU	2.0
1	D	91	GLN	2.0
1	A	301	GLY	2.0
1	D	328	ILE	2.0
1	B	262	TYR	2.0
1	C	242	ASN	2.0
1	D	373	ALA	2.0
1	C	296	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	PEG	B	502	7/7	0.66	0.30	88,89,94,96	0
2	GOL	B	501	6/6	0.81	0.38	83,83,83,83	0
2	GOL	C	502	6/6	0.85	0.29	57,57,57,58	0
2	GOL	A	501	6/6	0.88	0.34	66,66,66,67	0
2	GOL	C	501	6/6	0.88	0.25	74,74,75,76	0
2	GOL	D	501	6/6	0.88	0.22	61,61,61,61	0
3	PEG	D	503	7/7	0.91	0.19	67,67,68,68	0
3	PEG	A	502	7/7	0.91	0.38	78,78,79,79	0
3	PEG	C	503	7/7	0.92	0.23	65,65,65,66	0
2	GOL	D	502	6/6	0.92	0.15	64,65,66,74	0

6.5 Other polymers [i](#)

There are no such residues in this entry.