



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 2, 2022 – 10:31 AM JST

PDB ID : 7WKY
Title : Crystal Structure of the first bromodomain of human BRD4 in complex with the inhibitor Y13153
Authors : Li, J.; Zhang, C.; Xu, H.; Zhuang, X.; Wu, X.; Zhang, Y.; Xu, Y.
Deposited on : 2022-01-12
Resolution : 2.83 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at <http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.29
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.29


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.83 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1031 (2.86-2.82)
Clashscore	141614	1078 (2.86-2.82)
Ramachandran outliers	138981	1050 (2.86-2.82)
Sidechain outliers	138945	1051 (2.86-2.82)
RSRZ outliers	127900	1019 (2.86-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	141	<div> <div>2%</div> <div> <div></div> <div>74%</div> <div>•</div> <div>23%</div> </div> </div>
1	B	141	<div> <div>%</div> <div> <div></div> <div>74%</div> <div>•</div> <div>23%</div> </div> </div>
1	C	141	<div> <div>%</div> <div> <div></div> <div>73%</div> <div>• •</div> <div>22%</div> </div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2961 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bromodomain-containing protein 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	109	Total	C	N	O	S	0	0	0
			923	603	151	163	6			
1	B	109	Total	C	N	O	S	0	1	0
			931	607	152	166	6			
1	C	110	Total	C	N	O	S	0	0	0
			932	609	153	164	6			

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	28	MET	-	expression tag	UNP O60885
A	29	LYS	-	expression tag	UNP O60885
A	30	LYS	-	expression tag	UNP O60885
A	31	GLY	-	expression tag	UNP O60885
A	32	HIS	-	expression tag	UNP O60885
A	33	HIS	-	expression tag	UNP O60885
A	34	HIS	-	expression tag	UNP O60885
A	35	HIS	-	expression tag	UNP O60885
A	36	HIS	-	expression tag	UNP O60885
A	37	HIS	-	expression tag	UNP O60885
A	38	LEU	-	expression tag	UNP O60885
A	39	VAL	-	expression tag	UNP O60885
A	40	PRO	-	expression tag	UNP O60885
A	41	ARG	-	expression tag	UNP O60885
A	42	GLY	-	expression tag	UNP O60885
A	43	SER	-	expression tag	UNP O60885
B	28	MET	-	expression tag	UNP O60885
B	29	LYS	-	expression tag	UNP O60885
B	30	LYS	-	expression tag	UNP O60885
B	31	GLY	-	expression tag	UNP O60885
B	32	HIS	-	expression tag	UNP O60885
B	33	HIS	-	expression tag	UNP O60885
B	34	HIS	-	expression tag	UNP O60885

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	35	HIS	-	expression tag	UNP O60885
B	36	HIS	-	expression tag	UNP O60885
B	37	HIS	-	expression tag	UNP O60885
B	38	LEU	-	expression tag	UNP O60885
B	39	VAL	-	expression tag	UNP O60885
B	40	PRO	-	expression tag	UNP O60885
B	41	ARG	-	expression tag	UNP O60885
B	42	GLY	-	expression tag	UNP O60885
B	43	SER	-	expression tag	UNP O60885
C	28	MET	-	expression tag	UNP O60885
C	29	LYS	-	expression tag	UNP O60885
C	30	LYS	-	expression tag	UNP O60885
C	31	GLY	-	expression tag	UNP O60885
C	32	HIS	-	expression tag	UNP O60885
C	33	HIS	-	expression tag	UNP O60885
C	34	HIS	-	expression tag	UNP O60885
C	35	HIS	-	expression tag	UNP O60885
C	36	HIS	-	expression tag	UNP O60885
C	37	HIS	-	expression tag	UNP O60885
C	38	LEU	-	expression tag	UNP O60885
C	39	VAL	-	expression tag	UNP O60885
C	40	PRO	-	expression tag	UNP O60885
C	41	ARG	-	expression tag	UNP O60885
C	42	GLY	-	expression tag	UNP O60885
C	43	SER	-	expression tag	UNP O60885

- Molecule 2 is 2-(2-cyclopentyl-1 {H}-imidazol-5-yl)-7-[2-(4-fluoranyl-2,6-dimethyl-phenoxy)-5-(2-oxidanylpropan-2-yl)phenyl]-5-methyl-furo[3,2-c]pyridin-4-one (three-letter code: JFR) (formula: C₃₃H₃₄FN₃O₄) (labeled as "Ligand of Interest" by depositor).



- GOL
-
- The diagram shows the skeletal structure of 1,2,3-propanetriol (glycerol). The carbon backbone is represented by three vertices labeled C1, C2, and C3 in green. C1 is on the left, C2 is in the middle, and C3 is on the right and slightly higher. Each carbon is bonded to a hydroxyl group (OH) in red. The OH group on C1 is labeled O1 in green below it. The OH group on C2 is labeled O2 in green below it. The OH group on C3 is labeled O3 in green to its right. The bonds between the carbons and between the carbons and the hydroxyl groups are shown as thin grey lines.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		




Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	9	Total	O	0	0
			9	9		
4	B	4	Total	O	0	0
			4	4		
4	C	3	Total	O	0	0
			3	3		


- Molecule 1: Bromodomain-containing protein 4



- Chain B: 74% 23% 3%



- Chain C:  73% 2% 22%



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	140.85Å 140.85Å 131.83Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	121.98 – 2.83 70.42 – 2.83	Depositor EDS
% Data completeness (in resolution range)	98.4 (121.98-2.83) 98.4 (70.42-2.83)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.27 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.178 , 0.246 0.186 , 0.239	Depositor DCC
R_{free} test set	909 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å ²)	62.8	Xtriage
Anisotropy	0.859	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 48.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2961	wwPDB-VP
Average B, all atoms (Å ²)	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, JFR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.58	0/947	0.72	0/1285
1	B	0.52	0/955	0.69	0/1296
1	C	0.52	0/956	0.69	0/1296
All	All	0.54	0/2858	0.70	0/3877

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	923	0	926	0	0
1	B	931	0	929	0	0
1	C	932	0	939	1	0
2	A	41	0	0	0	0
2	B	41	0	0	0	0
2	C	41	0	0	0	0
3	A	18	0	24	0	0
3	B	12	0	16	0	0
3	C	6	0	8	0	0
4	A	9	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	4	0	0	0	0
4	C	3	0	0	0	0
All	All	2961	0	2842	1	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All (1) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:113:ARG:NH1	1:C:128:ASP:OD2	2.50	0.44

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	107/141 (76%)	105 (98%)	2 (2%)	0	100	100
1	B	108/141 (77%)	105 (97%)	2 (2%)	1 (1%)	17	34
1	C	108/141 (77%)	104 (96%)	4 (4%)	0	100	100
All	All	323/423 (76%)	314 (97%)	8 (2%)	1 (0%)	41	61

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	61	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	102/132 (77%)	97 (95%)	5 (5%)	25	47
1	B	103/132 (78%)	100 (97%)	3 (3%)	42	67
1	C	103/132 (78%)	97 (94%)	6 (6%)	20	38
All	All	308/396 (78%)	294 (96%)	14 (4%)	27	51

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	58	ARG
1	A	63	LEU
1	A	72	LYS
1	A	78	GLN
1	A	141	LYS
1	B	141	LYS
1	B	155	LYS
1	B	166	THR
1	C	57	LYS
1	C	60	THR
1	C	72	LYS
1	C	91	LYS
1	C	94	LEU
1	C	128	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	64	GLN
1	A	78	GLN
1	A	116	ASN
1	A	162	ASN
1	B	61	ASN
1	C	127	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	JFR	C	201	-	37,46,46	3.37	12 (32%)	51,70,70	1.20	6 (11%)
2	JFR	A	201	-	37,46,46	3.12	9 (24%)	51,70,70	1.40	9 (17%)
3	GOL	B	203	-	5,5,5	0.39	0	5,5,5	0.72	0
3	GOL	C	202	-	5,5,5	0.23	0	5,5,5	0.25	0
3	GOL	A	204	-	5,5,5	0.71	0	5,5,5	0.98	0
3	GOL	B	202	-	5,5,5	0.43	0	5,5,5	0.48	0
2	JFR	B	201	-	37,46,46	3.47	12 (32%)	51,70,70	1.32	10 (19%)
3	GOL	A	202	-	5,5,5	0.30	0	5,5,5	0.84	0
3	GOL	A	203	-	5,5,5	0.27	0	5,5,5	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	JFR	C	201	-	-	0/14/29/29	0/6/6/6

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	JFR	A	201	-	-	0/14/29/29	0/6/6/6
3	GOL	B	203	-	-	2/4/4/4	-
3	GOL	C	202	-	-	2/4/4/4	-
3	GOL	A	204	-	-	2/4/4/4	-
3	GOL	B	202	-	-	2/4/4/4	-
2	JFR	B	201	-	-	0/14/29/29	0/6/6/6
3	GOL	A	202	-	-	1/4/4/4	-
3	GOL	A	203	-	-	2/4/4/4	-

All (33) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	201	JFR	CAQ-CAM	-14.79	1.43	1.53
2	C	201	JFR	CAQ-CAM	-13.39	1.44	1.53
2	A	201	JFR	CAQ-CAM	-11.95	1.45	1.53
2	B	201	JFR	CAK-CAH	-6.46	1.34	1.49
2	C	201	JFR	CAK-CAH	-6.30	1.35	1.49
2	A	201	JFR	CAK-CAH	-5.78	1.36	1.49
2	B	201	JFR	CAB-CAC	-5.78	1.35	1.47
2	C	201	JFR	CAB-CAC	-5.63	1.36	1.47
2	A	201	JFR	CAA-CAF	-5.60	1.34	1.47
2	A	201	JFR	CBA-CAX	-5.58	1.39	1.51
2	B	201	JFR	CBA-CAX	-5.46	1.40	1.51
2	A	201	JFR	CAB-CAC	-5.38	1.36	1.47
2	C	201	JFR	CAZ-CAT	-5.35	1.40	1.51
2	C	201	JFR	FAY-CAV	-5.34	1.23	1.36
2	C	201	JFR	CBA-CAX	-5.34	1.40	1.51
2	A	201	JFR	CAZ-CAT	-5.32	1.40	1.51
2	B	201	JFR	CAA-CAF	-5.31	1.34	1.47
2	C	201	JFR	CAA-CAF	-5.05	1.35	1.47
2	B	201	JFR	CAZ-CAT	-5.05	1.40	1.51
2	B	201	JFR	CAJ-CAF	-4.00	1.42	1.49
2	A	201	JFR	CAJ-CAF	-3.90	1.42	1.49
2	C	201	JFR	CAJ-CAF	-3.74	1.42	1.49
2	B	201	JFR	CAI-CAB	-3.24	1.33	1.40
2	A	201	JFR	CAI-CAB	-2.92	1.34	1.40
2	B	201	JFR	FAY-CAV	2.85	1.43	1.36
2	C	201	JFR	CAE-NAD	2.60	1.43	1.36
2	C	201	JFR	CAI-CAB	-2.60	1.34	1.40
2	A	201	JFR	OBN-CAQ	2.54	1.48	1.44
2	B	201	JFR	CBM-CAQ	2.48	1.55	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	201	JFR	CAE-NAD	2.31	1.42	1.36
2	B	201	JFR	CBB-CAK	-2.23	1.34	1.37
2	C	201	JFR	CBB-CAK	-2.05	1.34	1.37
2	C	201	JFR	OBN-CAQ	2.03	1.47	1.44

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	201	JFR	OBN-CAQ-CAM	3.57	114.97	109.13
2	A	201	JFR	CBM-CAQ-CAM	-3.22	105.15	110.91
2	C	201	JFR	CBI-CBJ-CBF	-3.10	99.10	104.71
2	B	201	JFR	CAP-OAR-CAS	3.02	124.50	117.79
2	C	201	JFR	CAP-OAR-CAS	2.98	124.40	117.79
2	B	201	JFR	CAT-CAU-CAV	2.97	121.41	119.26
2	B	201	JFR	CAW-CAV-CAU	-2.88	119.89	123.52
2	A	201	JFR	CAL-CAM-CAQ	2.61	123.70	120.43
2	B	201	JFR	CBI-CBH-CBG	-2.58	97.32	105.99
2	A	201	JFR	CAP-CAJ-CAF	2.57	124.71	121.57
2	B	201	JFR	FAY-CAV-CAW	2.48	121.80	118.25
2	B	201	JFR	CAL-CAM-CAQ	2.44	123.48	120.43
2	C	201	JFR	CAP-CAJ-CAF	2.38	124.48	121.57
2	A	201	JFR	CAW-CAV-CAU	-2.28	120.64	123.52
2	C	201	JFR	OBN-CAQ-CAM	2.27	112.85	109.13
2	A	201	JFR	OBN-CAQ-CBO	-2.27	102.51	107.08
2	C	201	JFR	CAF-CAE-NAD	-2.27	120.06	122.96
2	A	201	JFR	CBF-CBD-NBC	2.23	129.33	125.08
2	A	201	JFR	CBF-CBD-NBE	-2.18	120.93	125.08
2	A	201	JFR	CBI-CBH-CBG	-2.16	98.71	105.99
2	B	201	JFR	CAP-CAJ-CAF	2.16	124.20	121.57
2	B	201	JFR	CAN-CAM-CAQ	-2.14	118.82	121.55
2	C	201	JFR	CAW-CAV-CAU	-2.11	120.85	123.52
2	B	201	JFR	CAF-CAE-NAD	-2.11	120.26	122.96
2	B	201	JFR	OBN-CAQ-CAM	2.10	112.57	109.13

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	203	GOL	O1-C1-C2-C3
3	A	202	GOL	C1-C2-C3-O3
3	A	203	GOL	C1-C2-C3-O3
3	A	204	GOL	C1-C2-C3-O3

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	B	202	GOL	C1-C2-C3-O3
3	C	202	GOL	C1-C2-C3-O3
3	A	203	GOL	O2-C2-C3-O3
3	C	202	GOL	O2-C2-C3-O3
3	A	204	GOL	O2-C2-C3-O3
3	B	203	GOL	O1-C1-C2-O2
3	B	202	GOL	O2-C2-C3-O3

There are no ring outliers.


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand JFR C 201



Ligand JFR A 201





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	109/141 (77%)	0.09	3 (2%) 53 47	46, 62, 101, 139	0
1	B	109/141 (77%)	0.10	2 (1%) 68 63	50, 65, 99, 122	0
1	C	110/141 (78%)	-0.02	1 (0%) 84 83	55, 76, 110, 129	0
All	All	328/423 (77%)	0.06	6 (1%) 68 63	46, 69, 107, 139	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	58	ARG	4.6
1	A	59	GLN	3.9
1	B	59	GLN	3.7
1	A	58	ARG	3.3
1	A	60	THR	3.2
1	C	166	THR	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	GOL	B	203	6/6	0.78	0.33	86,93,98,101	0
3	GOL	C	202	6/6	0.85	0.45	102,110,111,116	0
3	GOL	B	202	6/6	0.86	0.31	85,96,100,102	0
3	GOL	A	204	6/6	0.88	0.37	62,67,71,77	0
3	GOL	A	203	6/6	0.90	0.35	85,90,97,102	0
3	GOL	A	202	6/6	0.96	0.30	60,67,70,70	0
2	JFR	A	201	41/41	0.96	0.20	47,67,86,90	0
2	JFR	C	201	41/41	0.96	0.22	66,79,110,114	0
2	JFR	B	201	41/41	0.97	0.23	55,76,108,110	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around JFR A 201:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around JFR C 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around JFR B 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.