



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 8, 2020 – 09:12 PM BST

PDB ID : 2WLZ  
Title : Chitinase A from *Serratia marcescens* ATCC990 in complex with Chitobiothiazoline.  
Authors : Taylor, E.J.; Dennis, R.J.; Macdonald, J.M.; Tarling, C.A.; Knapp, S.; Withers, S.G.; Davies, G.J.  
Deposited on : 2009-06-29  
Resolution : 1.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.13.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	7484 (1.84-1.80)
Clashscore	141614	8401 (1.84-1.80)
Ramachandran outliers	138981	8290 (1.84-1.80)
Sidechain outliers	138945	8290 (1.84-1.80)
RSRZ outliers	127900	7371 (1.84-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	548	

## 2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 4663 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CHITINASE A.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	538	4148	2636	699	799	14	0	5	0

- Molecule 2 is 1,4-DIETHYLENE DIOXIDE (three-letter code: DIO) (formula: C<sub>4</sub>H<sub>8</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	6	4	2	0	0
2	A	1	6	4	2	0	0
2	A	1	6	4	2	0	0
2	A	1	6	4	2	0	0
2	A	1	6	4	2	0	0

- Molecule 3 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula:  $C_4H_{10}O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	4	3		
3	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 4 is 3AR,5R,6S,7R,7AR-5-HYDROXYMETHYL-2-METHYL-5,6,7,7A-TETRAHYDRO-3AH-PYRANO[3,2-D]THIAZOLE-6,7-DIOL (three-letter code: NGT) (formula:  $C_8H_{13}NO_4S$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			14	8	1	4	1		

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	443	Total	O	0	0
			443	443		

**i**

- Molecule 1: CHITINASE A

Protein	Accession	Score	Rank
Protein A	P00923	1.00	1
Protein B	P00923	0.99	2
Protein C	P00923	0.98	3
Protein D	P00923	0.97	4
Protein E	P00923	0.96	5
Protein F	P00923	0.95	6
Protein G	P00923	0.94	7
Protein H	P00923	0.93	8
Protein I	P00923	0.92	9
Protein J	P00923	0.91	10
Protein K	P00923	0.90	11
Protein L	P00923	0.89	12
Protein M	P00923	0.88	13
Protein N	P00923	0.87	14
Protein O	P00923	0.86	15
Protein P	P00923	0.85	16
Protein Q	P00923	0.84	17
Protein R	P00923	0.83	18
Protein S	P00923	0.82	19
Protein T	P00923	0.81	20
Protein U	P00923	0.80	21
Protein V	P00923	0.79	22
Protein W	P00923	0.78	23
Protein X	P00923	0.77	24
Protein Y	P00923	0.76	25
Protein Z	P00923	0.75	26
Protein AA	P00923	0.74	27
Protein AB	P00923	0.73	28
Protein AC	P00923	0.72	29
Protein AD	P00923	0.71	30
Protein AE	P00923	0.70	31
Protein AF	P00923	0.69	32
Protein AG	P00923	0.68	33
Protein AH	P00923	0.67	34
Protein AI	P00923	0.66	35
Protein AJ	P00923	0.65	36
Protein AK	P00923	0.64	37
Protein AL	P00923	0.63	38
Protein AM	P00923	0.62	39
Protein AN	P00923	0.61	40
Protein AO	P00923	0.60	41
Protein AP	P00923	0.59	42
Protein AQ	P00923	0.58	43
Protein AR	P00923	0.57	44
Protein AS	P00923	0.56	45
Protein AT	P00923	0.55	46
Protein AU	P00923	0.54	47
Protein AV	P00923	0.53	48
Protein AW	P00923	0.52	49
Protein AX	P00923	0.51	50
Protein AY	P00923	0.50	51
Protein AZ	P00923	0.49	52
Protein BA	P00923	0.48	53
Protein BB	P00923	0.47	54
Protein BC	P00923	0.46	55
Protein BD	P00923	0.45	56
Protein BE	P00923	0.44	57
Protein BF	P00923	0.43	58
Protein BG	P00923	0.42	59
Protein BH	P00923	0.41	60
Protein BI	P00923	0.40	61
Protein BJ	P00923	0.39	62
Protein BK	P00923	0.38	63
Protein BL	P00923	0.37	64
Protein BM	P00923	0.36	65
Protein BN	P00923	0.35	66
Protein BO	P00923	0.34	67
Protein BP	P00923	0.33	68
Protein BQ	P00923	0.32	69
Protein BR	P00923	0.31	70
Protein BS	P00923	0.30	71
Protein BT	P00923	0.29	72
Protein BU	P00923	0.28	73
Protein BV	P00923	0.27	74
Protein BW	P00923	0.26	75
Protein BX	P00923	0.25	76
Protein BY	P00923	0.24	77
Protein BZ	P00923	0.23	78
Protein CA	P00923	0.22	79
Protein CB	P00923	0.21	80
Protein CC	P00923	0.20	81
Protein CD	P00923	0.19	82
Protein CE	P00923	0.18	83
Protein CF	P00923	0.17	84
Protein CG	P00923	0.16	85
Protein CH	P00923	0.15	86
Protein CI	P00923	0.14	87
Protein CJ	P00923	0.13	88
Protein CK	P00923	0.12	89
Protein CL	P00923	0.11	90

## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	130.64Å 201.27Å 59.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	109.76 – 1.82 43.90 – 1.82	Depositor EDS
% Data completeness (in resolution range)	98.3 (109.76-1.82) 98.3 (43.90-1.82)	Depositor EDS
$R_{merge}$	0.17	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.19 (at 1.83Å)	Xtriage
Refinement program	REFMAC 5.5.0097	Depositor
R, $R_{free}$	0.184 , 0.218 0.185 , 0.217	Depositor DCC
$R_{free}$ test set	3456 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.6	Xtriage
Anisotropy	0.099	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 49.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4663	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.12% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NGT, DIO, PEG, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.92	1/4265 (0.0%)	0.82	3/5788 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	213	ALA	CA-CB	5.07	1.63	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	132	ASP	CB-CG-OD1	5.94	123.65	118.30
1	A	279	ASP	CB-CG-OD1	5.60	123.34	118.30
1	A	340	LYS	CD-CE-NZ	5.53	124.42	111.70

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	560	ALA	Peptide



## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4148	0	4014	27	0
2	A	30	0	40	5	0
3	A	14	0	20	2	0
4	A	14	0	12	2	0
5	A	14	0	13	3	0
6	A	443	0	0	6	0
All	All	4663	0	4099	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:476:ILE:HD13	5:A:1570:NAG:H83	1.51	0.92
1:A:148:LYS:HE2	6:A:2429:HOH:O	1.68	0.92
1:A:54:VAL:HB	2:A:1562:DIO:H1'1	1.76	0.66
1:A:169:VAL:HG11	1:A:252:ASN:HD21	1.63	0.64
1:A:207:ILE:HD13	2:A:1564:DIO:H2'1	1.81	0.62
3:A:1568:PEG:H12	6:A:2318:HOH:O	1.99	0.61
1:A:313[B]:ASP:OD1	4:A:1569:NGT:HC83	2.04	0.58
2:A:1564:DIO:H2'2	5:A:1570:NAG:H62	1.90	0.53
1:A:207:ILE:HD11	1:A:276:THR:HG21	1.91	0.52
3:A:1568:PEG:C1	6:A:2318:HOH:O	2.61	0.49
1:A:343:ARG:HD2	1:A:381:SER:O	2.12	0.49
1:A:493:TYR:OH	1:A:517:ASP:OD2	2.24	0.49
1:A:340:LYS:HE2	6:A:2297:HOH:O	2.13	0.48
4:A:1569:NGT:HC61	5:A:1570:NAG:C7	2.45	0.47
1:A:299:GLU:CD	2:A:1562:DIO:H21	2.35	0.46
1:A:129:VAL:HG13	1:A:137:HIS:HB2	1.99	0.45
1:A:173:ASN:ND2	6:A:2176:HOH:O	2.49	0.45
1:A:193:PRO:HD2	1:A:227:SER:O	2.17	0.45
1:A:131:ALA:HA	1:A:137:HIS:CG	2.53	0.43
1:A:299:GLU:OE2	2:A:1562:DIO:H21	2.19	0.43
1:A:319:GLY:O	1:A:320:LYS:HB2	2.18	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:492:GLN:HG3	6:A:2395:HOH:O	2.19	0.42
1:A:42:GLU:HG3	1:A:57:LYS:HD2	2.01	0.42
1:A:270:PRO:HD3	1:A:307:PHE:CD1	2.55	0.41
1:A:190:GLY:HA2	1:A:191:PHE:HA	1.84	0.41
1:A:434:PRO:O	1:A:533:LEU:HA	2.19	0.41
1:A:311:ASP:OD2	1:A:313[A]:ASP:OD1	2.38	0.41
1:A:207:ILE:CD1	1:A:276:THR:HG21	2.49	0.41
1:A:313[B]:ASP:OD2	1:A:315:GLU:OE2	2.38	0.41
1:A:521:VAL:HG21	1:A:548:ILE:HG23	2.03	0.40
1:A:528:VAL:HG13	1:A:533:LEU:HB2	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	541/548 (99%)	528 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	434/439 (99%)	425 (98%)	9 (2%)	53	41

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	57	LYS
1	A	75[A]	THR
1	A	75[B]	THR
1	A	188	LEU
1	A	328	SER
1	A	399	LYS
1	A	455	GLN
1	A	492	GLN
1	A	514	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	173	ASN
1	A	215	GLN
1	A	239	GLN
1	A	252	ASN
1	A	481	GLN
1	A	485	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	NAG	A	1570	4	14,14,15	0.94	0	17,19,21	1.73	4 (23%)
2	DIO	A	1565	-	6,6,6	0.58	0	6,6,6	0.72	0
4	NGT	A	1569	5	13,15,15	3.27	2 (15%)	12,22,22	1.91	3 (25%)
3	PEG	A	1567	-	6,6,6	0.73	0	5,5,5	0.68	0
2	DIO	A	1566	-	6,6,6	0.62	0	6,6,6	1.25	2 (33%)
3	PEG	A	1568	-	6,6,6	0.90	0	5,5,5	0.97	0
2	DIO	A	1562	-	6,6,6	0.61	0	6,6,6	0.56	0
2	DIO	A	1563	-	6,6,6	0.53	0	6,6,6	1.00	0
2	DIO	A	1564	-	6,6,6	0.70	0	6,6,6	0.83	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	A	1570	4	-	0/6/23/26	0/1/1/1
2	DIO	A	1565	-	-	-	0/1/1/1
4	NGT	A	1569	5	-	0/2/30/30	0/2/2/2
3	PEG	A	1567	-	-	2/4/4/4	-
2	DIO	A	1566	-	-	-	0/1/1/1
3	PEG	A	1568	-	-	2/4/4/4	-
2	DIO	A	1562	-	-	-	0/1/1/1
2	DIO	A	1563	-	-	-	0/1/1/1
2	DIO	A	1564	-	-	-	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1569	NGT	C7-S1	-11.08	1.67	1.77
4	A	1569	NGT	C4-C3	2.01	1.57	1.52

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1569	NGT	O4-C4-C5	-4.11	99.09	109.30
5	A	1570	NAG	O7-C7-C8	-3.85	114.91	122.06
5	A	1570	NAG	C4-C3-C2	2.71	114.99	111.02
2	A	1566	DIO	C2'-O1'-C1'	2.27	117.45	109.89
5	A	1570	NAG	O3-C3-C4	-2.22	105.22	110.35
4	A	1569	NGT	C8-C7-S1	2.21	122.28	118.96
4	A	1569	NGT	O5-C1-C2	2.19	120.08	115.27
5	A	1570	NAG	O7-C7-N2	2.13	125.86	121.95
2	A	1566	DIO	C2-O1-C1	2.04	116.71	109.89

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1568	PEG	C4-C3-O2-C2
3	A	1567	PEG	O1-C1-C2-O2
3	A	1568	PEG	O2-C3-C4-O4
3	A	1567	PEG	C4-C3-O2-C2

There are no ring outliers.

5 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1570	NAG	3	0
4	A	1569	NGT	2	0
3	A	1568	PEG	2	0
2	A	1562	DIO	3	0
2	A	1564	DIO	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	538/548 (98%)	-0.31	2 (0%) 92 91	9, 17, 30, 41	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	561	GLY	3.3
1	A	509	THR	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	DIO	A	1566	6/6	0.73	0.22	64,64,65,66	0
2	DIO	A	1564	6/6	0.74	0.20	60,61,62,62	0
3	PEG	A	1568	7/7	0.80	0.19	26,39,45,46	0
2	DIO	A	1563	6/6	0.83	0.19	65,66,67,68	0
3	PEG	A	1567	7/7	0.83	0.15	28,36,39,39	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	NAG	A	1570	14/15	0.87	0.13	28,35,39,40	0
2	DIO	A	1562	6/6	0.89	0.14	43,44,44,45	0
2	DIO	A	1565	6/6	0.93	0.12	46,46,46,47	0
4	NGT	A	1569	14/14	0.95	0.10	23,28,32,32	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.