



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 21, 2020 – 03:35 PM BST

PDB ID : 4WOJ
Title : Aspartate Semialdehyde Dehydrogenase from Francisella tularensis
Authors : Mank, N.J.; Arnette, A.K.; Klapper, V.G.; Chruszcz, M.
Deposited on : 2014-10-15
Resolution : 2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1544 (2.48-2.44)
Clashscore	141614	1613 (2.48-2.44)
Ramachandran outliers	138981	1598 (2.48-2.44)
Sidechain outliers	138945	1598 (2.48-2.44)
RSRZ outliers	127900	1523 (2.48-2.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	370	<div> <div>2%</div> <div> <div></div> <div>86%</div> <div>11%</div> <div>..</div> </div> </div>
1	B	370	<div> <div>%</div> <div> <div></div> <div>85%</div> <div>13%</div> <div>..</div> </div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5797 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aspartate semialdehyde dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	364	Total	C	N	O	S	0	1	0
			2827	1788	480	545	14			
1	B	365	Total	C	N	O	S	0	1	0
			2842	1798	481	549	14			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	HIS	-	expression tag	UNP Q8G8T7
A	-4	HIS	-	expression tag	UNP Q8G8T7
A	-3	HIS	-	expression tag	UNP Q8G8T7
A	-2	HIS	-	expression tag	UNP Q8G8T7
A	-1	HIS	-	expression tag	UNP Q8G8T7
A	0	HIS	-	expression tag	UNP Q8G8T7
B	-5	HIS	-	expression tag	UNP Q8G8T7
B	-4	HIS	-	expression tag	UNP Q8G8T7
B	-3	HIS	-	expression tag	UNP Q8G8T7
B	-2	HIS	-	expression tag	UNP Q8G8T7
B	-1	HIS	-	expression tag	UNP Q8G8T7
B	0	HIS	-	expression tag	UNP Q8G8T7

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Na	0	0
			1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	55	Total	O	0	0
			55	55		
4	B	57	Total	O	0	0
			57	57		

- Molecule 1: Aspartate semialdehyde dehydrogenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	46.59 Å 60.57 Å 67.68 Å 115.43° 100.98° 91.74°	Depositor
Resolution (Å)	37.22 – 2.45 37.20 – 2.44	Depositor EDS
% Data completeness (in resolution range)	97.4 (37.22-2.45) 96.8 (37.20-2.44)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.64 (at 2.45 Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.183 , 0.238 0.188 , 0.239	Depositor DCC
R_{free} test set	1190 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	33.2	Xtriage
Anisotropy	0.556	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 47.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.137 for -h,k,-k-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5797	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.64	0/2871	0.76	1/3886 (0.0%)
1	B	0.65	0/2886	0.77	3/3904 (0.1%)
All	All	0.65	0/5757	0.77	4/7790 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	322	ASP	CB-CG-OD1	5.39	123.15	118.30
1	A	307	ASP	CB-CG-OD1	5.24	123.02	118.30
1	B	102	ARG	NE-CZ-NH2	-5.04	117.78	120.30
1	B	117	ASP	CB-CG-OD1	5.02	122.82	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2827	0	2853	30	0
1	B	2842	0	2876	27	0
2	A	10	0	0	0	0
2	B	5	0	0	0	0
3	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	55	0	0	0	0
4	B	57	0	0	0	0
All	All	5797	0	5729	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:58:TYR:CD1	1:B:81:ILE:HD11	2.23	0.73
1:B:172:MET:HE3	1:B:267:LEU:HB3	1.76	0.67
1:A:185:ILE:HG21	1:A:202:LEU:CD2	2.24	0.66
1:A:283:THR:OG1	1:A:286:GLU:HG3	2.00	0.61
1:B:37:THR:O	1:B:37:THR:HG22	2.01	0.60
1:B:185:ILE:HG21	1:B:202:LEU:HD22	1.86	0.58
1:B:104:ASP:OD1	1:B:105:LYS:N	2.38	0.57
1:B:185:ILE:HG21	1:B:202:LEU:CD2	2.35	0.56
1:B:302:ASP:OD2	1:B:304:ASN:ND2	2.36	0.56
1:A:110:VAL:HG11	1:A:358:VAL:HG21	1.88	0.56
1:A:185:ILE:HG21	1:A:202:LEU:HD23	1.87	0.56
1:B:172:MET:CE	1:B:267:LEU:HB3	2.35	0.55
1:B:110:VAL:HG11	1:B:358:VAL:HG21	1.88	0.54
1:B:283:THR:OG1	1:B:286:GLU:HG3	2.09	0.53
1:B:16:VAL:CG1	1:B:348:LEU:HD23	2.39	0.52
1:A:222:LEU:HD21	1:B:161:TYR:HB3	1.92	0.51
1:B:360:ASN:OD1	1:B:363:LYS:HE3	2.10	0.51
1:A:186:ASP:HB2	1:A:199:LEU:HD21	1.93	0.50
1:A:117:ASP:N	1:A:117:ASP:OD2	2.45	0.50
1:B:186:ASP:HB2	1:B:199:LEU:HD21	1.94	0.50
1:A:222:LEU:HD21	1:B:161:TYR:CB	2.44	0.48
1:A:240:LYS:O	1:A:241:ALA:C	2.53	0.47
1:A:39:GLN:O	1:A:42:GLN:HB2	2.15	0.47
1:A:185:ILE:HG21	1:A:202:LEU:HD22	1.95	0.46
1:B:225:TRP:CE2	1:B:235:THR:HG22	2.52	0.45
1:A:193:LEU:CD2	1:B:12:MET:HE2	2.46	0.45
1:A:173:GLN:HG3	1:B:192:ILE:HD13	1.99	0.45
1:A:223:LEU:HB3	1:B:309:LEU:HD22	2.00	0.44
1:A:239:TYR:O	1:A:240:LYS:C	2.55	0.44
1:A:12:MET:HE2	1:B:193:LEU:CD2	2.47	0.44
1:A:57:ALA:O	1:A:63:LEU:HD11	2.18	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:57:ALA:O	1:B:63:LEU:HD11	2.19	0.43
1:A:236:LYS:O	1:A:240:LYS:HG3	2.19	0.42
1:A:75:GLY:O	1:A:79:LYS:HG3	2.20	0.42
1:A:46:GLY:O	1:A:49:GLN:HB3	2.19	0.42
1:A:12:MET:CE	1:A:172:MET:CE	2.98	0.42
1:A:38:SER:OG	1:A:39:GLN:N	2.50	0.42
1:A:102:ARG:HG2	1:A:109:LEU:HD21	2.01	0.42
1:A:220:TYR:HA	1:B:265:PRO:O	2.19	0.41
1:B:112:ASP:O	1:B:116:HIS:HB2	2.20	0.41
1:A:284:ILE:HA	1:A:284:ILE:HD12	1.95	0.41
1:B:353:GLU:N	1:B:354:PRO:CD	2.84	0.41
1:A:139:LEU:HA	1:A:139:LEU:HD23	1.88	0.41
1:A:188:ARG:HB3	1:A:188:ARG:CZ	2.51	0.40
1:A:80:GLU:O	1:A:84:LYS:HE2	2.21	0.40
1:B:32:PRO:HD2	1:B:51:TYR:CD1	2.56	0.40
1:A:11:GLY:O	1:A:15:SER:OG	2.36	0.40
1:A:94:TRP:HB3	1:A:129:ASP:OD1	2.21	0.40
1:B:102:ARG:HG2	1:B:109:LEU:HD21	2.03	0.40
1:B:97:ALA:HB2	1:B:355:LEU:HD11	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	363/370 (98%)	346 (95%)	16 (4%)	1 (0%)	41	49
1	B	364/370 (98%)	347 (95%)	16 (4%)	1 (0%)	41	49
All	All	727/740 (98%)	693 (95%)	32 (4%)	2 (0%)	41	49

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	227	ASP
1	B	227	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	312/324 (96%)	306 (98%)	6 (2%)	57	69
1	B	315/324 (97%)	308 (98%)	7 (2%)	52	64
All	All	627/648 (97%)	614 (98%)	13 (2%)	53	66

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	SER
1	A	80	GLU
1	A	128	LYS
1	A	206	SER
1	A	212	GLN
1	A	227	ASP
1	B	15	SER
1	B	45	THR
1	B	79	LYS
1	B	206	SER
1	B	227	ASP
1	B	228	VAL
1	B	239	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	125	ASN
1	B	83	HIS
1	B	125	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	402	-	4,4,4	0.51	0	6,6,6	0.28	0
2	SO4	B	401	-	4,4,4	0.57	0	6,6,6	0.36	0
2	SO4	A	401	-	4,4,4	0.45	0	6,6,6	0.24	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	364/370 (98%)	-0.15	6 (1%) 72 69	18, 35, 63, 85	0
1	B	365/370 (98%)	-0.21	3 (0%) 86 86	16, 36, 62, 82	0
All	All	729/740 (98%)	-0.18	9 (1%) 79 77	16, 36, 62, 85	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	231	PRO	4.5
1	A	228	VAL	3.2
1	B	41	GLY	3.2
1	A	61	ASP	2.8
1	B	88	ALA	2.4
1	A	232	SER	2.3
1	A	239	TYR	2.2
1	A	88	ALA	2.1
1	B	207	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	B	401	5/5	0.77	0.24	71,83,88,91	0
3	NA	B	402	1/1	0.87	0.09	31,31,31,31	0
2	SO4	A	402	5/5	0.90	0.33	103,105,110,112	0
2	SO4	A	401	5/5	0.93	0.12	81,83,86,92	0

6.5 Other polymers [i](#)

There are no such residues in this entry.