



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 1, 2021 – 11:14 PM EDT

PDB ID : 1WPV  
Title : Crystal Structure of Activated Binary complex of HutP, an RNA binding anti-termination protein  
Authors : Kumarevel, T.S.; Mizuno, H.; Kumar, P.K.R.  
Deposited on : 2004-09-14  
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.23.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	147	<div> <div>6%</div> <div>80%</div> <div>18%</div> <div>.</div> </div>
1	B	147	<div> <div>5%</div> <div>85%</div> <div>13%</div> <div>.</div> </div>
1	C	147	<div> <div>14%</div> <div>77%</div> <div>23%</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3899 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hut operon positive regulatory protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	147	Total	C	N	O	S	27	0	0
			1095	690	198	203	4			
1	B	147	Total	C	N	O	S	27	0	0
			1106	697	201	204	4			
1	C	147	Total	C	N	O	S	18	0	0
			1084	684	195	201	4			


There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	51	ILE	VAL	engineered mutation	UNP P10943
B	51	ILE	VAL	engineered mutation	UNP P10943
C	51	ILE	VAL	engineered mutation	UNP P10943

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		
2	B	1	Total	Mg	0	0
			1	1		
2	C	1	Total	Mg	0	0
			1	1		

- Molecule 3 is HISTIDINE (three-letter code: HIS) (formula: C<sub>6</sub>H<sub>10</sub>N<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	1	0
			11	6	3	2		
3	A	1	Total	C	N	O	1	0
			11	6	3	2		
3	C	1	Total	C	N	O	1	0
			11	6	3	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	193	Total	O	0	0
			193	193		
4	B	179	Total	O	0	0
			179	179		
4	C	206	Total	O	0	0
			206	206		



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	77.79Å 80.88Å 75.17Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.53 – 1.70 24.69 – 1.68	Depositor EDS
% Data completeness (in resolution range)	96.4 (19.53-1.70) 94.4 (24.69-1.68)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.71 (at 1.68Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.233 , 0.259 0.236 , 0.222	Depositor DCC
$R_{free}$ test set	2635 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	24.0	Xtriage
Anisotropy	0.153	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 52.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.015 for l,-k,h	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3899	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.18% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.32	0/1113	0.61	1/1505 (0.1%)
1	B	0.32	0/1124	0.64	1/1517 (0.1%)
1	C	0.32	0/1102	0.64	1/1492 (0.1%)
All	All	0.32	0/3339	0.63	3/4514 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	101	GLY	N-CA-C	-5.67	98.91	113.10
1	C	101	GLY	N-CA-C	-5.51	99.31	113.10
1	A	101	GLY	N-CA-C	-5.14	100.25	113.10

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1095	0	1077	21	0
1	B	1106	0	1103	21	0
1	C	1084	0	1060	26	1
2	A	1	0	0	0	0
2	B	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	1	0	0	0	0
3	A	22	0	12	0	0
3	C	11	0	6	1	0
4	A	193	0	0	6	0
4	B	179	0	0	9	0
4	C	206	0	0	6	0
All	All	3899	0	3258	66	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:20:GLU:HG3	4:B:3166:HOH:O	1.72	0.88
1:B:20:GLU:H	1:B:20:GLU:CD	1.77	0.86
1:B:20:GLU:CD	1:B:20:GLU:N	2.43	0.71
1:C:45:MET:HG2	1:C:99:THR:HB	1.72	0.69
1:B:13:SER:HB2	4:B:3176:HOH:O	1.90	0.69
1:C:103:ARG:HG3	4:C:3086:HOH:O	1.92	0.69
1:C:148:ILE:OXT	1:C:148:ILE:HG12	1.95	0.66
1:A:37:VAL:HG22	1:A:107:LEU:HD22	1.80	0.64
4:B:3140:HOH:O	1:C:133:ILE:HD12	1.99	0.62
1:A:45:MET:HG2	1:A:99:THR:HB	1.82	0.61
1:B:41:LYS:HE2	1:B:103:ARG:HD2	1.83	0.59
1:C:27:VAL:O	1:C:31:GLU:HG3	2.03	0.58
1:B:133:ILE:HG13	4:B:3096:HOH:O	2.03	0.58
1:C:38:CYS:HB2	1:C:63:VAL:HG21	1.86	0.57
1:A:38:CYS:HB2	1:A:63:VAL:HG21	1.86	0.56
1:A:40:GLY:HA3	1:A:104:PHE:CZ	2.43	0.54
1:C:22:GLU:O	1:C:24:SER:N	2.41	0.54
1:B:134:LYS:HA	1:B:134:LYS:HE2	1.89	0.53
1:C:131:ALA:HB1	4:C:3105:HOH:O	2.08	0.53
1:C:103:ARG:HB3	1:C:103:ARG:NH1	2.24	0.52
1:B:4:HIS:HB3	1:B:7:ARG:HB2	1.91	0.52
1:C:9:ILE:HD11	1:C:107:LEU:O	2.09	0.52
1:C:24:SER:CB	4:C:3190:HOH:O	2.57	0.51
1:B:45:MET:HG2	4:B:3082:HOH:O	2.10	0.51
1:A:37:VAL:HG22	1:A:107:LEU:CD2	2.41	0.51
1:A:60:LYS:N	1:A:60:LYS:HD3	2.27	0.50
1:C:61:SER:OG	1:C:63:VAL:HG23	2.12	0.50

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:25:THR:HA	4:A:2105:HOH:O	2.12	0.49
1:A:51:ILE:HD11	1:A:80:MET:HE1	1.95	0.49
1:C:111:PRO:HG2	1:C:112:TYR:CE1	2.47	0.49
1:B:18:LEU:HD21	1:B:141:PHE:CE2	2.49	0.48
1:B:148:ILE:HG12	1:B:148:ILE:OXT	2.13	0.48
1:C:38:CYS:HB3	1:C:106:VAL:HB	1.95	0.48
1:A:27:VAL:O	1:A:31:GLU:HG3	2.14	0.48
1:C:103:ARG:HB3	1:C:103:ARG:CZ	2.43	0.47
1:C:111:PRO:HG2	1:C:112:TYR:CD1	2.48	0.47
1:A:92:LEU:HD21	1:B:92:LEU:HD21	1.96	0.47
1:B:92:LEU:HD21	1:C:92:LEU:HD21	1.96	0.47
1:A:38:CYS:CB	1:A:63:VAL:HG21	2.45	0.47
1:B:87:THR:HB	1:B:90:GLU:O	2.14	0.46
1:C:35:TRP:HB3	1:C:107:LEU:HD11	1.96	0.46
1:B:135:GLY:HA3	4:B:3146:HOH:O	2.15	0.46
1:A:87:THR:HB	1:A:90:GLU:O	2.16	0.45
1:B:98:ARG:HD2	1:B:129:ILE:HD11	1.98	0.45
1:C:128:THR:HB	1:C:137:GLU:HG3	1.99	0.45
1:A:46:ASP:HB3	1:A:48:HIS:CE1	2.52	0.44
1:A:48:HIS:HD2	4:A:2169:HOH:O	1.99	0.44
1:B:13:SER:CB	4:B:3176:HOH:O	2.60	0.44
1:B:21:ALA:HB3	4:B:3049:HOH:O	2.17	0.44
1:C:114:SER:HB3	4:C:3193:HOH:O	2.17	0.43
1:A:101:GLY:O	1:A:127:GLY:HA3	2.18	0.43
1:C:36:LYS:HB2	1:C:108:ARG:HB3	2.01	0.43
1:B:101:GLY:O	1:B:127:GLY:HA3	2.18	0.43
1:C:134:LYS:HD3	4:C:3131:HOH:O	2.17	0.43
1:A:128:THR:HB	1:A:137:GLU:HG3	2.01	0.42
1:C:77:HIS:O	1:C:81:GLU:HG3	2.19	0.42
1:A:67:GLU:HB3	4:A:2188:HOH:O	2.19	0.42
1:B:45:MET:HE2	4:B:3163:HOH:O	2.20	0.41
1:C:87:THR:HB	1:C:90:GLU:O	2.20	0.41
1:A:90:GLU:HG2	4:A:2081:HOH:O	2.21	0.41
1:A:27:VAL:HG23	4:A:2111:HOH:O	2.20	0.41
1:B:84:HIS:HE1	4:C:3108:HOH:O	2.01	0.41
1:C:38:CYS:HB2	1:C:63:VAL:CG2	2.49	0.41
1:C:130:GLY:HA2	3:C:3001:HIS:CG	2.56	0.41
1:A:7:ARG:NH2	4:A:2037:HOH:O	2.54	0.40
1:A:36:LYS:HB2	1:A:108:ARG:HB3	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:70:ARG:NH2	1:C:148:ILE:O[2_655]	2.14	0.06

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	145/147 (99%)	139 (96%)	6 (4%)	0	100	100
1	B	145/147 (99%)	137 (94%)	8 (6%)	0	100	100
1	C	145/147 (99%)	136 (94%)	8 (6%)	1 (1%)	22	8
All	All	435/441 (99%)	412 (95%)	22 (5%)	1 (0%)	47	30

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	23	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	108/118 (92%)	102 (94%)	6 (6%)	21	7
1	B	111/118 (94%)	109 (98%)	2 (2%)	59	43
1	C	106/118 (90%)	106 (100%)	0	100	100
All	All	325/354 (92%)	317 (98%)	8 (2%)	49	29

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	LEU
1	A	44	SER
1	A	60	LYS
1	A	70	ARG
1	A	108	ARG
1	A	137	GLU
1	B	20	GLU
1	B	103	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	26	GLN
1	A	65	GLN
1	B	4	HIS
1	B	84	HIS
1	C	48	HIS
1	C	65	GLN
1	C	84	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 3 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	HIS	A	1001	2	4,11,11	0.66	0	3,14,14	1.28	1 (33%)
3	HIS	C	3001	2	4,11,11	0.63	0	3,14,14	1.30	1 (33%)
3	HIS	A	2001	2	4,11,11	0.61	0	3,14,14	1.29	1 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HIS	A	1001	2	-	0/4/8/8	0/1/1/1
3	HIS	C	3001	2	-	0/4/8/8	0/1/1/1
3	HIS	A	2001	2	-	0/4/8/8	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	3001	HIS	CD2-NE2-CE1	2.07	109.02	105.78
3	A	1001	HIS	CD2-NE2-CE1	2.04	108.97	105.78
3	A	2001	HIS	CD2-NE2-CE1	2.03	108.95	105.78

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	3001	HIS	1	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	147/147 (100%)	0.54	9 (6%) 21 23	17, 26, 45, 66	14 (9%)
1	B	147/147 (100%)	0.46	7 (4%) 30 34	16, 24, 40, 67	15 (10%)
1	C	147/147 (100%)	0.92	21 (14%) 2 2	21, 31, 52, 68	12 (8%)
All	All	441/441 (100%)	0.64	37 (8%) 11 12	16, 27, 49, 68	41 (9%)

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	24	SER	8.7
1	C	24	SER	8.3
1	B	21	ALA	8.0
1	C	23	GLU	7.6
1	A	21	ALA	7.5
1	B	23	GLU	7.4
1	A	24	SER	6.8
1	C	21	ALA	6.1
1	A	20	GLU	5.8
1	A	23	GLU	5.8
1	C	25	THR	5.6
1	B	22	GLU	5.2
1	C	22	GLU	4.9
1	C	116	ALA	4.0
1	A	25	THR	4.0
1	B	20	GLU	3.7
1	A	22	GLU	3.7
1	C	114	SER	3.4
1	C	4	HIS	3.3
1	C	20	GLU	3.2
1	C	134	LYS	3.0
1	C	33	ASP	2.9
1	C	30	LEU	2.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	2	THR	2.9
1	C	6	GLU	2.7
1	C	56	THR	2.6
1	A	19	ASN	2.6
1	C	133	ILE	2.5
1	B	116	ALA	2.4
1	A	18	LEU	2.4
1	B	145	ILE	2.3
1	C	112	TYR	2.3
1	C	121	ILE	2.3
1	C	148	ILE	2.2
1	C	63	VAL	2.1
1	C	68	GLY	2.1
1	A	123	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MG	C	1002	1/1	0.94	0.14	21,21,21,21	0
3	HIS	A	1001	11/11	0.95	0.10	17,21,24,25	1
3	HIS	A	2001	11/11	0.95	0.10	16,17,20,20	1
3	HIS	C	3001	11/11	0.96	0.09	15,17,21,22	1
2	MG	B	3002	1/1	0.97	0.14	15,15,15,15	0
2	MG	A	2002	1/1	1.00	0.12	13,13,13,13	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.