



## wwPDB EM Validation Summary Report ⓘ

Nov 13, 2022 – 08:55 AM EST

PDB ID : 6WR4  
EMDB ID : EMD-21876  
Title : Structure of human ATG9A, the only transmembrane protein of the core autophagy machinery  
Authors : Guardia, C.M.; Tan, X.; Lian, T.; Rana, M.S.; Zhou, W.; Christenson, E.T.; Lowry, A.J.; Faraldo-Gomez, J.D.; Bonifacino, J.S.; Jiang, J.; Banerjee, A.  
Deposited on : 2020-04-29  
Resolution : 2.90 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
MolProbity : 4.02b-467  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
MapQ : 1.9.9  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.31.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	839	
1	B	839	
1	C	839	

2 Entry composition ⓘ


There are 2 unique types of molecules in this entry. The entry contains 11934 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Autophagy-related protein 9A.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	470	Total	C	N	O	S	0	0
			3862	2550	647	645	20		
1	B	470	Total	C	N	O	S	0	0
			3862	2550	647	645	20		
1	C	470	Total	C	N	O	S	0	0
			3862	2550	647	645	20		

- Molecule 2 is Lauryl Maltose Neopentyl Glycol (three-letter code: LMN) (formula: C<sub>47</sub>H<sub>88</sub>O<sub>22</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			AltConf
2	A	1	Total	C	O	0
			116	82	34	
2	A	1	Total	C	O	0
			116	82	34	
2	B	1	Total	C	O	0
			116	82	34	

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			AltConf
2	B	1	Total 116	C 82	O 34	0
2	C	1	Total 116	C 82	O 34	0
2	C	1	Total 116	C 82	O 34	0





HIS	PHE	SER	ARG	LEU	PRO	LEU	GLY	GLY	GLY	TRP	ALA	ALA	GLU	ASP	ARG	ALA	PRO	GLN	SER	ILE	SER	ARG	PRO	PRO	CHS	ALA	ALA	PRO	ARG	PRO	GLY	ALA	GLY	PRO	GLU	THR	THR	ALA	LEU	HIS	GLY	GLY	PHE	GLN	ARG	ARG	THR	GLY	GLY	ILE	THR	ASP	PRO	THR	VAL	ARG	VAL	PRO	SER
GLU	ALA	PRO	ASP	GLY	GLY	GLY	GLY	GLY	GLY	GLY	ALA	ALA	ARG	ALA	PRO	GLN	SER	ILE	SER	ARG	PRO	PRO	CHS	ALA	ALA	PRO	ARG	PRO	PRO	GLY	ALA	GLY	PRO	GLU	THR	THR	ALA	LEU	HIS	GLY	GLY	PHE	GLN	ARG	ARG	THR	GLY	GLY	ILE	THR	ASP	PRO	THR	VAL	ARG	VAL	PRO	SER	

## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	646291	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	57	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	6.751	Depositor
Minimum map value	-4.148	Depositor
Average map value	0.009	Depositor
Map value standard deviation	0.160	Depositor
Recommended contour level	0.538	Depositor
Map size ( $\text{\AA}$ )	203.51999, 203.51999, 203.51999	wwPDB
Map dimensions	192, 192, 192	wwPDB
Map angles ( $^\circ$ )	90.0, 90.0, 90.0	wwPDB
Pixel spacing ( $\text{\AA}$ )	1.06, 1.06, 1.06	Depositor



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: LMN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.38	0/3969	0.65	0/5392
1	B	0.38	0/3969	0.64	0/5392
1	C	0.38	0/3969	0.64	0/5392
All	All	0.38	0/11907	0.64	0/16176

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
1	B	0	4
1	C	0	4
All	All	0	12

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 12 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	36	PRO	Peptide
1	A	404	VAL	Peptide
1	A	434	PRO	Peptide
1	A	436	GLN	Peptide
1	B	36	PRO	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3862	0	3894	33	0
1	B	3862	0	3894	36	0
1	C	3862	0	3894	32	0
2	A	116	0	155	6	0
2	B	116	0	155	7	0
2	C	116	0	155	7	0
All	All	11934	0	12147	100	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 100 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:217:ASN:HD21	1:B:344:ASN:H	1.40	0.69
1:A:217:ASN:HD21	1:A:344:ASN:H	1.40	0.69
1:C:217:ASN:HD21	1:C:344:ASN:H	1.40	0.68
1:B:254:ILE:O	1:B:289:ARG:NH1	2.30	0.65
1:A:254:ILE:O	1:A:289:ARG:NH1	2.30	0.65

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	464/839 (55%)	438 (94%)	24 (5%)	2 (0%)	34 66

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	B	464/839 (55%)	438 (94%)	24 (5%)	2 (0%)	34 66
1	C	464/839 (55%)	438 (94%)	24 (5%)	2 (0%)	34 66
All	All	1392/2517 (55%)	1314 (94%)	72 (5%)	6 (0%)	38 66

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	203	LEU
1	B	203	LEU
1	C	203	LEU
1	A	229	PRO
1	B	229	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	416/717 (58%)	400 (96%)	16 (4%)	33 67
1	B	416/717 (58%)	400 (96%)	16 (4%)	33 67
1	C	416/717 (58%)	400 (96%)	16 (4%)	33 67
All	All	1248/2151 (58%)	1200 (96%)	48 (4%)	36 67

5 of 48 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	423	SER
1	C	186	ARG
1	B	437	LEU
1	C	80	THR
1	C	205	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	57	ASN
1	C	278	GLN
1	C	217	ASN
1	C	348	HIS
1	A	457	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	LMN	C	902	-	48,48,72	1.87	10 (20%)	61,62,98	1.09	4 (6%)
2	LMN	C	901	-	72,72,72	1.67	12 (16%)	96,98,98	1.49	15 (15%)
2	LMN	B	901	-	72,72,72	1.67	12 (16%)	96,98,98	1.49	15 (15%)
2	LMN	A	901	-	72,72,72	1.68	12 (16%)	96,98,98	1.49	15 (15%)
2	LMN	B	902	-	48,48,72	1.87	10 (20%)	61,62,98	1.09	4 (6%)
2	LMN	A	902	-	48,48,72	1.87	10 (20%)	61,62,98	1.08	4 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LMN	C	902	-	-	25/39/79/130	0/2/2/4
2	LMN	C	901	-	-	23/50/130/130	0/4/4/4
2	LMN	B	901	-	-	23/50/130/130	0/4/4/4
2	LMN	A	901	-	-	23/50/130/130	0/4/4/4
2	LMN	B	902	-	-	25/39/79/130	0/2/2/4
2	LMN	A	902	-	-	25/39/79/130	0/2/2/4

The worst 5 of 66 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	902	LMN	O5-C1	5.83	1.56	1.41
2	C	902	LMN	O5-C1	5.82	1.56	1.41
2	B	902	LMN	O5-C1	5.80	1.56	1.41
2	A	901	LMN	O5-C1	5.39	1.55	1.41
2	C	901	LMN	O5-C1	5.39	1.55	1.41

The worst 5 of 57 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	901	LMN	O1-C1-C2	5.10	116.26	108.30
2	C	901	LMN	O1-C1-C2	5.07	116.21	108.30
2	B	901	LMN	O1-C1-C2	5.05	116.19	108.30
2	B	901	LMN	CCL-CCH-CCQ	4.06	118.96	109.68
2	C	901	LMN	CCL-CCH-CCQ	4.05	118.93	109.68

There are no chirality outliers.

5 of 144 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	901	LMN	C2-C1-O1-CBS
2	A	901	LMN	O5-C1-O1-CBS
2	A	901	LMN	OBX-CCJ-OBV-CBT
2	A	901	LMN	CCL-CCJ-OBV-CBT
2	A	902	LMN	CBK-CBQ-CCM-CBR


There are no ring outliers.


6 monomers are involved in 20 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	902	LMN	3	0
2	C	901	LMN	4	0
2	B	901	LMN	4	0
2	A	901	LMN	3	0
2	B	902	LMN	3	0
2	A	902	LMN	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-21876. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X



Y




Z

The images above show the map projected in three orthogonal directions.

### 6.2 Central slices [i](#)

#### 6.2.1 Primary map



X Index: 96



Y Index: 96



Z Index: 96

The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)

### 6.3.1 Primary map



X Index: 88



Y Index: 107




Z Index: 83


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal surface views [i](#)

### 6.4.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.538. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


## 6.5 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 73 nm<sup>3</sup>; this corresponds to an approximate mass of 66 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum ⓘ



\*Reported resolution corresponds to spatial frequency of 0.345 Å<sup>-1</sup>




## 8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.


## 9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-21876 and PDB model 6WR4. Per-residue inclusion information can be found in [section 3](#) on [page 5](#).


### 9.1 Map-model overlay [i](#)



X




Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.538 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.538).

## 9.4 Atom inclusion [i](#)



At the recommended contour level, 85% of all backbone atoms, 79% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.538) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div></div> 0.7929	<div></div> 0.5260
A	<div></div> 0.7915	<div></div> 0.5250
B	<div></div> 0.7918	<div></div> 0.5250
C	<div></div> 0.7954	<div></div> 0.5270

