



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 26, 2020 – 06:44 pm BST

PDB ID : 4WU9  
Title : Structure of cisPtNAP-NCP145  
Authors : Chua, E.Y.D.; Davey, G.E.; Chin, C.F.; Droge, P.; Ang, W.H.; Davey, C.A.  
Deposited on : 2014-10-31  
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	135	<div> <div>2%</div> <div> <div></div> <div>57%</div> <div>14%</div> <div>29%</div> </div> </div>
1	E	135	<div> <div>2%</div> <div> <div></div> <div>61%</div> <div>9%</div> <div>29%</div> </div> </div>
2	B	102	<div> <div>2%</div> <div> <div></div> <div>63%</div> <div>18%</div> <div>20%</div> </div> </div>
2	F	102	<div> <div>%</div> <div> <div></div> <div>70%</div> <div>16%</div> <div>15%</div> </div> </div>
3	C	129	<div> <div>2%</div> <div> <div></div> <div>64%</div> <div>16%</div> <div>18%</div> </div> </div>
3	G	129	<div> <div>2%</div> <div> <div></div> <div>70%</div> <div>12%</div> <div>18%</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	D	125	<div><div><div></div><div></div><div></div></div><div><div>%</div><div>64%</div><div>11%</div><div>24%</div></div></div>
4	H	125	<div><div><div></div><div></div><div></div></div><div><div>%</div><div>66%</div><div>10%</div><div>24%</div></div></div>
5	I	145	<div><div><div></div><div></div><div></div></div><div><div>3%</div><div>40%</div><div>46%</div><div>14%</div></div></div>
6	J	145	<div><div><div></div><div></div><div></div></div><div><div>2%</div><div>48%</div><div>38%</div><div>14%</div></div></div>

## 2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 12110 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone H3.2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	96	Total	C	N	O	S	0	0	0
			791	500	151	137	3			
1	E	96	Total	C	N	O	S	0	0	0
			791	500	151	137	3			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	102	ALA	GLY	engineered mutation	UNP P84233
E	102	ALA	GLY	engineered mutation	UNP P84233

- Molecule 2 is a protein called Histone H4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	82	Total	C	N	O	S	0	0	0
			653	412	127	113	1			
2	F	87	Total	C	N	O	S	0	0	0
			703	442	142	118	1			

- Molecule 3 is a protein called Histone H2A type 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	106	Total	C	N	O		0	0	0
			818	516	160	142				
3	G	106	Total	C	N	O		0	0	0
			818	516	160	142				

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	99	ARG	GLY	engineered mutation	UNP P06897
C	123	SER	ALA	engineered mutation	UNP P06897

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
G	99	ARG	GLY	engineered mutation	UNP P06897
G	123	SER	ALA	engineered mutation	UNP P06897

- Molecule 4 is a protein called Histone H2B 1.1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	95	Total	C	N	O	S	0	0	0
			745	469	134	140	2			
4	H	95	Total	C	N	O	S	0	0	0
			745	469	134	140	2			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	29	THR	SER	engineered mutation	UNP P02281
H	29	THR	SER	engineered mutation	UNP P02281

- Molecule 5 is a DNA chain called DNA (145-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	I	145	Total	C	N	O	P	0	0	0
			2970	1421	538	867	144			

- Molecule 6 is a DNA chain called DNA (145-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	J	145	Total	C	N	O	P	0	0	0
			2969	1421	535	869	144			

- Molecule 7 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	C	1	Total	O	S	0	0
			5	4	1		
7	H	1	Total	O	S	0	0
			5	4	1		

- Molecule 8 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	E	1	Total	Mg	0	0
			1	1		

- Molecule 9 is [2-{3-[(2-{[2-(amino-kappaN)ethyl]amino-kappaN}ethyl)amino-kappaN]propyl}-1H-benzo[de]isoquinoline-1,3(2H)-dionato(3-)]platinum (three-letter code: CX8) (formula: C<sub>19</sub>H<sub>21</sub>N<sub>4</sub>O<sub>2</sub>Pt).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
9	I	1	Total	C	N	O	Pt	0	0
			26	19	4	2	1		
9	I	1	Total	C	N	O	Pt	0	0
			26	19	4	2	1		
9	J	1	Total	C	N	O	Pt	0	0
			26	19	4	2	1		


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	2	Total	O	0	0
			2	2		
10	C	5	Total	O	0	0
			5	5		
10	D	1	Total	O	0	0
			1	1		
10	E	4	Total	O	0	0
			4	4		
10	F	4	Total	O	0	0
			4	4		
10	G	1	Total	O	0	0
			1	1		
10	H	1	Total	O	0	0
			1	1		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Histone H3.2




- Molecule 1: Histone H3.2



- Molecule 2: Histone H4




- Molecule 2: Histone H4




- Molecule 3: Histone H2A type 1









• Molecule 3: Histone H2A type 1



• Molecule 4: Histone H2B 1.1




• Molecule 4: Histone H2B 1.1



• Molecule 5: DNA (145-MER)



• Molecule 6: DNA (145-MER)



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	106.57Å 109.33Å 181.28Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	70.33 – 2.60 58.38 – 2.60	Depositor EDS
% Data completeness (in resolution range)	88.6 (70.33-2.60) 88.6 (58.38-2.60)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.77 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.216 , 0.273 0.215 , 0.265	Depositor DCC
$R_{free}$ test set	1174 reflections (2.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	74.1	Xtriage
Anisotropy	0.064	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.27 , 50.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.022 for k,h,-l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	12110	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	102.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.88% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CX8, MG, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.48	0/803	0.63	0/1078
1	E	0.56	0/803	0.70	0/1078
2	B	0.45	0/660	0.67	0/883
2	F	0.60	0/711	0.82	0/948
3	C	0.54	0/828	0.76	0/1117
3	G	0.46	0/828	0.62	0/1117
4	D	0.57	0/756	0.69	0/1015
4	H	0.49	0/756	0.64	0/1015
5	I	0.53	6/3332 (0.2%)	1.15	44/5141 (0.9%)
6	J	0.53	6/3330 (0.2%)	1.14	40/5138 (0.8%)
All	All	0.52	12/12807 (0.1%)	0.97	84/18530 (0.5%)

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	I	-14	DG	P-O5'	13.59	1.73	1.59
6	J	-14	DG	P-O5'	13.42	1.73	1.59
6	J	-14	DG	C5-C4	12.69	1.47	1.38
5	I	-14	DG	C5-C4	12.42	1.47	1.38
5	I	-14	DG	N7-C5	-11.46	1.32	1.39

The worst 5 of 84 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	I	-14	DG	N3-C4-C5	-22.97	117.11	128.60
6	J	-14	DG	N3-C4-C5	-21.91	117.65	128.60
5	I	-14	DG	N3-C4-N9	17.16	136.30	126.00
5	I	-14	DG	C2-N3-C4	17.11	120.45	111.90
6	J	-14	DG	C2-N3-C4	16.78	120.29	111.90

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	791	0	828	18	0
1	E	791	0	828	10	0
2	B	653	0	696	16	0
2	F	703	0	755	10	0
3	C	818	0	877	18	0
3	G	818	0	877	11	0
4	D	745	0	773	8	0
4	H	745	0	773	11	0
5	I	2970	0	1640	62	0
6	J	2969	0	1641	63	0
7	C	5	0	0	0	0
7	H	5	0	0	0	0
8	E	1	0	0	0	0
9	I	52	0	40	2	0
9	J	26	0	20	3	0
10	A	2	0	0	1	0
10	C	5	0	0	1	0
10	D	1	0	0	0	0
10	E	4	0	0	0	0
10	F	4	0	0	0	0
10	G	1	0	0	0	0
10	H	1	0	0	0	0
All	All	12110	0	9748	189	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 189 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:G:17:ARG:HH12	3:G:31:HIS:HD2	1.25	0.85
9:J:100:CX8:H11	9:J:100:CX8:O1	1.74	0.85

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:49:ARG:HG3	1:E:49:ARG:HH11	1.44	0.81
2:B:23:ARG:HH11	2:B:23:ARG:HG2	1.43	0.81
5:I:-44:DA:H2'	5:I:-43:DA:C8	2.18	0.79

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	94/135 (70%)	92 (98%)	1 (1%)	1 (1%)	14	30
1	E	94/135 (70%)	90 (96%)	4 (4%)	0	100	100
2	B	80/102 (78%)	74 (92%)	6 (8%)	0	100	100
2	F	85/102 (83%)	82 (96%)	3 (4%)	0	100	100
3	C	104/129 (81%)	95 (91%)	9 (9%)	0	100	100
3	G	104/129 (81%)	96 (92%)	8 (8%)	0	100	100
4	D	93/125 (74%)	86 (92%)	5 (5%)	2 (2%)	6	12
4	H	93/125 (74%)	89 (96%)	3 (3%)	1 (1%)	14	30
All	All	747/982 (76%)	704 (94%)	39 (5%)	4 (0%)	29	52

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	H	101	GLY
1	A	81	ASP
4	D	101	GLY
4	D	116	THR

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	84/110 (76%)	82 (98%)	2 (2%)	49	74
1	E	84/110 (76%)	81 (96%)	3 (4%)	35	61
2	B	67/78 (86%)	67 (100%)	0	100	100
2	F	72/78 (92%)	67 (93%)	5 (7%)	15	31
3	C	84/101 (83%)	79 (94%)	5 (6%)	19	39
3	G	84/101 (83%)	80 (95%)	4 (5%)	25	49
4	D	81/105 (77%)	74 (91%)	7 (9%)	10	20
4	H	81/105 (77%)	78 (96%)	3 (4%)	34	60
All	All	637/788 (81%)	608 (95%)	29 (5%)	27	51

5 of 29 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	D	102	GLU
1	E	120	MET
4	H	33	SER
1	E	49	ARG
2	F	19	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	68	GLN
2	F	18	HIS
4	H	46	HIS
4	D	92	GLN
4	H	44	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
7	SO4	C	201	-	4,4,4	0.50	0	6,6,6	0.33	0
7	SO4	H	201	-	4,4,4	0.50	0	6,6,6	0.58	0
9	CX8	I	102	-	26,30,30	1.74	4 (15%)	26,44,44	1.74	3 (11%)
9	CX8	J	100	6	26,30,30	1.66	3 (11%)	26,44,44	1.86	2 (7%)
9	CX8	I	101	5	26,30,30	1.89	5 (19%)	26,44,44	1.90	2 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	CX8	I	102	-	-	3/5/25/25	0/5/5/5
9	CX8	J	100	6	-	1/5/25/25	0/5/5/5
9	CX8	I	101	5	-	3/5/25/25	0/5/5/5

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	I	101	CX8	C12-C11	5.18	1.50	1.41

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	I	101	CX8	C8-C9	4.97	1.50	1.41
9	J	100	CX8	C12-C11	4.75	1.50	1.41
9	I	102	CX8	C8-C9	4.63	1.49	1.41
9	I	102	CX8	C12-C11	4.54	1.49	1.41

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	J	100	CX8	C9-C8-N4	6.48	120.57	116.15
9	I	101	CX8	C11-C12-N4	6.45	120.55	116.15
9	I	101	CX8	C9-C8-N4	6.16	120.36	116.15
9	J	100	CX8	C11-C12-N4	5.76	120.08	116.15
9	I	102	CX8	C9-C8-N4	5.30	119.77	116.15

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
9	I	102	CX8	C6-C7-N4-C8
9	I	102	CX8	C5-C6-C7-N4
9	I	101	CX8	C6-C7-N4-C12
9	I	101	CX8	C5-C6-C7-N4
9	J	100	CX8	N1-C5-C6-C7

There are no ring outliers.


3 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
9	I	102	CX8	1	0
9	J	100	CX8	3	0
9	I	101	CX8	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.



The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	96/135 (71%)	0.31	3 (3%) 49 42	51, 72, 107, 159	0
1	E	96/135 (71%)	0.26	3 (3%) 49 42	38, 53, 83, 144	0
2	B	82/102 (80%)	0.47	2 (2%) 59 53	52, 70, 97, 154	0
2	F	87/102 (85%)	0.36	1 (1%) 80 78	42, 52, 75, 157	0
3	C	106/129 (82%)	0.26	2 (1%) 66 62	39, 60, 90, 146	0
3	G	106/129 (82%)	0.24	3 (2%) 53 46	49, 72, 112, 147	0
4	D	95/125 (76%)	0.41	1 (1%) 80 78	44, 61, 96, 145	0
4	H	95/125 (76%)	0.30	1 (1%) 80 78	47, 72, 116, 160	0
5	I	145/145 (100%)	0.05	4 (2%) 53 46	77, 132, 180, 217	0
6	J	145/145 (100%)	0.07	3 (2%) 63 58	79, 135, 181, 199	0
All	All	1053/1272 (82%)	0.25	23 (2%) 62 56	38, 73, 160, 217	0

The worst 5 of 23 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	H	122	LYS	4.8
5	I	-72	DA	4.4
4	D	122	LYS	4.1
1	E	39	HIS	3.6
2	F	16	LYS	3.6

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ


There are no carbohydrates in this entry.

### 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
9	CX8	J	100	26/26	0.79	0.23	122,151,197,200	0
9	CX8	I	101	26/26	0.79	0.25	145,163,230,238	0
9	CX8	I	102	26/26	0.94	0.17	152,170,181,185	0
7	SO4	C	201	5/5	0.95	0.14	90,98,103,104	0
7	SO4	H	201	5/5	0.97	0.12	84,85,93,102	0
8	MG	E	201	1/1	0.98	0.15	45,45,45,45	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around CX8 I 101:**

$2mF_o - DF_c$  (at 0.7 rmsd) in gray  
 $mF_o - DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around CX8 I 102:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.