



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 19, 2021 – 10:07 AM EDT

PDB ID : 6WVZ
Title : Crystal structure of anti-MET Fab arm of amivantamab in complex with human MET
Authors : Cardoso, R.M.F.
Deposited on : 2020-05-07
Resolution : 3.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.18
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.18


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1094 (3.10-3.10)
Clashscore	141614	1184 (3.10-3.10)
Ramachandran outliers	138981	1141 (3.10-3.10)
Sidechain outliers	138945	1141 (3.10-3.10)
RSRZ outliers	127900	1067 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	228	 86% 8% 6%
2	L	214	 2% 84% 16%
3	M	534	 2% 82% 15% .
4	A	2	 50% 50%
4	C	2	 50% 50%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	B	3	 33% 67%
5	D	3	 33% 67%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NAG	A	2	-	-	-	X
4	NAG	C	2	-	-	-	X
5	NAG	B	2	-	-	-	X
5	FUC	B	3	-	-	-	X
5	NAG	D	2	-	-	-	X
5	FUC	D	3	-	-	-	X
6	NAG	M	606	-	-	-	X

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 7260 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heavy Chain of anti-MET Fab of amivantamab.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	H	215	Total	C	N	O	S	0	0	0
			1588	1002	262	316	8			

- Molecule 2 is a protein called Light Chain of anti-MET Fab of amivantamab.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	213	Total	C	N	O	S	0	0	0
			1556	975	266	310	5			

- Molecule 3 is a protein called Hepatocyte growth factor receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	M	518	Total	C	N	O	S	0	1	0
			3915	2505	661	719	30			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
M	565	HIS	-	expression tag	UNP P08581
M	566	HIS	-	expression tag	UNP P08581
M	567	HIS	-	expression tag	UNP P08581
M	568	HIS	-	expression tag	UNP P08581
M	569	HIS	-	expression tag	UNP P08581
M	570	HIS	-	expression tag	UNP P08581
M	571	HIS	-	expression tag	UNP P08581
M	572	HIS	-	expression tag	UNP P08581

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	A	2	Total	C	N	O	0	0	0
			28	16	2	10			
4	C	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 5 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	B	3	Total	C	N	O	0	0	0
			38	22	2	14			
5	D	3	Total	C	N	O	0	0	0
			38	22	2	14			

- Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	M	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 7 is FORMIC ACID (three-letter code: FMT) (formula: CH_2O_2).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	M	1	Total	C	O	0	0
			3	1	2		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	H	9	Total	O	0	0
			9	9		
8	L	13	Total	O	0	0
			13	13		
8	M	30	Total	O	0	0
			30	30		

Chain A:  50% 50%


MAG1
MAG2

- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:  50% 50%

MAG1
MAG2

- Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain B:  33% 67%

MAG1
MAG2
FUC3

- Molecule 5: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:  33% 67%

MAG1
MAG2
FUC3

4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	86.83Å 86.83Å 457.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.45 – 3.10 40.58 – 3.10	Depositor EDS
% Data completeness (in resolution range)	94.9 (40.45-3.10) 94.9 (40.58-3.10)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.64 (at 3.12Å)	Xtriage
Refinement program	PHENIX 1.10_2155	Depositor
R, R_{free}	0.190 , 0.235 0.191 , 0.236	Depositor DCC
R_{free} test set	1592 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	88.5	Xtriage
Anisotropy	0.455	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 74.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7260	wwPDB-VP
Average B, all atoms (Å ²)	98.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FUC, NAG, FMT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	H	0.26	0/1628	0.47	0/2231
2	L	0.26	0/1592	0.46	0/2176
3	M	0.26	0/4017	0.46	0/5480
All	All	0.26	0/7237	0.46	0/9887

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1588	0	1487	11	0
2	L	1556	0	1424	19	0
3	M	3915	0	3584	46	0
4	A	28	0	25	0	0
4	C	28	0	25	1	0
5	B	38	0	34	0	0
5	D	38	0	34	3	0
6	M	14	0	13	1	0
7	M	3	0	1	0	0
8	H	9	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	L	13	0	0	0	0
8	M	30	0	0	1	0
All	All	7260	0	6627	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:120:PRO:HD3	2:L:132:VAL:HG22	1.68	0.74
5:D:1:NAG:H83	5:D:1:NAG:H3	1.72	0.72
3:M:68:ASN:HB3	3:M:87:GLY:HA2	1.71	0.71
1:H:187:VAL:HG21	2:L:135:LEU:HD11	1.75	0.68
2:L:118:PHE:HB2	2:L:133:VAL:HG22	1.76	0.68
3:M:299:ILE:HG22	3:M:313:VAL:HG12	1.76	0.67
1:H:6:GLN:HE21	1:H:113:THR:HG23	1.60	0.66
2:L:2:ILE:HG13	2:L:26:SER:HB3	1.82	0.61
3:M:269:LEU:HD11	3:M:387:GLN:HB3	1.80	0.61
3:M:325:PRO:HD3	3:M:341:ILE:HD13	1.82	0.61
3:M:279:ILE:HG12	3:M:293:GLU:HG2	1.81	0.61
4:C:2:NAG:H3	4:C:2:NAG:H83	1.85	0.58
3:M:180:LEU:HD22	3:M:202:ASN:HB2	1.87	0.56
3:M:481:LEU:HD11	3:M:513:ILE:HD11	1.88	0.56
3:M:260:TYR:HA	3:M:279:ILE:O	2.06	0.55
3:M:175:CYS:O	3:M:217:ARG:NH1	2.32	0.55
2:L:166:GLN:HE21	2:L:171:SER:HB3	1.70	0.55
3:M:290:SER:N	3:M:415:GLU:OE2	2.26	0.55
3:M:464:GLN:HB2	3:M:477:VAL:HB	1.89	0.54
3:M:541:CYS:O	3:M:543:ASP:N	2.41	0.54
1:H:91:THR:HG23	1:H:116:THR:HA	1.91	0.53
5:D:1:NAG:H61	5:D:2:NAG:C7	2.37	0.53
3:M:72:VAL:HB	3:M:82:ALA:HB3	1.90	0.53
1:H:35:SER:HG	1:H:106:PHE:HE1	1.57	0.52
3:M:339:ASP:OD2	3:M:368:LYS:HE2	2.10	0.52
3:M:378:VAL:HG23	3:M:383:VAL:HG11	1.92	0.51
3:M:235:ILE:HG23	3:M:389:PHE:CD1	2.45	0.51
2:L:108:ARG:HH12	2:L:111:ALA:HB2	1.75	0.51
3:M:274:PHE:O	3:M:315:ASN:ND2	2.43	0.51
3:M:123:ASP:O	3:M:128:ASP:HA	2.10	0.51
1:H:154:GLU:OE2	1:H:174:ALA:HB3	2.11	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:M:526:CYS:HB2	3:M:540:TRP:CG	2.46	0.51
3:M:277:ARG:HD2	3:M:293:GLU:HB3	1.93	0.50
3:M:212:HIS:HA	3:M:236:ASP:OD2	2.11	0.50
2:L:92:ASN:ND2	3:M:99:GLN:HA	2.28	0.49
3:M:429:LEU:HG	3:M:469:ARG:HH21	1.78	0.48
2:L:161:GLU:HG2	2:L:175:LEU:HD21	1.95	0.48
3:M:297:GLU:O	3:M:425:GLN:HA	2.13	0.48
3:M:204:SER:O	3:M:207:PRO:HD2	2.14	0.48
1:H:6:GLN:NE2	1:H:113:THR:HG23	2.28	0.48
3:M:118:MET:CB	3:M:177:VAL:HG11	2.44	0.47
2:L:147:GLN:NE2	2:L:195:GLU:OE1	2.35	0.47
3:M:442:ILE:HA	3:M:454:ASN:O	2.14	0.47
3:M:243:ASP:OD1	3:M:243:ASP:N	2.48	0.47
2:L:66:GLY:HA3	2:L:71:PHE:HA	1.97	0.46
3:M:64:LEU:HD11	3:M:73:LEU:HD11	1.96	0.46
2:L:18:ARG:HA	2:L:75:ILE:O	2.16	0.45
3:M:245:TYR:OH	3:M:277:ARG:NH2	2.49	0.45
5:D:1:NAG:H3	5:D:1:NAG:C8	2.44	0.45
3:M:172:CYS:SG	3:M:175:CYS:N	2.90	0.44
3:M:345:VAL:HG11	3:M:439:LEU:HB2	1.99	0.44
3:M:54[B]:ASN:ND2	8:M:702:HOH:O	2.50	0.43
3:M:267:GLU:HA	3:M:275:HIS:CE1	2.53	0.43
1:H:2:VAL:HG13	1:H:27:TYR:HD1	1.83	0.43
3:M:221:GLU:H	3:M:221:GLU:CD	2.22	0.43
3:M:166:ILE:HD12	3:M:166:ILE:H	1.83	0.43
2:L:33:LEU:HD13	2:L:71:PHE:CG	2.54	0.43
1:H:48:MET:HG2	1:H:64:LEU:HD11	2.01	0.42
3:M:68:ASN:CB	3:M:87:GLY:HA2	2.45	0.42
2:L:155:GLN:HG3	2:L:158:ASN:HD21	1.84	0.42
3:M:205:TYR:HB3	3:M:209:HIS:O	2.20	0.42
3:M:121:VAL:O	3:M:130:LEU:HD12	2.19	0.42
2:L:89:GLN:HG2	2:L:90:GLN:N	2.34	0.42
3:M:540:TRP:NE1	3:M:542:HIS:O	2.52	0.42
2:L:134:CYS:HB2	2:L:148:TRP:CH2	2.55	0.42
3:M:328:GLN:O	3:M:331:ARG:HG2	2.19	0.42
2:L:146:VAL:HG22	2:L:196:VAL:HG22	2.01	0.42
3:M:222:THR:OG1	3:M:224:ASP:OD1	2.29	0.42
2:L:37:GLN:HB2	2:L:47:LEU:HD11	2.02	0.41
3:M:247:ILE:HG23	3:M:265:GLN:HG2	2.01	0.41
3:M:438:LEU:HD23	3:M:458:SER:HB2	2.01	0.41
1:H:71:THR:OG1	1:H:80:TYR:HB2	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:M:433:GLN:HG2	3:M:473:SER:O	2.21	0.41
3:M:149:ASN:OD1	6:M:606:NAG:N2	2.53	0.41
1:H:128:PHE:CZ	2:L:124:GLN:HB3	2.55	0.40
2:L:20:THR:HA	2:L:73:LEU:O	2.21	0.40
1:H:98:ARG:O	1:H:106:PHE:HA	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	211/228 (92%)	203 (96%)	8 (4%)	0	100	100
2	L	211/214 (99%)	200 (95%)	11 (5%)	0	100	100
3	M	513/534 (96%)	488 (95%)	24 (5%)	1 (0%)	47	79
All	All	935/976 (96%)	891 (95%)	43 (5%)	1 (0%)	51	83

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	M	542	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	169/194 (87%)	168 (99%)	1 (1%)	86	94
2	L	160/186 (86%)	160 (100%)	0	100	100
3	M	399/484 (82%)	395 (99%)	4 (1%)	76	90
All	All	728/864 (84%)	723 (99%)	5 (1%)	84	93

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	96	CYS
3	M	206	PHE
3	M	282	CYS
3	M	288	LEU
3	M	526	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
3	M	332	GLN
3	M	484	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

10 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	A	1	3,4	14,14,15	0.65	1 (7%)	17,19,21	0.58	0
4	NAG	A	2	4	14,14,15	0.32	0	17,19,21	0.40	0
5	NAG	B	1	3,5	14,14,15	0.80	1 (7%)	17,19,21	0.72	0
5	NAG	B	2	5	14,14,15	0.29	0	17,19,21	0.40	0
5	FUC	B	3	5	10,10,11	0.92	1 (10%)	14,14,16	1.75	3 (21%)
4	NAG	C	1	3,4	14,14,15	1.00	1 (7%)	17,19,21	0.79	0
4	NAG	C	2	4	14,14,15	0.56	0	17,19,21	1.28	1 (5%)
5	NAG	D	1	3,5	14,14,15	1.33	2 (14%)	17,19,21	1.42	2 (11%)
5	NAG	D	2	5	14,14,15	0.38	0	17,19,21	0.86	1 (5%)
5	FUC	D	3	5	10,10,11	0.82	0	14,14,16	1.71	3 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	1	3,4	-	0/6/23/26	0/1/1/1
4	NAG	A	2	4	-	0/6/23/26	0/1/1/1
5	NAG	B	1	3,5	-	4/6/23/26	0/1/1/1
5	NAG	B	2	5	-	0/6/23/26	0/1/1/1
5	FUC	B	3	5	-	-	0/1/1/1
4	NAG	C	1	3,4	-	4/6/23/26	0/1/1/1
4	NAG	C	2	4	-	3/6/23/26	0/1/1/1
5	NAG	D	1	3,5	-	3/6/23/26	0/1/1/1
5	NAG	D	2	5	-	2/6/23/26	0/1/1/1
5	FUC	D	3	5	-	-	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	D	1	NAG	O5-C1	-3.96	1.37	1.43
4	C	1	NAG	O5-C1	-3.57	1.38	1.43
5	B	1	NAG	O5-C1	-2.93	1.39	1.43
4	A	1	NAG	O5-C1	-2.33	1.40	1.43
5	D	1	NAG	C1-C2	2.31	1.55	1.52
5	B	3	FUC	O5-C5	2.07	1.48	1.43

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	2	NAG	C2-N2-C7	4.33	129.07	122.90
5	D	1	NAG	C2-N2-C7	3.99	128.59	122.90
5	B	3	FUC	C1-O5-C5	3.94	121.70	112.78
5	D	3	FUC	C1-O5-C5	3.55	120.82	112.78
5	D	3	FUC	O5-C1-C2	3.43	116.06	110.77
5	B	3	FUC	O5-C5-C4	3.14	115.15	109.52
5	B	3	FUC	O5-C1-C2	2.97	115.36	110.77
5	D	2	NAG	C1-O5-C5	2.75	115.92	112.19
5	D	3	FUC	O5-C5-C4	2.70	114.37	109.52
5	D	1	NAG	O4-C4-C5	-2.54	103.00	109.30

There are no chirality outliers.

All (16) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	C	1	NAG	C4-C5-C6-O6
5	B	1	NAG	O5-C5-C6-O6
5	D	2	NAG	O5-C5-C6-O6
4	C	2	NAG	C8-C7-N2-C2
4	C	2	NAG	O7-C7-N2-C2
5	B	1	NAG	C8-C7-N2-C2
5	B	1	NAG	O7-C7-N2-C2
5	D	1	NAG	C8-C7-N2-C2
5	D	1	NAG	O7-C7-N2-C2
4	C	1	NAG	O5-C5-C6-O6
4	C	1	NAG	C1-C2-N2-C7
5	B	1	NAG	C4-C5-C6-O6
5	D	2	NAG	C4-C5-C6-O6
4	C	1	NAG	C3-C2-N2-C7
5	D	1	NAG	C3-C2-N2-C7
4	C	2	NAG	C3-C2-N2-C7

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	2	NAG	1	0
4	C	2	NAG	1	0
5	D	1	NAG	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	NAG	M	606	3	14,14,15	0.28	0	17,19,21	0.53	0
7	FMT	M	612	-	0,2,2	0.00	-	0,1,1	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	M	606	3	-	4/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	M	606	NAG	O5-C5-C6-O6
6	M	606	NAG	C8-C7-N2-C2
6	M	606	NAG	O7-C7-N2-C2
6	M	606	NAG	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	M	606	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	H	215/228 (94%)	-0.36	0 100 100	59, 91, 117, 187	0
2	L	213/214 (99%)	-0.11	3 (1%) 75 56	65, 95, 147, 161	0
3	M	518/534 (97%)	-0.14	10 (1%) 66 46	53, 93, 154, 205	0
All	All	946/976 (96%)	-0.18	13 (1%) 75 56	53, 93, 147, 205	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	M	392	PRO	5.0
3	M	565	HIS	2.8
3	M	150	HIS	2.7
3	M	244	SER	2.5
2	L	191	VAL	2.5
2	L	151	ASP	2.2
3	M	352	ASP	2.2
3	M	241	PHE	2.1
2	L	181	LEU	2.1
3	M	206	PHE	2.1
3	M	153	ASP	2.0
3	M	391	GLY	2.0
3	M	357	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	FUC	B	3	10/11	0.53	0.61	179,185,190,190	0
5	FUC	D	3	10/11	0.57	0.50	131,147,154,156	0
5	NAG	D	2	14/15	0.59	0.50	188,205,220,234	0
5	NAG	B	2	14/15	0.63	0.46	154,174,179,179	0
5	NAG	D	1	14/15	0.66	0.24	110,150,160,178	0
4	NAG	C	2	14/15	0.76	0.66	203,211,213,213	0
5	NAG	B	1	14/15	0.79	0.22	132,152,170,176	0
4	NAG	A	2	14/15	0.80	0.45	158,185,194,196	0
4	NAG	C	1	14/15	0.84	0.46	135,171,180,192	0
4	NAG	A	1	14/15	0.91	0.23	101,134,155,170	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	NAG	M	606	14/15	0.58	0.54	157,169,181,183	0
7	FMT	M	612	3/3	0.90	0.17	81,81,88,91	0

6.5 Other polymers ⓘ

There are no such residues in this entry.