



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 29, 2020 – 01:38 am BST

PDB ID : 2WWR  
Title : Crystal Structure of Human Glyoxylate Reductase Hydroxypyruvate Reductase  
Authors : Booth, M.P.S.; Connors, R.; Rumsby, G.; Brady, R.L.  
Deposited on : 2009-10-26  
Resolution : 2.82 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3617 (2.84-2.80)
Clashscore	141614	4060 (2.84-2.80)
Ramachandran outliers	138981	3978 (2.84-2.80)
Sidechain outliers	138945	3980 (2.84-2.80)
RSRZ outliers	127900	3552 (2.84-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	330	<div> <div>3%</div> <div>84%</div> <div>11%</div> <div>• 5%</div> </div>
1	B	330	<div> <div>%</div> <div>82%</div> <div>13%</div> <div>• 5%</div> </div>
1	C	330	<div> <div>3%</div> <div>81%</div> <div>13%</div> <div>• 5%</div> </div>
1	D	330	<div> <div>5%</div> <div>78%</div> <div>15%</div> <div>•• 5%</div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 9640 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GLYOXYLATE REDUCTASE/HYDROXYPYRUVATE REDUCTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	315	Total	C	N	O	S	44	0	0
			2402	1518	426	446	12			
1	B	315	Total	C	N	O	S	36	0	0
			2402	1518	426	446	12			
1	C	315	Total	C	N	O	S	80	0	0
			2402	1518	426	446	12			
1	D	313	Total	C	N	O	S	170	0	0
			2385	1506	423	444	12			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	ALA	-	expression tag	UNP Q9UBQ7
A	0	SER	-	expression tag	UNP Q9UBQ7
B	-1	ALA	-	expression tag	UNP Q9UBQ7
B	0	SER	-	expression tag	UNP Q9UBQ7
C	-1	ALA	-	expression tag	UNP Q9UBQ7
C	0	SER	-	expression tag	UNP Q9UBQ7
D	-1	ALA	-	expression tag	UNP Q9UBQ7
D	0	SER	-	expression tag	UNP Q9UBQ7

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		
2	D	1	Total	Mg	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	18	Total 18	O 18	0	0
3	B	16	Total 16	O 16	0	0
3	C	6	Total 6	O 6	0	0
3	D	7	Total 7	O 7	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: GLYOXYLATE REDUCTASE/HYDROXYPYRUVATE REDUCTASE




- Molecule 1: GLYOXYLATE REDUCTASE/HYDROXYPYRUVATE REDUCTASE




- Molecule 1: GLYOXYLATE REDUCTASE/HYDROXYPYRUVATE REDUCTASE



- Molecule 1: GLYOXYLATE REDUCTASE/HYDROXYPYRUVATE REDUCTASE





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	67.36Å 116.08Å 198.23Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	100.00 – 2.82 45.58 – 2.82	Depositor EDS
% Data completeness (in resolution range)	90.6 (100.00-2.82) 90.7 (45.58-2.82)	Depositor EDS
$R_{merge}$	0.18	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.65 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.217 , 0.279 0.219 , 0.277	Depositor DCC
$R_{free}$ test set	1737 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	50.8	Xtriage
Anisotropy	0.413	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 56.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	9640	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 46.18 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.1874e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.66	3/2444 (0.1%)	0.63	5/3316 (0.2%)
1	B	0.77	5/2444 (0.2%)	0.70	5/3316 (0.2%)
1	C	1.13	13/2444 (0.5%)	1.74	20/3316 (0.6%)
1	D	1.63	19/2427 (0.8%)	1.40	35/3294 (1.1%)
All	All	1.11	40/9759 (0.4%)	1.21	65/13242 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	C	0	2
1	D	1	4
All	All	1	7

The worst 5 of 40 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	245	ARG	CD-NE	-46.78	0.67	1.46
1	B	66	ARG	CB-CG	-25.64	0.83	1.52
1	D	66	ARG	CA-C	24.10	2.15	1.52
1	A	178	GLN	CG-CD	-21.20	1.02	1.51
1	C	185	ARG	CB-CG	-20.23	0.97	1.52

The worst 5 of 65 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	188	ARG	NE-CZ-NH1	-54.32	93.14	120.30

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	188	ARG	NE-CZ-NH2	46.25	143.42	120.30
1	D	95	LYS	CA-C-O	-33.93	48.84	120.10
1	C	66	ARG	NE-CZ-NH1	-33.09	103.75	120.30
1	C	66	ARG	NE-CZ-NH2	32.45	136.52	120.30

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	D	95	LYS	CA

5 of 7 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	245	ARG	Sidechain
1	C	163	ARG	Sidechain
1	C	188	ARG	Sidechain
1	D	94	LYS	Peptide
1	D	95	LYS	Mainchain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2402	0	2455	14	0
1	B	2402	0	2455	25	0
1	C	2402	0	2455	15	0
1	D	2385	0	2426	18	0
2	B	1	0	0	0	0
2	D	1	0	0	0	0
3	A	18	0	0	0	0
3	B	16	0	0	0	0
3	C	6	0	0	0	0
3	D	7	0	0	0	0
All	All	9640	0	9791	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 64 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:27:ALA:HA	1:B:29:CYS:H	1.32	0.94
1:B:27:ALA:HA	1:B:29:CYS:N	2.01	0.76
1:D:10:PHE:HD1	1:D:55:LEU:CD1	2.03	0.71
1:C:85:ILE:HD13	1:C:326:LEU:HD22	1.79	0.64
1:B:27:ALA:CA	1:B:29:CYS:H	2.09	0.62

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	311/330 (94%)	299 (96%)	11 (4%)	1 (0%)	41	70
1	B	311/330 (94%)	295 (95%)	15 (5%)	1 (0%)	41	70
1	C	311/330 (94%)	296 (95%)	13 (4%)	2 (1%)	25	54
1	D	309/330 (94%)	290 (94%)	15 (5%)	4 (1%)	12	34
All	All	1242/1320 (94%)	1180 (95%)	54 (4%)	8 (1%)	25	54

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	95	LYS
1	D	244	SER
1	A	244	SER
1	C	106	VAL
1	D	65	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	257/270 (95%)	245 (95%)	12 (5%)	26	57
1	B	257/270 (95%)	249 (97%)	8 (3%)	40	72
1	C	257/270 (95%)	250 (97%)	7 (3%)	44	77
1	D	255/270 (94%)	248 (97%)	7 (3%)	44	77
All	All	1026/1080 (95%)	992 (97%)	34 (3%)	38	70

5 of 34 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	156	VAL
1	B	286	LYS
1	D	121	THR
1	B	216	CYS
1	A	109	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	166	GLN
1	B	33	GLN
1	C	74	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	D	4

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	D	46:GLU	C	47:ARG	N	1.77
1	D	91:ASP	C	92:GLU	N	1.71
1	D	73:ALA	C	74:ASN	N	1.15
1	D	63:VAL	C	64:ASP	N	1.10

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	315/330 (95%)	0.01	10 (3%) 47 37	22, 31, 44, 46	30 (9%)
1	B	315/330 (95%)	0.03	4 (1%) 77 72	20, 29, 51, 54	30 (9%)
1	C	315/330 (95%)	0.32	10 (3%) 47 37	34, 44, 61, 65	57 (18%)
1	D	303/330 (91%)	0.38	15 (4%) 28 20	25, 40, 53, 57	61 (20%)
All	All	1248/1320 (94%)	0.18	39 (3%) 49 39	20, 36, 54, 65	178 (14%)

The worst 5 of 39 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	29	CYS	3.9
1	C	6	LEU	3.7
1	D	78	ILE	3.4
1	D	67	ILE	3.3
1	D	98	ILE	3.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MG	B	1328	1/1	0.89	0.29	54,54,54,54	0
2	MG	D	1328	1/1	0.91	0.37	50,50,50,50	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.