



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 22, 2020 – 10:30 AM BST

PDB ID : 5WWV  
Title : Crystal structure of porcine kidney D-amino acid oxidase mutant (I230A/R283G)  
Authors : Motojima, F.; Yasukawa, K.; Ohno, A.; Asano, Y.  
Deposited on : 2017-01-05  
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1133 (3.20-3.20)
Clashscore	141614	1253 (3.20-3.20)
Ramachandran outliers	138981	1234 (3.20-3.20)
Sidechain outliers	138945	1233 (3.20-3.20)
RSRZ outliers	127900	1095 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	347	<div> <div>86%</div> <div>11%</div> <div>..</div> </div>
1	B	347	<div> <div>83%</div> <div>12%</div> <div>..</div> </div>
1	C	347	<div> <div>86%</div> <div>12%</div> <div>.</div> </div>
1	D	347	<div> <div>86%</div> <div>11%</div> <div>..</div> </div>
1	E	347	<div> <div>83%</div> <div>13%</div> <div>..</div> </div>
1	F	347	<div> <div>2%</div> <div>84%</div> <div>14%</div> <div>.</div> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	G	347	<div><div>%</div><div><div></div><div>86%</div><div>9%</div><div></div></div><div></div></div>
1	H	347	<div><div>%</div><div><div></div><div>81%</div><div>15%</div><div></div></div><div></div></div>

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 22427 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called D-amino-acid oxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	339	Total	C	N	O	S	0	1	0
			2720	1748	473	490	9			
1	B	339	Total	C	N	O	S	0	0	0
			2710	1742	470	489	9			
1	C	339	Total	C	N	O	S	0	0	0
			2710	1742	470	489	9			
1	D	339	Total	C	N	O	S	0	1	0
			2719	1750	470	490	9			
1	E	338	Total	C	N	O	S	0	2	0
			2718	1749	471	489	9			
1	F	338	Total	C	N	O	S	0	0	0
			2702	1736	469	488	9			
1	G	338	Total	C	N	O	S	0	0	0
			2702	1736	469	488	9			
1	H	337	Total	C	N	O	S	0	1	0
			2704	1738	470	487	9			

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	230	ALA	ILE	engineered mutation	UNP P00371
A	283	GLY	ARG	engineered mutation	UNP P00371
B	230	ALA	ILE	engineered mutation	UNP P00371
B	283	GLY	ARG	engineered mutation	UNP P00371
C	230	ALA	ILE	engineered mutation	UNP P00371
C	283	GLY	ARG	engineered mutation	UNP P00371
D	230	ALA	ILE	engineered mutation	UNP P00371
D	283	GLY	ARG	engineered mutation	UNP P00371
E	230	ALA	ILE	engineered mutation	UNP P00371
E	283	GLY	ARG	engineered mutation	UNP P00371
F	230	ALA	ILE	engineered mutation	UNP P00371
F	283	GLY	ARG	engineered mutation	UNP P00371
G	230	ALA	ILE	engineered mutation	UNP P00371


*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	283	GLY	ARG	engineered mutation	UNP P00371
H	230	ALA	ILE	engineered mutation	UNP P00371
H	283	GLY	ARG	engineered mutation	UNP P00371

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: C<sub>27</sub>H<sub>33</sub>N<sub>9</sub>O<sub>15</sub>P<sub>2</sub>).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	Cl	N	0	0
			15	13	1	1		
3	C	1	Total	C	Cl	N	0	0
			15	13	1	1		
3	D	1	Total	C	Cl	N	0	0
			15	13	1	1		
3	F	1	Total	C	Cl	N	0	0
			15	13	1	1		
3	G	1	Total	C	Cl	N	0	0
			15	13	1	1		

- Molecule 4 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			8	6	2		
4	A	1	Total	C	O	0	0
			8	6	2		
4	A	1	Total	C	O	0	0
			8	6	2		
4	A	1	Total	C	O	0	0
			8	6	2		
4	B	1	Total	C	O	0	0
			8	6	2		
4	B	1	Total	C	O	0	0
			8	6	2		
4	C	1	Total	C	O	0	0
			8	6	2		
4	D	1	Total	C	O	0	0
			8	6	2		
4	D	1	Total	C	O	0	0
			8	6	2		
4	E	1	Total	C	O	0	0
			8	6	2		
4	E	1	Total	C	O	0	0
			8	6	2		
4	E	1	Total	C	O	0	0
			8	6	2		
4	E	1	Total	C	O	0	0
			8	6	2		
4	F	1	Total	C	O	0	0
			8	6	2		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	F	1	Total	C	O	0	0
			8	6	2		
4	F	1	Total	C	O	0	0
			8	6	2		
4	G	1	Total	C	O	0	0
			8	6	2		
4	G	1	Total	C	O	0	0
			8	6	2		
4	H	1	Total	C	O	0	0
			8	6	2		
4	H	1	Total	C	O	0	0
			8	6	2		
4	H	1	Total	C	O	0	0
			8	6	2		

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	C	1	Total	O	S	0	0
			5	4	1		
5	D	1	Total	O	S	0	0
			5	4	1		
5	E	1	Total	O	S	0	0
			5	4	1		

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	F	1	Total	O	S	0	0
			5	4	1		
5	H	1	Total	O	S	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	7	Total	O	0	0
			7	7		
6	B	8	Total	O	0	0
			8	8		
6	C	9	Total	O	0	0
			9	9		
6	D	9	Total	O	0	0
			9	9		
6	E	5	Total	O	0	0
			5	5		
6	F	2	Total	O	0	0
			2	2		
6	H	5	Total	O	0	0
			5	5		

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: D-amino-acid oxidase






- Molecule 1: D-amino-acid oxidase

Chain E: 83% 13% . .




- Molecule 1: D-amino-acid oxidase

Chain F: 84% 14% 2% .





- Molecule 1: D-amino-acid oxidase

Chain G: 86% 9% . .



- Molecule 1: D-amino-acid oxidase

Chain H: 81% 15% . .



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	170.06Å 273.95Å 136.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.73 – 3.20 48.73 – 3.20	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.73-3.20) 100.0 (48.73-3.20)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	0.12	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.95 (at 3.19Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, $R_{free}$	0.145 , 0.242 0.152 , 0.241	Depositor DCC
$R_{free}$ test set	2657 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	79.1	Xtriage
Anisotropy	0.023	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 45.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	22427	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	79.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.02% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: 7V3, MPD, SO4, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.70	1/2797 (0.0%)	0.87	3/3810 (0.1%)
1	B	0.67	0/2786	0.84	1/3795 (0.0%)
1	C	0.64	1/2786 (0.0%)	0.83	2/3795 (0.1%)
1	D	0.65	0/2799	0.84	3/3813 (0.1%)
1	E	0.60	0/2802	0.82	4/3817 (0.1%)
1	F	0.64	0/2778	0.81	0/3784
1	G	0.59	0/2778	0.79	0/3784
1	H	0.61	1/2781 (0.0%)	0.77	2/3788 (0.1%)
All	All	0.64	3/22307 (0.0%)	0.82	15/30386 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	247	TRP	CB-CG	6.38	1.61	1.50
1	C	247	TRP	CB-CG	5.80	1.60	1.50
1	A	147	TRP	CB-CG	5.24	1.59	1.50

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	120	ARG	NE-CZ-NH1	6.94	123.77	120.30
1	H	120	ARG	NE-CZ-NH1	6.47	123.54	120.30
1	A	120	ARG	NE-CZ-NH1	6.29	123.45	120.30
1	E	151	ARG	NE-CZ-NH1	6.16	123.38	120.30
1	B	221	ARG	NE-CZ-NH1	5.81	123.20	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2720	0	2665	10	0
1	B	2710	0	2659	19	0
1	C	2710	0	2659	14	0
1	D	2719	0	2668	11	0
1	E	2718	0	2664	16	0
1	F	2702	0	2648	17	0
1	G	2702	0	2648	16	0
1	H	2704	0	2648	23	0
2	A	53	0	31	0	0
2	B	53	0	31	0	0
2	C	53	0	31	0	0
2	D	53	0	31	1	0
2	E	53	0	31	1	0
2	F	53	0	31	1	0
2	G	53	0	31	1	0
2	H	53	0	31	0	0
3	A	15	0	0	0	0
3	C	15	0	0	2	0
3	D	15	0	0	1	0
3	F	15	0	0	1	0
3	G	15	0	0	0	0
4	A	32	0	56	0	0
4	B	16	0	28	0	0
4	C	8	0	14	1	0
4	D	16	0	28	0	0
4	E	32	0	56	2	0
4	F	24	0	42	0	0
4	G	16	0	28	0	0
4	H	24	0	42	3	0
5	A	5	0	0	0	0
5	C	5	0	0	0	0
5	D	5	0	0	0	0
5	E	5	0	0	0	0
5	F	5	0	0	1	0
5	H	5	0	0	0	0
6	A	7	0	0	0	0
6	B	8	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	C	9	0	0	1	0
6	D	9	0	0	0	0
6	E	5	0	0	0	0
6	F	2	0	0	0	0
6	H	5	0	0	0	0
All	All	22427	0	21801	121	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 121 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:E:404:MPD:H52	4:E:404:MPD:H12	1.71	0.73
1:H:334:LEU:HA	1:H:335:GLU:C	2.13	0.69
1:D:96:ASN:ND2	1:D:217:HIS:NE2	2.38	0.69
1:A:120:ARG:NH2	1:B:110:MET:O	2.28	0.66
1:H:330:PHE:O	1:H:333:VAL:HG22	2.00	0.62

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	338/347 (97%)	310 (92%)	24 (7%)	4 (1%)	13	49
1	B	337/347 (97%)	308 (91%)	27 (8%)	2 (1%)	25	64
1	C	337/347 (97%)	314 (93%)	23 (7%)	0	100	100
1	D	338/347 (97%)	317 (94%)	20 (6%)	1 (0%)	41	74
1	E	338/347 (97%)	308 (91%)	29 (9%)	1 (0%)	41	74

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	F	336/347 (97%)	299 (89%)	34 (10%)	3 (1%)	17	56
1	G	336/347 (97%)	304 (90%)	31 (9%)	1 (0%)	41	74
1	H	336/347 (97%)	309 (92%)	25 (7%)	2 (1%)	25	64
All	All	2696/2776 (97%)	2469 (92%)	213 (8%)	14 (0%)	29	67

5 of 14 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	333	VAL
1	D	338	ASN
1	B	56	THR
1	F	29	PRO
1	G	29	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	289/296 (98%)	268 (93%)	21 (7%)	14	46
1	B	288/296 (97%)	265 (92%)	23 (8%)	12	42
1	C	288/296 (97%)	271 (94%)	17 (6%)	19	54
1	D	289/296 (98%)	267 (92%)	22 (8%)	13	45
1	E	289/296 (98%)	267 (92%)	22 (8%)	13	45
1	F	287/296 (97%)	266 (93%)	21 (7%)	14	46
1	G	287/296 (97%)	266 (93%)	21 (7%)	14	46
1	H	287/296 (97%)	265 (92%)	22 (8%)	13	44
All	All	2304/2368 (97%)	2135 (93%)	169 (7%)	13	46

5 of 169 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	218	ASP

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
1	E	221	ARG
1	H	202	ILE
1	D	291	LEU
1	E	81	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 32 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	96	ASN
1	E	190	GLN
1	H	96	ASN
1	E	134	ASN
1	F	20	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

40 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	MPD	G	403	-	7,7,7	0.58	0	9,10,10	0.65	0
4	MPD	H	402	-	7,7,7	0.63	0	9,10,10	0.66	0
4	MPD	C	403	-	7,7,7	0.51	0	9,10,10	0.86	0
4	MPD	A	404	-	7,7,7	0.45	0	9,10,10	1.36	1 (11%)
3	7V3	C	402	-	15,16,16	2.72	4 (26%)	15,21,21	0.96	1 (6%)
4	MPD	D	403	-	7,7,7	0.52	0	9,10,10	1.26	1 (11%)
4	MPD	F	403	-	7,7,7	0.29	0	9,10,10	0.29	0
2	FAD	C	401	-	51,58,58	1.76	9 (17%)	60,89,89	1.63	8 (13%)
5	SO4	A	407	-	4,4,4	0.30	0	6,6,6	0.52	0
4	MPD	B	402	-	7,7,7	0.50	0	9,10,10	1.10	0
3	7V3	A	402	-	15,16,16	1.82	2 (13%)	15,21,21	0.95	1 (6%)
5	SO4	F	406	-	4,4,4	0.39	0	6,6,6	0.38	0
5	SO4	E	405	-	4,4,4	0.31	0	6,6,6	0.36	0
3	7V3	F	402	-	15,16,16	2.08	5 (33%)	15,21,21	0.97	1 (6%)
4	MPD	F	405	-	7,7,7	0.59	0	9,10,10	1.48	1 (11%)
2	FAD	H	401	-	51,58,58	1.76	8 (15%)	60,89,89	1.86	10 (16%)
4	MPD	A	405	-	7,7,7	0.52	0	9,10,10	0.70	0
4	MPD	B	403	-	7,7,7	0.37	0	9,10,10	0.52	0
5	SO4	H	405	-	4,4,4	0.36	0	6,6,6	0.16	0
4	MPD	E	406	-	7,7,7	0.36	0	9,10,10	0.46	0
4	MPD	A	403	-	7,7,7	0.33	0	9,10,10	0.77	0
3	7V3	D	402	-	15,16,16	1.66	3 (20%)	15,21,21	1.22	1 (6%)
4	MPD	F	404	-	7,7,7	0.53	0	9,10,10	0.74	0
4	MPD	D	404	-	7,7,7	0.41	0	9,10,10	0.76	0
2	FAD	F	401	-	51,58,58	1.87	7 (13%)	60,89,89	2.18	10 (16%)
2	FAD	A	401	-	51,58,58	1.79	6 (11%)	60,89,89	1.88	12 (20%)
2	FAD	G	401	-	51,58,58	1.92	8 (15%)	60,89,89	2.28	13 (21%)
4	MPD	A	406	-	7,7,7	0.44	0	9,10,10	0.57	0
4	MPD	E	403	-	7,7,7	0.51	0	9,10,10	0.77	0
4	MPD	H	403	-	7,7,7	0.40	0	9,10,10	0.89	0
3	7V3	G	402	-	15,16,16	1.47	4 (26%)	15,21,21	0.91	0
4	MPD	E	404	-	7,7,7	0.51	0	9,10,10	0.77	0
5	SO4	C	404	-	4,4,4	0.36	0	6,6,6	0.63	0
4	MPD	G	404	-	7,7,7	0.48	0	9,10,10	1.03	0
4	MPD	H	404	-	7,7,7	0.46	0	9,10,10	1.55	2 (22%)
5	SO4	D	405	-	4,4,4	0.44	0	6,6,6	0.48	0
2	FAD	B	401	-	51,58,58	1.85	8 (15%)	60,89,89	1.86	10 (16%)
2	FAD	D	401	-	51,58,58	1.82	5 (9%)	60,89,89	1.90	13 (21%)
4	MPD	E	402	-	7,7,7	0.75	0	9,10,10	1.11	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	FAD	E	401	-	51,58,58	1.86	5 (9%)	60,89,89	2.10	14 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MPD	G	403	-	-	2/5/5/5	-
4	MPD	H	402	-	-	2/5/5/5	-
4	MPD	C	403	-	-	3/5/5/5	-
4	MPD	A	404	-	-	0/5/5/5	-
3	7V3	C	402	-	-	2/8/8/8	0/2/2/2
4	MPD	D	403	-	-	1/5/5/5	-
4	MPD	F	403	-	-	2/5/5/5	-
2	FAD	C	401	-	-	1/30/50/50	0/6/6/6
4	MPD	B	402	-	-	0/5/5/5	-
3	7V3	A	402	-	-	1/8/8/8	0/2/2/2
3	7V3	F	402	-	-	2/8/8/8	0/2/2/2
4	MPD	F	405	-	-	2/5/5/5	-
2	FAD	H	401	-	-	1/30/50/50	0/6/6/6
4	MPD	A	405	-	-	2/5/5/5	-
4	MPD	B	403	-	-	2/5/5/5	-
4	MPD	E	406	-	-	2/5/5/5	-
4	MPD	A	403	-	-	2/5/5/5	-
3	7V3	D	402	-	-	0/8/8/8	0/2/2/2
4	MPD	F	404	-	-	2/5/5/5	-
4	MPD	D	404	-	-	2/5/5/5	-
2	FAD	F	401	-	-	1/30/50/50	0/6/6/6
2	FAD	A	401	-	-	2/30/50/50	0/6/6/6
2	FAD	G	401	-	-	2/30/50/50	0/6/6/6
4	MPD	A	406	-	-	0/5/5/5	-
4	MPD	E	403	-	-	1/5/5/5	-
4	MPD	H	403	-	-	1/5/5/5	-
3	7V3	G	402	-	-	0/8/8/8	0/2/2/2
4	MPD	E	404	-	-	2/5/5/5	-
4	MPD	G	404	-	-	2/5/5/5	-
4	MPD	H	404	-	-	0/5/5/5	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	B	401	-	-	3/30/50/50	0/6/6/6
2	FAD	D	401	-	-	4/30/50/50	0/6/6/6
4	MPD	E	402	-	-	3/5/5/5	-
2	FAD	E	401	-	-	1/30/50/50	0/6/6/6

The worst 5 of 74 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	FAD	C4X-C10	9.77	1.48	1.38
2	F	401	FAD	C4X-C10	9.22	1.48	1.38
2	G	401	FAD	C4X-C10	9.15	1.48	1.38
2	A	401	FAD	C4X-C10	9.15	1.48	1.38
2	E	401	FAD	C4X-C10	9.05	1.47	1.38

The worst 5 of 99 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	FAD	C4-N3-C2	8.06	121.95	115.14
2	F	401	FAD	C4-N3-C2	8.00	121.90	115.14
2	F	401	FAD	C1'-N10-C9A	7.92	124.52	118.29
2	G	401	FAD	C1'-N10-C9A	7.84	124.46	118.29
2	G	401	FAD	C4-N3-C2	7.78	121.71	115.14

There are no chirality outliers.

5 of 53 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	H	402	MPD	C2-C3-C4-O4
4	C	403	MPD	C1-C2-C3-C4
4	C	403	MPD	O2-C2-C3-C4
4	A	403	MPD	C2-C3-C4-O4
4	E	403	MPD	C2-C3-C4-O4

There are no ring outliers.

12 monomers are involved in 14 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	403	MPD	1	0
3	C	402	7V3	2	0
5	F	406	SO4	1	0


*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	402	7V3	1	0
3	D	402	7V3	1	0
2	F	401	FAD	1	0
2	G	401	FAD	1	0
4	H	403	MPD	3	0
4	E	404	MPD	1	0
2	D	401	FAD	1	0
4	E	402	MPD	1	0
2	E	401	FAD	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.























## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	339/347 (97%)	-0.61	0 <span>100</span> <span>100</span>	38, 59, 111, 146	0
1	B	339/347 (97%)	-0.54	1 (0%) <span>94</span> <span>92</span>	39, 64, 125, 180	0
1	C	339/347 (97%)	-0.51	1 (0%) <span>94</span> <span>92</span>	42, 63, 111, 161	0
1	D	339/347 (97%)	-0.50	0 <span>100</span> <span>100</span>	43, 67, 120, 172	1 (0%)
1	E	338/347 (97%)	-0.42	0 <span>100</span> <span>100</span>	44, 78, 124, 157	0
1	F	338/347 (97%)	-0.34	6 (1%) <span>68</span> <span>55</span>	51, 81, 120, 152	0
1	G	338/347 (97%)	-0.29	2 (0%) <span>89</span> <span>83</span>	60, 86, 131, 149	0
1	H	337/347 (97%)	-0.30	2 (0%) <span>89</span> <span>83</span>	57, 87, 132, 194	0
All	All	2707/2776 (97%)	-0.44	12 (0%) <span>92</span> <span>89</span>	38, 75, 125, 194	1 (0%)

The worst 5 of 12 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	300	SER	4.5
1	H	300	SER	3.2
1	F	301	SER	2.8
1	F	223	ILE	2.8
1	F	60	SER	2.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	SO4	H	405	5/5	0.82	0.22	144,146,156,157	0
5	SO4	D	405	5/5	0.86	0.21	115,120,122,132	0
5	SO4	F	406	5/5	0.87	0.28	177,191,197,221	0
4	MPD	E	406	8/8	0.88	0.40	87,106,113,118	0
4	MPD	A	406	8/8	0.89	0.26	94,98,102,104	0
4	MPD	E	404	8/8	0.89	0.65	101,112,121,122	0
5	SO4	A	407	5/5	0.89	0.25	121,123,131,136	0
4	MPD	G	403	8/8	0.90	0.24	85,95,101,107	0
3	7V3	G	402	15/15	0.90	0.53	114,122,140,160	0
4	MPD	B	402	8/8	0.91	0.30	85,87,88,97	0
4	MPD	G	404	8/8	0.91	0.43	88,98,104,106	0
4	MPD	B	403	8/8	0.91	0.33	112,120,123,125	0
4	MPD	F	403	8/8	0.92	0.25	102,109,111,114	0
4	MPD	H	403	8/8	0.93	0.25	68,74,83,83	0
5	SO4	E	405	5/5	0.93	0.15	122,126,131,141	0
3	7V3	F	402	15/15	0.93	0.54	87,94,104,111	0
4	MPD	D	404	8/8	0.93	0.23	84,93,99,104	0
4	MPD	H	404	8/8	0.93	0.26	71,75,78,79	0
4	MPD	D	403	8/8	0.93	0.24	74,93,108,123	0
4	MPD	E	403	8/8	0.94	0.21	82,84,89,95	0
3	7V3	C	402	15/15	0.94	0.53	95,102,117,125	0
4	MPD	F	405	8/8	0.94	0.20	67,73,79,80	0
3	7V3	D	402	15/15	0.94	0.44	87,98,106,120	0
4	MPD	F	404	8/8	0.94	0.25	65,77,92,94	0
4	MPD	A	405	8/8	0.94	0.25	84,89,97,103	0
3	7V3	A	402	15/15	0.94	0.24	82,89,101,106	0
4	MPD	E	402	8/8	0.94	0.19	82,97,105,107	0
4	MPD	A	403	8/8	0.95	0.27	70,81,88,92	0
5	SO4	C	404	5/5	0.95	0.13	91,97,108,110	0
4	MPD	H	402	8/8	0.95	0.24	92,99,101,105	0
4	MPD	A	404	8/8	0.96	0.19	47,49,58,69	0
2	FAD	F	401	53/53	0.97	0.15	51,60,67,74	0
2	FAD	G	401	53/53	0.97	0.14	65,75,88,91	0
4	MPD	C	403	8/8	0.97	0.16	56,61,63,64	0
2	FAD	A	401	53/53	0.98	0.14	36,42,58,71	0
2	FAD	H	401	53/53	0.98	0.16	63,69,79,80	0
2	FAD	B	401	53/53	0.98	0.15	44,56,84,95	0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	FAD	D	401	53/53	0.98	0.15	43,55,61,65	0
2	FAD	C	401	53/53	0.98	0.15	40,48,56,64	0
2	FAD	E	401	53/53	0.98	0.14	45,58,69,78	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around FAD F 401:**




2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around FAD G 401:**

$2mF_o - DF_c$  (at 0.7 rmsd) in gray  
 $mF_o - DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around FAD A 401:**

$2mF_o - DF_c$  (at 0.7 rmsd) in gray  
 $mF_o - DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around FAD H 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around FAD B 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around FAD D 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around FAD C 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around FAD E 401:**

$2mF_o - DF_c$  (at 0.7 rmsd) in gray  
 $mF_o - DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.