



wwPDB X-ray Structure Validation Summary Report ⓘ

May 21, 2020 – 03:24 am BST

PDB ID : 3W XK
Title : Crystal structure of trypanosoma brucei gambiense glycerol kinase in complex with glycerol
Authors : Balogun, E.O.; Inaoka, D.K.; Shiba, T.; Kido, Y.; Tsuge, C.; Nara, T.; Aoki, T.; Honma, T.; Tanaka, A.; Inoue, M.; Matsuoka, S.; Michels, P.A.M.; Kita, K.; Harada, S.
Deposited on : 2014-08-01
Resolution : 2.37 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.37 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5509 (2.40-2.36)
Clashscore	141614	6082 (2.40-2.36)
Ramachandran outliers	138981	5973 (2.40-2.36)
Sidechain outliers	138945	5975 (2.40-2.36)
RSRZ outliers	127900	5397 (2.40-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	518	<div> <div>%</div> <div> <div></div> <div>74%</div> <div>21%</div> <div>• •</div> </div> </div>
1	B	518	<div> <div>2%</div> <div> <div></div> <div>75%</div> <div>21%</div> <div>• •</div> </div> </div>
1	C	518	<div> <div>2%</div> <div> <div></div> <div>74%</div> <div>23%</div> <div>• •</div> </div> </div>
1	D	518	<div> <div>3%</div> <div> <div></div> <div>75%</div> <div>20%</div> <div>• •</div> </div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 16111 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycerol kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	513	Total	C	N	O	S	0	0	0
			3957	2499	694	731	33			
1	B	513	Total	C	N	O	S	0	0	0
			3957	2499	694	731	33			
1	C	513	Total	C	N	O	S	0	0	0
			3957	2499	694	731	33			
1	D	513	Total	C	N	O	S	0	1	0
			3962	2502	695	732	33			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	GLY	-	EXPRESSION TAG	UNP D3KVM3
A	-4	ILE	-	EXPRESSION TAG	UNP D3KVM3
A	-3	ASP	-	EXPRESSION TAG	UNP D3KVM3
A	-2	PRO	-	EXPRESSION TAG	UNP D3KVM3
A	-1	PHE	-	EXPRESSION TAG	UNP D3KVM3
A	0	THR	-	EXPRESSION TAG	UNP D3KVM3
B	-5	GLY	-	EXPRESSION TAG	UNP D3KVM3
B	-4	ILE	-	EXPRESSION TAG	UNP D3KVM3
B	-3	ASP	-	EXPRESSION TAG	UNP D3KVM3
B	-2	PRO	-	EXPRESSION TAG	UNP D3KVM3
B	-1	PHE	-	EXPRESSION TAG	UNP D3KVM3
B	0	THR	-	EXPRESSION TAG	UNP D3KVM3
C	-5	GLY	-	EXPRESSION TAG	UNP D3KVM3
C	-4	ILE	-	EXPRESSION TAG	UNP D3KVM3
C	-3	ASP	-	EXPRESSION TAG	UNP D3KVM3
C	-2	PRO	-	EXPRESSION TAG	UNP D3KVM3
C	-1	PHE	-	EXPRESSION TAG	UNP D3KVM3
C	0	THR	-	EXPRESSION TAG	UNP D3KVM3
D	-5	GLY	-	EXPRESSION TAG	UNP D3KVM3
D	-4	ILE	-	EXPRESSION TAG	UNP D3KVM3
D	-3	ASP	-	EXPRESSION TAG	UNP D3KVM3

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
D	-2	PRO	-	EXPRESSION TAG	UNP D3KVM3
D	-1	PHE	-	EXPRESSION TAG	UNP D3KVM3
D	0	THR	-	EXPRESSION TAG	UNP D3KVM3

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	C	1	Total C O 6 3 3	0	0
2	C	1	Total C O 6 3 3	0	0
2	D	1	Total C O 6 3 3	0	0
2	D	1	Total C O 6 3 3	0	0

- Molecule 3 is HEXANE-1,6-DIOL (three-letter code: HEZ) (formula: $C_6H_{14}O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			8	6	2		
3	B	1	Total	C	O	0	0
			8	6	2		
3	C	1	Total	C	O	0	0
			8	6	2		
3	D	1	Total	C	O	0	0
			8	6	2		

- Molecule 4 is water.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	52	Total	O	0	0
			52	52		
4	B	52	Total	O	0	0
			52	52		
4	C	50	Total	O	0	0
			50	50		
4	D	50	Total	O	0	0
			50	50		

- Molecule 1: Glycerol kinase





• Molecule 1: Glycerol kinase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	62.48 Å 122.14 Å 154.51 Å 90.00° 90.76° 90.00°	Depositor
Resolution (Å)	30.00 – 2.37 47.91 – 2.37	Depositor EDS
% Data completeness (in resolution range)	90.3 (30.00-2.37) 93.5 (47.91-2.37)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.25 (at 2.37 Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.190 , 0.251 0.207 , 0.264	Depositor DCC
R_{free} test set	4415 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	55.2	Xtriage
Anisotropy	0.033	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 46.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.37$	Xtriage
Estimated twinning fraction	0.008 for h,-k,-l	Xtriage
Reported twinning fraction	0.914 for H, K, L 0.086 for -h,-k,l	Depositor
Outliers	0 of 88201 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	16111	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, HEZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.93	2/4039 (0.0%)	0.82	2/5465 (0.0%)
1	B	0.94	0/4039	0.83	2/5465 (0.0%)
1	C	1.04	3/4039 (0.1%)	0.80	1/5465 (0.0%)
1	D	0.88	2/4047 (0.0%)	0.79	0/5476
All	All	0.95	7/16164 (0.0%)	0.81	5/21871 (0.0%)

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	309	CYS	CB-SG	-5.86	1.72	1.81
1	A	311	TYR	CD1-CE1	-5.50	1.31	1.39
1	C	310	TYR	CD2-CE2	-5.41	1.31	1.39
1	D	279	PHE	CD1-CE1	-5.25	1.28	1.39
1	A	311	TYR	CD2-CE2	-5.23	1.31	1.39

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	46	ASP	CB-CG-OD2	5.66	123.39	118.30
1	A	345	ARG	NE-CZ-NH2	-5.44	117.58	120.30
1	A	98	LEU	CA-CB-CG	5.35	127.61	115.30
1	C	242	LEU	N-CA-C	-5.24	96.86	111.00
1	B	384	ARG	NE-CZ-NH1	5.14	122.87	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3957	0	3970	99	0
1	B	3957	0	3970	81	0
1	C	3957	0	3970	95	0
1	D	3962	0	3976	107	0
2	A	12	0	16	1	0
2	B	6	0	8	0	0
2	C	12	0	16	2	0
2	D	12	0	16	1	0
3	B	16	0	28	4	0
3	C	8	0	14	0	0
3	D	8	0	14	0	0
4	A	52	0	0	7	0
4	B	52	0	0	3	0
4	C	50	0	0	6	0
4	D	50	0	0	2	0
All	All	16111	0	15998	383	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 383 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:364:ALA:CB	1:C:365:PRO:HD2	1.63	1.28
1:B:364:ALA:HB1	1:B:365:PRO:CD	1.64	1.27
1:C:364:ALA:HB1	1:C:365:PRO:CD	1.64	1.27
1:B:420:ASP:OD1	1:B:449:THR:HG21	1.40	1.21
1:B:364:ALA:CB	1:B:365:PRO:HD2	1.70	1.18

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	511/518 (99%)	485 (95%)	25 (5%)	1 (0%)	47	61
1	B	511/518 (99%)	480 (94%)	27 (5%)	4 (1%)	19	27
1	C	511/518 (99%)	492 (96%)	17 (3%)	2 (0%)	34	46
1	D	512/518 (99%)	476 (93%)	33 (6%)	3 (1%)	25	34
All	All	2045/2072 (99%)	1933 (94%)	102 (5%)	10 (0%)	29	39

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	364	ALA
1	B	364	ALA
1	C	364	ALA
1	D	364	ALA
1	B	215	MET

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	421/425 (99%)	379 (90%)	42 (10%)	7	10
1	B	421/425 (99%)	383 (91%)	38 (9%)	9	12
1	C	421/425 (99%)	385 (91%)	36 (9%)	10	14
1	D	422/425 (99%)	382 (90%)	40 (10%)	8	11
All	All	1685/1700 (99%)	1529 (91%)	156 (9%)	9	11

5 of 156 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	317	ILE
1	C	106	ASP
1	D	366	TYR
1	B	329	ARG
1	B	476	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 31 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	23	GLN
1	C	105	ASN
1	D	243	ASN
1	C	35	GLN
1	C	224	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	D	602	-	5,5,5	0.28	0	5,5,5	0.63	0
3	HEZ	B	602	-	7,7,7	0.49	0	6,6,6	0.39	0
3	HEZ	C	603	-	7,7,7	0.52	0	6,6,6	0.48	0
2	GOL	C	602	-	5,5,5	0.44	0	5,5,5	0.45	0
2	GOL	A	602	-	5,5,5	0.46	0	5,5,5	0.33	0
3	HEZ	D	603	-	7,7,7	0.50	0	6,6,6	0.31	0
2	GOL	A	601	-	5,5,5	0.50	0	5,5,5	0.43	0
2	GOL	C	601	-	5,5,5	0.33	0	5,5,5	0.31	0
3	HEZ	B	603	-	7,7,7	0.51	0	6,6,6	0.29	0
2	GOL	B	601	-	5,5,5	0.57	0	5,5,5	0.90	0
2	GOL	D	601	-	5,5,5	0.34	0	5,5,5	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	D	602	-	-	0/4/4/4	-
3	HEZ	B	602	-	-	4/5/5/5	-
3	HEZ	C	603	-	-	3/5/5/5	-
2	GOL	C	602	-	-	4/4/4/4	-
2	GOL	A	602	-	-	4/4/4/4	-
3	HEZ	D	603	-	-	5/5/5/5	-
2	GOL	A	601	-	-	2/4/4/4	-
2	GOL	C	601	-	-	0/4/4/4	-
3	HEZ	B	603	-	-	4/5/5/5	-
2	GOL	B	601	-	-	4/4/4/4	-
2	GOL	D	601	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 30 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	602	GOL	O1-C1-C2-C3
2	C	602	GOL	C1-C2-C3-O3
2	A	602	GOL	O1-C1-C2-C3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	602	GOL	C1-C2-C3-O3
2	B	601	GOL	C1-C2-C3-O3

There are no ring outliers.

5 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	602	GOL	1	0
3	B	602	HEZ	3	0
2	A	602	GOL	1	0
2	C	601	GOL	2	0
3	B	603	HEZ	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	513/518 (99%)	0.17	3 (0%) 89 89	34, 49, 63, 72	1 (0%)
1	B	513/518 (99%)	0.17	10 (1%) 66 68	30, 53, 69, 87	1 (0%)
1	C	513/518 (99%)	0.19	11 (2%) 63 65	32, 51, 68, 75	1 (0%)
1	D	513/518 (99%)	0.13	13 (2%) 57 59	31, 51, 69, 84	1 (0%)
All	All	2052/2072 (99%)	0.17	37 (1%) 68 70	30, 51, 67, 87	4 (0%)

The worst 5 of 37 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	491	MET	6.4
1	B	491	MET	5.3
1	A	278	CYS	4.4
1	B	152	VAL	4.2
1	D	511	LYS	3.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	HEZ	B	602	8/8	0.73	0.29	63,66,67,67	0
3	HEZ	B	603	8/8	0.75	0.28	58,62,63,64	0
3	HEZ	D	603	8/8	0.78	0.26	52,56,57,58	0
3	HEZ	C	603	8/8	0.85	0.18	49,51,52,53	0
2	GOL	A	602	6/6	0.86	0.21	58,60,63,65	0
2	GOL	D	602	6/6	0.87	0.19	60,61,61,62	0
2	GOL	C	602	6/6	0.88	0.27	50,58,61,65	0
2	GOL	C	601	6/6	0.89	0.25	72,73,74,75	0
2	GOL	A	601	6/6	0.94	0.14	58,59,60,63	0
2	GOL	B	601	6/6	0.96	0.14	31,34,36,37	0
2	GOL	D	601	6/6	0.96	0.13	38,40,43,45	0

6.5 Other polymers [i](#)

There are no such residues in this entry.