



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 9, 2020 – 07:27 AM BST

PDB ID : 3WY4
Title : Crystal structure of alpha-glucosidase mutant E271Q in complex with maltose
Authors : Shen, X.; Gai, Z.; Kato, K.; Yao, M.
Deposited on : 2014-08-18
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	538	<div> <div style="width: 100%; height: 10px; background-color: red;"></div> <div style="width: 100%; height: 10px; background-color: green; position: relative;"> % 88% 11% • </div> </div>
1	B	538	<div> <div style="width: 9%; height: 10px; background-color: red;"></div> <div style="width: 100%; height: 10px; background-color: green; position: relative;"> 9% 64% 12% • • 22% </div> </div>
2	C	2	<div> <div style="width: 100%; height: 10px; background-color: green; position: relative;"> 50% 50% </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	601	-	-	-	X
4	GOL	B	601	-	-	-	X

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8190 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha-glucosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	535	Total	C	N	O	S	0	0	0
			4292	2730	735	809	18			
1	B	421	Total	C	N	O	S	0	0	0
			3398	2172	578	634	14			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	271	GLN	GLU	engineered mutation	UNP H3K096
B	271	GLN	GLU	engineered mutation	UNP H3K096

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	C	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	293	Total	O	0	0
			293	293		
5	B	159	Total	O	0	0
			159	159		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Alpha-glucosidase



• Molecule 1: Alpha-glucosidase



• Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	111.46 Å 181.06 Å 51.93 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.46 – 2.50 47.46 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.6 (47.46-2.50) 99.6 (47.46-2.50)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.40 (at 2.51 Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.202 , 0.241 0.210 , 0.243	Depositor DCC
R_{free} test set	1865 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	37.1	Xtriage
Anisotropy	0.594	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 40.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8190	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.21	0/4423	0.39	0/6033
1	B	0.24	0/3496	0.49	0/4762
All	All	0.22	0/7919	0.43	0/10795

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	58	SER	Peptide
1	B	59	PRO	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4292	0	4053	40	0
1	B	3398	0	3223	58	0
2	C	23	0	21	0	0
3	A	1	0	0	0	0
4	A	18	0	24	3	0
4	B	6	0	8	2	0
5	A	293	0	0	11	0
5	B	159	0	0	11	0
All	All	8190	0	7329	97	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (97) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:44:SER:OG	5:B:800:HOH:O	1.81	0.96
1:B:480:THR:OG1	5:B:729:HOH:O	1.95	0.84
1:A:514:ALA:O	5:A:873:HOH:O	1.96	0.83
1:B:139:PRO:O	5:B:795:HOH:O	1.96	0.82
1:B:297:PHE:HA	1:B:298:ASP:HB2	1.61	0.80
1:A:396:GLU:OE1	5:A:934:HOH:O	2.03	0.77
1:B:305:SER:HB3	1:B:308:TYR:HB3	1.67	0.77
1:A:209:HIS:HB3	4:A:601:GOL:H32	1.66	0.76
1:B:499:GLN:OE1	5:B:731:HOH:O	2.05	0.74
1:B:297:PHE:HA	1:B:298:ASP:CB	2.16	0.74
1:B:179:GLU:OE1	5:B:777:HOH:O	2.06	0.73
1:A:311:GLU:OE1	5:A:827:HOH:O	2.07	0.72
1:A:315:ARG:NH1	5:A:962:HOH:O	2.19	0.71
1:A:289:ASP:OD2	5:A:888:HOH:O	2.09	0.70
1:A:328:ALA:O	5:A:802:HOH:O	2.07	0.70
1:A:25:ARG:NH2	1:A:33:ASN:OD1	2.25	0.70
1:B:19:ARG:HD3	1:B:372:GLU:HB3	1.73	0.69
1:B:447:ASN:HA	1:B:450:ARG:HD2	1.75	0.69
1:B:298:ASP:O	1:B:300:LEU:N	2.28	0.66
1:A:172:ASP:OD1	5:A:710:HOH:O	2.13	0.66
1:B:472:VAL:HG21	1:B:478:GLY:HA3	1.79	0.64
1:B:301:ASN:OD1	1:B:301:ASN:N	2.31	0.63
1:B:301:ASN:HB2	1:B:302:MET:HA	1.82	0.62
1:B:200:ARG:NH2	1:B:271:GLN:OE1	2.33	0.62
1:A:105:HIS:O	5:A:703:HOH:O	2.16	0.61
1:A:441:ASP:O	1:A:447:ASN:ND2	2.32	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:438:GLN:HG3	1:A:444:ALA:HB1	1.83	0.61
1:A:304:HIS:NE2	5:A:733:HOH:O	2.32	0.60
1:B:165:ASN:ND2	4:B:601:GOL:O3	2.34	0.60
1:B:360:SER:O	1:B:481:ARG:NH1	2.37	0.58
1:B:165:ASN:HD22	1:B:206:PHE:HD1	1.52	0.57
1:B:120:ARG:NH2	5:B:845:HOH:O	2.38	0.56
1:A:138:THR:HG21	1:B:142:ASN:HB2	1.88	0.56
1:B:304:HIS:HA	1:B:305:SER:HB2	1.88	0.56
1:B:61:LEU:HG	1:B:62:ASP:H	1.70	0.56
1:B:59:PRO:HG2	5:B:840:HOH:O	2.05	0.55
1:B:372:GLU:O	5:B:848:HOH:O	2.18	0.55
1:A:225:LYS:HE3	1:A:231:GLU:HG3	1.87	0.54
1:B:469:LEU:HD23	1:B:479:PHE:HB3	1.89	0.53
1:A:438:GLN:HG2	1:A:446:LEU:HB3	1.91	0.52
1:A:33:ASN:ND2	5:A:904:HOH:O	2.40	0.51
1:B:301:ASN:CB	1:B:302:MET:HA	2.40	0.51
1:B:329:THR:HG21	1:B:367:LEU:HD12	1.93	0.51
1:B:61:LEU:O	1:B:62:ASP:HB2	2.11	0.50
1:A:310:ARG:NH1	1:A:476:LEU:O	2.43	0.50
1:A:153:THR:OG1	1:A:164:HIS:NE2	2.37	0.50
1:B:61:LEU:HG	1:B:63:PHE:H	1.77	0.50
1:B:80:LEU:HG	1:B:84:LYS:HE3	1.94	0.49
1:B:58:SER:O	1:B:60:MET:HB2	2.12	0.49
1:A:165:ASN:ND2	1:A:172:ASP:OD2	2.41	0.49
1:A:129:VAL:HB	1:A:163:LEU:HB3	1.94	0.48
1:B:206:PHE:O	4:B:601:GOL:H12	2.14	0.48
1:B:442:PRO:HA	1:B:444:ALA:H	1.78	0.48
1:A:204:VAL:HG21	1:A:291:LEU:HD11	1.96	0.48
1:B:476:LEU:HD21	1:B:527:LEU:HD13	1.95	0.48
1:A:63:PHE:CZ	1:A:386:GLN:HB3	2.49	0.48
1:A:206:PHE:O	4:A:601:GOL:H12	2.14	0.48
1:B:329:THR:OG1	1:B:368:TYR:N	2.43	0.47
1:B:129:VAL:HB	1:B:163:LEU:HB3	1.96	0.47
1:B:347:PRO:HB3	1:B:350:TYR:HB3	1.96	0.47
1:B:209:HIS:HD2	5:B:833:HOH:O	1.97	0.47
1:A:408:PRO:HG3	1:A:430:HIS:HB3	1.96	0.47
1:B:297:PHE:CZ	1:B:327:TRP:HA	2.50	0.46
1:A:466:ASP:O	1:A:481:ARG:HA	2.15	0.46
1:B:442:PRO:CA	1:B:444:ALA:H	2.28	0.46
1:A:328:ALA:HB2	1:A:366:CYS:HB2	1.97	0.46
1:B:219:VAL:HG22	1:B:239:ARG:HG3	1.97	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:400:ARG:O	1:A:404:ARG:HD3	2.17	0.44
1:A:66:ASP:OD1	1:A:66:ASP:N	2.50	0.44
1:A:17:TYR:CE2	1:A:404:ARG:HG2	2.52	0.44
1:B:347:PRO:HB2	1:B:348:HIS:H	1.57	0.44
1:A:469:LEU:HD23	1:A:479:PHE:HB3	2.00	0.43
1:A:503:LEU:HD11	1:A:520:ARG:HD2	2.00	0.43
1:B:81:GLU:OE1	1:B:81:GLU:N	2.42	0.43
1:B:297:PHE:CE1	1:B:327:TRP:HA	2.53	0.43
1:A:219:VAL:HG22	1:A:239:ARG:HG3	1.99	0.43
1:A:112:TRP:CD1	1:A:124:LYS:HE2	2.54	0.43
1:B:4:ASN:N	5:B:719:HOH:O	2.51	0.43
1:B:153:THR:OG1	1:B:164:HIS:NE2	2.44	0.42
1:A:153:THR:HG1	1:A:164:HIS:CD2	2.32	0.42
1:B:508:ALA:N	1:B:536:GLN:O	2.49	0.42
1:B:175:PHE:HB3	1:B:208:PHE:CZ	2.54	0.42
1:B:235:TYR:O	1:B:240:HIS:NE2	2.41	0.42
1:B:326:CYS:SG	5:B:759:HOH:O	2.62	0.42
1:A:441:ASP:HB3	1:A:444:ALA:HB2	2.01	0.42
1:A:288:GLY:HA2	5:A:777:HOH:O	2.18	0.42
1:A:220:PRO:HB2	1:A:223:GLU:HG3	2.02	0.41
1:B:166:PHE:CZ	1:B:203:THR:HG21	2.55	0.41
1:A:172:ASP:HB3	4:A:601:GOL:H2	2.02	0.41
1:B:57:THR:O	1:B:58:SER:OG	2.33	0.41
1:B:466:ASP:O	1:B:481:ARG:HA	2.21	0.41
1:B:441:ASP:HA	1:B:442:PRO:HD2	1.67	0.40
1:A:438:GLN:HG3	1:A:444:ALA:CB	2.50	0.40
1:B:325:PRO:HG2	1:B:363:GLY:HA3	2.03	0.40
1:B:140:PRO:HG2	1:B:152:TRP:CE2	2.56	0.40
1:B:206:PHE:HA	1:B:244:LEU:HB2	2.03	0.40
1:B:453:LEU:HA	1:B:453:LEU:HD23	1.95	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	533/538 (99%)	516 (97%)	17 (3%)	0	100	100
1	B	409/538 (76%)	380 (93%)	16 (4%)	13 (3%)	4	5
All	All	942/1076 (88%)	896 (95%)	33 (4%)	13 (1%)	11	20

All (13) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	58	SER
1	B	62	ASP
1	B	220	PRO
1	B	298	ASP
1	B	299	LEU
1	B	302	MET
1	B	304	HIS
1	B	347	PRO
1	B	297	PHE
1	B	305	SER
1	B	348	HIS
1	B	442	PRO
1	B	321	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	458/461 (99%)	451 (98%)	7 (2%)	65	85
1	B	365/461 (79%)	356 (98%)	9 (2%)	47	73
All	All	823/922 (89%)	807 (98%)	16 (2%)	57	80

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	ARG
1	A	51	TRP
1	A	100	ASP
1	A	221	LYS
1	A	244	LEU
1	A	319	LEU
1	A	389	TYR
1	B	60	MET
1	B	61	LEU
1	B	62	ASP
1	B	63	PHE
1	B	244	LEU
1	B	300	LEU
1	B	301	ASN
1	B	304	HIS
1	B	366	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	165	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLC	C	1	2	12,12,12	0.53	0	17,17,17	0.56	0
2	GLC	C	2	2	11,11,12	0.51	0	15,15,17	1.73	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	C	1	2	-	0/2/22/22	0/1/1/1
2	GLC	C	2	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	2	GLC	C1-O5-C5	5.37	119.47	112.19


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	A	603	-	5,5,5	0.37	0	5,5,5	0.24	0
4	GOL	B	601	-	5,5,5	0.37	0	5,5,5	0.24	0
4	GOL	A	602	-	5,5,5	0.37	0	5,5,5	0.22	0
4	GOL	A	601	-	5,5,5	0.37	0	5,5,5	0.21	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	603	-	-	2/4/4/4	-
4	GOL	B	601	-	-	2/4/4/4	-
4	GOL	A	602	-	-	2/4/4/4	-
4	GOL	A	601	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	601	GOL	O1-C1-C2-C3
4	A	601	GOL	O1-C1-C2-C3
4	A	603	GOL	O1-C1-C2-C3
4	A	602	GOL	O1-C1-C2-C3
4	A	603	GOL	O1-C1-C2-O2
4	B	601	GOL	O1-C1-C2-O2
4	A	601	GOL	O1-C1-C2-O2
4	A	602	GOL	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	601	GOL	2	0
4	A	601	GOL	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	535/538 (99%)	0.01	6 (1%) 80 82	26, 34, 52, 68	0
1	B	421/538 (78%)	0.68	46 (10%) 5 5	28, 45, 86, 123	0
All	All	956/1076 (88%)	0.30	52 (5%) 25 27	26, 38, 71, 123	0

All (52) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	62	ASP	12.2
1	B	303	PRO	10.6
1	B	346	ASP	10.6
1	B	347	PRO	10.5
1	B	63	PHE	8.9
1	B	300	LEU	7.7
1	B	80	LEU	6.0
1	B	304	HIS	6.0
1	B	61	LEU	5.6
1	B	302	MET	5.5
1	B	231	GLU	5.4
1	B	60	MET	5.2
1	B	297	PHE	5.0
1	B	301	ASN	4.9
1	B	59	PRO	4.5
1	B	330	SER	4.5
1	B	441	ASP	4.5
1	B	529	ALA	3.7
1	B	329	THR	3.6
1	B	496	GLY	3.3
1	B	369	GLN	3.3
1	B	30	GLY	3.3
1	A	221	LYS	3.3
1	B	275	ASP	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	232	ALA	3.1
1	B	442	PRO	3.1
1	B	79	THR	3.0
1	B	495	THR	3.0
1	A	222	GLY	2.9
1	B	305	SER	2.9
1	B	527	LEU	2.6
1	B	22	LEU	2.6
1	B	494	LEU	2.6
1	B	371	GLU	2.5
1	B	504	PRO	2.4
1	A	521	ASP	2.4
1	B	296	THR	2.4
1	A	497	GLN	2.3
1	B	525	LEU	2.3
1	B	295	TYR	2.3
1	A	348	HIS	2.3
1	A	514	ALA	2.3
1	B	219	VAL	2.2
1	B	55	PHE	2.2
1	B	328	ALA	2.2
1	B	58	SER	2.2
1	B	444	ALA	2.2
1	B	298	ASP	2.2
1	B	85	ALA	2.2
1	B	372	GLU	2.0
1	B	348	HIS	2.0
1	B	4	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GLC	C	1	12/12	0.90	0.21	32,36,40,40	0
2	GLC	C	2	11/12	0.91	0.19	30,32,35,36	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	GOL	A	601	6/6	0.72	0.43	35,35,37,40	0
4	GOL	A	603	6/6	0.77	0.29	39,40,41,42	0
4	GOL	B	601	6/6	0.79	0.58	42,45,46,46	0
4	GOL	A	602	6/6	0.89	0.30	42,43,44,47	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	MG	A	600	1/1	0.89	0.11	39,39,39,39	0

6.5 Other polymers [i](#)

There are no such residues in this entry.