



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 15, 2021 – 02:33 PM EDT

PDB ID : 6WY8  
Title : Tcur3481-Tcur3483 steroid ACAD  
Authors : Kimber, M.S.; Stirling, A.J.; Seah, S.Y.K.  
Deposited on : 2020-05-12  
Resolution : 2.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.17.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.17.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	364	<div> <div>%</div> <div> <div></div> <div>87%</div> <div>12%</div> <div>.</div> </div> </div>
1	C	364	<div> <div>%</div> <div> <div></div> <div>90%</div> <div>10%</div> </div> </div>
2	B	407	<div> <div>%</div> <div> <div></div> <div>84%</div> <div>10%</div> <div>6%</div> </div> </div>
2	D	407	<div> <div>%</div> <div> <div></div> <div>80%</div> <div>14%</div> <div>6%</div> </div> </div>

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 12709 atoms, of which 130 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acyl-CoA dehydrogenase domain protein Tcur3481.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	363	Total	C	N	O	S	0	6	0
			2718	1711	481	518	8			
1	C	364	Total	C	N	O	S	0	5	0
			2720	1710	484	517	9			

- Molecule 2 is a protein called Acyl-CoA dehydrogenase domain protein Tcur3483.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	384	Total	C	N	O	S	0	5	0
			2976	1884	522	560	10			
2	D	384	Total	C	N	O	S	0	6	0
			2989	1889	524	565	11			

There are 40 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	-19	MET	-	initiating methionine	UNP D1AB78
B	-18	GLY	-	expression tag	UNP D1AB78
B	-17	SER	-	expression tag	UNP D1AB78
B	-16	SER	-	expression tag	UNP D1AB78
B	-15	HIS	-	expression tag	UNP D1AB78
B	-14	HIS	-	expression tag	UNP D1AB78
B	-13	HIS	-	expression tag	UNP D1AB78
B	-12	HIS	-	expression tag	UNP D1AB78
B	-11	HIS	-	expression tag	UNP D1AB78
B	-10	HIS	-	expression tag	UNP D1AB78
B	-9	SER	-	expression tag	UNP D1AB78
B	-8	SER	-	expression tag	UNP D1AB78
B	-7	GLY	-	expression tag	UNP D1AB78
B	-6	LEU	-	expression tag	UNP D1AB78
B	-5	VAL	-	expression tag	UNP D1AB78
B	-4	PRO	-	expression tag	UNP D1AB78

*Continued on next page...*

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	ARG	-	expression tag	UNP D1AB78
B	-2	GLY	-	expression tag	UNP D1AB78
B	-1	SER	-	expression tag	UNP D1AB78
B	0	HIS	-	expression tag	UNP D1AB78
D	-19	MET	-	initiating methionine	UNP D1AB78
D	-18	GLY	-	expression tag	UNP D1AB78
D	-17	SER	-	expression tag	UNP D1AB78
D	-16	SER	-	expression tag	UNP D1AB78
D	-15	HIS	-	expression tag	UNP D1AB78
D	-14	HIS	-	expression tag	UNP D1AB78
D	-13	HIS	-	expression tag	UNP D1AB78
D	-12	HIS	-	expression tag	UNP D1AB78
D	-11	HIS	-	expression tag	UNP D1AB78
D	-10	HIS	-	expression tag	UNP D1AB78
D	-9	SER	-	expression tag	UNP D1AB78
D	-8	SER	-	expression tag	UNP D1AB78
D	-7	GLY	-	expression tag	UNP D1AB78
D	-6	LEU	-	expression tag	UNP D1AB78
D	-5	VAL	-	expression tag	UNP D1AB78
D	-4	PRO	-	expression tag	UNP D1AB78
D	-3	ARG	-	expression tag	UNP D1AB78
D	-2	GLY	-	expression tag	UNP D1AB78
D	-1	SER	-	expression tag	UNP D1AB78
D	0	HIS	-	expression tag	UNP D1AB78

- Molecule 3 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula:  $C_6H_{14}O_4$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			24	6	14	4		
3	B	1	Total	C	H	O	0	0
			24	6	14	4		
3	D	1	Total	C	H	O	0	0
			24	6	14	4		

- Molecule 4 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	B	1	Total 84	C 27	H 31	N 9	O 15	P 2	0	0
4	D	1	Total 84	C 27	H 31	N 9	O 15	P 2	0	0

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Cl	0	0
			1	1		
5	C	2	Total	Cl	0	0
			2	2		
5	D	1	Total	Cl	0	0
			1	1		


- Molecule 6 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES)

(formula:  $C_6H_{13}NO_4S$ ).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
6	B	1	Total	C	H	N	O	S	0	0
			25	6	13	1	4	1		
6	D	1	Total	C	H	N	O	S	0	0
			25	6	13	1	4	1		

- Molecule 7 is SULFATE ION (three-letter code: SO<sub>4</sub>) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	C	1	Total	O	S	0	0
			5	4	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	C	1	Total	O	S	0	0
			5	4	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	224	Total	O	0	0
			224	224		
8	B	254	Total	O	0	0
			254	254		
8	C	229	Total	O	0	0
			229	229		
8	D	295	Total	O	0	0
			295	295		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Acyl-CoA dehydrogenase domain protein Tcur3481




- Molecule 1: Acyl-CoA dehydrogenase domain protein Tcur3481




- Molecule 2: Acyl-CoA dehydrogenase domain protein Tcur3483



- Molecule 2: Acyl-CoA dehydrogenase domain protein Tcur3483







## 4 Data and refinement statistics

Property	Value	Source
Space group	P 2 <sub>1</sub> 2 <sub>1</sub> 2 <sub>1</sub>	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	76.62Å 138.61Å 147.03Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.52 – 2.10 48.52 – 1.73	Depositor EDS
% Data completeness (in resolution range)	99.6 (48.52-2.10) 98.8 (48.52-1.73)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.14	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.36 (at 1.73Å)	Xtriage
Refinement program	PHENIX 1.15.2 _3472	Depositor
R, $R_{free}$	0.150 , 0.193 0.150 , 0.193	Depositor DCC
$R_{free}$ test set	8071 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	24.7	Xtriage
Anisotropy	0.340	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 45.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	12709	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.10% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PGE, SO4, FAD, MES, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.55	1/2773 (0.0%)	0.67	0/3787
1	C	0.56	0/2776	0.67	0/3788
2	B	0.55	0/3038	0.65	0/4123
2	D	0.56	0/3049	0.65	0/4136
All	All	0.55	1/11636 (0.0%)	0.66	0/15834

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	7	GLU	CD-OE2	5.26	1.31	1.25

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2718	0	2686	42	0
1	C	2720	0	2684	29	0
2	B	2976	0	2977	38	0
2	D	2989	0	2974	43	0
3	A	10	14	14	3	0
3	B	10	14	14	3	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	10	14	14	0	0
4	B	53	31	31	3	0
4	D	53	31	31	2	0
5	B	1	0	0	0	0
5	C	2	0	0	0	0
5	D	1	0	0	0	0
6	B	12	13	13	0	0
6	D	12	13	13	0	0
7	C	10	0	0	1	0
8	A	224	0	0	3	0
8	B	254	0	0	6	0
8	C	229	0	0	2	1
8	D	295	0	0	8	1
All	All	12579	130	11451	145	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 145 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:185:LEU:HD11	1:A:201:LEU:HD22	1.35	1.08
2:D:1:MET:N	8:D:501:HOH:O	1.91	1.02
1:C:185[A]:LEU:HD11	1:C:201:LEU:HD22	1.57	0.86
1:A:41:GLU:OE2	8:A:501:HOH:O	1.95	0.83
1:A:257:GLY:HA3	2:B:381:MET:CE	2.11	0.81

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:C:540:HOH:O	8:D:716:HOH:O[1_455]	2.07	0.13

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	367/364 (101%)	358 (98%)	9 (2%)	0	100	100
1	C	367/364 (101%)	361 (98%)	6 (2%)	0	100	100
2	B	387/407 (95%)	383 (99%)	4 (1%)	0	100	100
2	D	388/407 (95%)	383 (99%)	5 (1%)	0	100	100
All	All	1509/1542 (98%)	1485 (98%)	24 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	270/265 (102%)	268 (99%)	2 (1%)	84	88
1	C	270/265 (102%)	269 (100%)	1 (0%)	91	94
2	B	307/321 (96%)	304 (99%)	3 (1%)	76	82
2	D	308/321 (96%)	305 (99%)	3 (1%)	76	82
All	All	1155/1172 (98%)	1146 (99%)	9 (1%)	84	86

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	270[A]	GLU
2	D	270[B]	GLU
2	B	254	ARG
2	B	271	GLN
1	C	249	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	356	GLN
1	C	23	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

Of 13 ligands modelled in this entry, 4 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	MES	D	404	-	12,12,12	2.16	1 (8%)	14,16,16	1.68	3 (21%)
4	FAD	D	401	-	51,58,58	1.26	4 (7%)	60,89,89	2.31	10 (16%)
3	PGE	A	401	-	9,9,9	0.36	0	8,8,8	0.30	0
7	SO4	C	403	-	4,4,4	0.15	0	6,6,6	0.21	0
6	MES	B	404	-	12,12,12	2.11	1 (8%)	14,16,16	1.69	3 (21%)
3	PGE	D	403	-	9,9,9	0.34	0	8,8,8	0.33	0
3	PGE	B	403	-	9,9,9	0.36	0	8,8,8	0.40	0
4	FAD	B	401	-	51,58,58	1.32	5 (9%)	60,89,89	2.23	9 (15%)
7	SO4	C	404	-	4,4,4	0.20	0	6,6,6	0.34	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.  
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	MES	D	404	-	-	0/6/14/14	0/1/1/1
4	FAD	D	401	-	-	2/30/50/50	0/6/6/6
3	PGE	A	401	-	-	4/7/7/7	-
6	MES	B	404	-	-	1/6/14/14	0/1/1/1
3	PGE	D	403	-	-	1/7/7/7	-
3	PGE	B	403	-	-	3/7/7/7	-
4	FAD	B	401	-	-	1/30/50/50	0/6/6/6

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	D	404	MES	C8-S	-7.10	1.67	1.77
6	B	404	MES	C8-S	-6.92	1.67	1.77
4	D	401	FAD	C4X-C10	5.83	1.44	1.38
4	B	401	FAD	C4X-C10	5.23	1.44	1.38
4	B	401	FAD	C1'-N10	-3.44	1.44	1.48

The worst 5 of 25 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	401	FAD	C4-N3-C2	12.56	125.75	115.14
4	D	401	FAD	C4-N3-C2	12.31	125.54	115.14
4	B	401	FAD	C4X-C4-N3	-7.03	113.82	123.43
4	D	401	FAD	C4X-C4-N3	-6.77	114.17	123.43
4	D	401	FAD	C10-C4X-N5	4.72	124.52	121.26

There are no chirality outliers.

5 of 12 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	B	401	FAD	C2'-C1'-N10-C10
3	B	403	PGE	O2-C3-C4-O3
3	D	403	PGE	O1-C1-C2-O2
3	A	401	PGE	O2-C3-C4-O3
3	A	401	PGE	C6-C5-O3-C4

There are no ring outliers.


5 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	401	FAD	2	0
3	A	401	PGE	3	0
3	B	403	PGE	3	0
4	B	401	FAD	3	0
7	C	404	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	363/364 (99%)	-0.71	3 (0%) 86 88	16, 27, 47, 70	0
1	C	364/364 (100%)	-0.72	3 (0%) 86 88	17, 27, 50, 74	0
2	B	384/407 (94%)	-0.72	3 (0%) 86 88	15, 28, 52, 85	0
2	D	384/407 (94%)	-0.69	5 (1%) 77 80	18, 27, 49, 89	0
All	All	1495/1542 (96%)	-0.71	14 (0%) 84 86	15, 27, 50, 89	0

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	291	ALA	2.9
2	D	176	GLU	2.8
1	A	363	ALA	2.7
2	D	383	ARG	2.7
2	B	177	ALA	2.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	PGE	A	401	10/10	0.73	0.21	60,72,76,77	24
3	PGE	D	403	10/10	0.81	0.16	28,37,46,49	24
3	PGE	B	403	10/10	0.82	0.21	48,60,69,69	24
6	MES	D	404	12/12	0.84	0.17	45,54,61,67	25
6	MES	B	404	12/12	0.87	0.14	49,64,75,79	25
5	CL	C	402	1/1	0.89	0.15	76,76,76,76	0
7	SO4	C	404	5/5	0.93	0.12	36,41,42,46	5
7	SO4	C	403	5/5	0.94	0.18	44,49,52,55	5
5	CL	B	402	1/1	0.96	0.15	51,51,51,51	0
5	CL	D	402	1/1	0.96	0.14	52,52,52,52	0
4	FAD	D	401	53/53	0.97	0.07	21,29,53,59	0
4	FAD	B	401	53/53	0.97	0.07	18,29,48,61	0
5	CL	C	401	1/1	0.98	0.03	29,29,29,29	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.