



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 14, 2020 – 04:23 PM BST

PDB ID : 6WYA
Title : RTX (Reverse Transcription Xenopolymerase) in complex with a DNA duplex and dAMPNPP
Authors : Choi, W.S.; He, P.; Pothukuchy, A.; Gollihar, J.; Ellington, A.D.; Yang, W.
Deposited on : 2020-05-12
Resolution : 2.41 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.14.4.dev1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.14.4.dev1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.41 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4647 (2.44-2.40)
Clashscore	141614	5161 (2.44-2.40)
Ramachandran outliers	138981	5073 (2.44-2.40)
Sidechain outliers	138945	5074 (2.44-2.40)
RSRZ outliers	127900	4543 (2.44-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	774	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 15%, green 82%, grey 4%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> % 82% 15% •• </div> </div>
2	B	23	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 4%, orange 1%, yellow 43%, green 52%, grey 2%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 4% 52% 43% • </div> </div>
3	C	20	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, green 55%, yellow 35%, orange 10%);"></div> <div style="display: flex; justify-content: space-between; width: 100%;"> 55% 35% 10% </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
8	ACT	A	808	-	-	X	-

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 7443 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	756	Total	C	N	O	S	0	8	0
			6270	4042	1063	1149	16			

There are 19 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	38	LEU	PHE	conflict	UNP D0VWU9
A	97	MET	ARG	conflict	UNP D0VWU9
A	118	ILE	LYS	conflict	UNP D0VWU9
A	137	LEU	MET	conflict	UNP D0VWU9
A	147	HIS	GLU	conflict	UNP D0VWU9
A	210	ASP	ASN	conflict	UNP D0VWU9
A	381	HIS	ARG	conflict	UNP D0VWU9
A	384	HIS	TYR	conflict	UNP D0VWU9
A	389	ILE	VAL	conflict	UNP D0VWU9
A	466	ARG	LYS	conflict	UNP D0VWU9
A	493	LEU	TYR	conflict	UNP D0VWU9
A	514	ILE	THR	conflict	UNP D0VWU9
A	521	LEU	ILE	conflict	UNP D0VWU9
A	584	LYS	GLU	conflict	UNP D0VWU9
A	587	LEU	PHE	conflict	UNP D0VWU9
A	664	LYS	GLU	conflict	UNP D0VWU9
A	711	VAL	GLY	conflict	UNP D0VWU9
A	735	LYS	ASN	conflict	UNP D0VWU9
A	768	ARG	TRP	conflict	UNP D0VWU9

- Molecule 2 is a DNA chain called DNA strand 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	22	Total	C	N	O	P	0	0	0
			447	214	86	126	21			


- Molecule 3 is a DNA chain called DNA strand 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	20	Total	C	N	O	P	0	0	0
			408	196	71	122	19			

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Mg	0	0
			1	1		
4	A	4	Total	Mg	0	0
			4	4		

- Molecule 5 is 2'-deoxy-5'-O-[(R)-hydroxy{[(R)-hydroxy(phosphonooxy)phosphoryl]amino}phosphoryl]adenosine (three-letter code: DZ4) (formula: C₁₀H₁₇N₆O₁₁P₃) (labeled as "Ligand of Interest" by author).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			30	10	6	11	3		

- Molecule 6 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: C₄H₁₂NO₃).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 7 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			4	2	2		
7	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 8 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	230	Total	O	0	0
			230	230		
9	B	16	Total	O	0	0
			16	16		
9	C	17	Total	O	0	0
			17	17		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA polymerase



- Molecule 2: DNA strand 1



- Molecule 3: DNA strand 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	92.96 Å 96.36 Å 112.90 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.78 – 2.41 73.29 – 2.41	Depositor EDS
% Data completeness (in resolution range)	100.0 (42.78-2.41) 92.5 (73.29-2.41)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.18 (at 2.42 Å)	Xtriage
Refinement program	PHENIX 1.15.2_3472	Depositor
R, R_{free}	0.166 , 0.218 0.171 , 0.219	Depositor DCC
R_{free} test set	1004 reflections (2.52%)	wwPDB-VP
Wilson B-factor (Å ²)	45.0	Xtriage
Anisotropy	0.286	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 47.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.022 for k,h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7443	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, MG, ACT, EDO, DZ4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.47	0/6423	0.59	1/8665 (0.0%)
2	B	0.93	0/502	1.00	1/772 (0.1%)
3	C	0.87	0/456	1.14	6/703 (0.9%)
All	All	0.55	0/7381	0.68	8/10140 (0.1%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	20	DT	O4'-C4'-C3'	-6.15	102.04	104.50
3	C	18	DC	O5'-P-OP1	6.13	118.05	110.70
3	C	19	DC	O5'-P-OP2	-5.66	100.61	105.70
1	A	493	LEU	CA-CB-CG	-5.45	102.77	115.30
2	B	4	DA	O5'-P-OP2	-5.33	100.91	105.70
3	C	2	DC	O4'-C1'-N1	5.31	111.72	108.00
3	C	18	DC	C1'-O4'-C4'	-5.03	105.07	110.10
3	C	18	DC	O4'-C4'-C3'	-5.00	102.50	104.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6270	0	6355	85	0
2	B	447	0	248	10	0
3	C	408	0	229	8	0
4	A	4	0	0	0	0
4	B	1	0	0	0	0
5	A	30	0	13	0	0
6	A	8	0	12	1	0
7	A	4	0	6	0	0
7	B	4	0	6	2	0
8	A	4	0	3	2	0
9	A	230	0	0	7	0
9	B	16	0	0	0	0
9	C	17	0	0	0	0
All	All	7443	0	6872	97	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (97) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:48:ILE:CD1	1:A:83:TRP:CE3	2.52	0.92
1:A:243:ARG:HG3	1:A:243:ARG:HH11	1.34	0.89
1:A:48:ILE:HD12	1:A:83:TRP:CD2	2.08	0.88
1:A:48:ILE:HD13	1:A:83:TRP:CZ3	2.16	0.79
1:A:265:ARG:NH1	1:A:696:GLY:O	2.16	0.77
1:A:186:THR:HG23	1:A:189:GLU:H	1.49	0.76
1:A:591:LYS:HD3	2:B:7:DC:H5"	1.67	0.75
1:A:501[A]:ARG:HH22	2:B:2:DA:H5"	1.53	0.74
1:A:321:GLU:OE2	9:A:901:HOH:O	2.04	0.74
2:B:11:DC:H2"	2:B:12:DG:C8	2.25	0.72
1:A:354:GLU:OE1	9:A:902:HOH:O	2.09	0.70
1:A:48:ILE:CD1	1:A:83:TRP:CD2	2.76	0.69
1:A:247:ARG:NH1	1:A:694[B]:ARG:HE	1.93	0.67
1:A:4:ASP:OD1	6:A:806:TRS:H22	1.95	0.67
1:A:713:ARG:NH2	1:A:728:ASP:OD2	2.27	0.67
1:A:145:LEU:HB2	1:A:158:LEU:HD21	1.77	0.65
1:A:585:ARG:NH1	1:A:631:LEU:O	2.30	0.64
1:A:243:ARG:NH1	1:A:243:ARG:HG3	2.07	0.64
1:A:48:ILE:HD13	1:A:83:TRP:CE3	2.26	0.62
1:A:491[B]:ASN:O	9:A:903:HOH:O	2.15	0.62
1:A:523:MET:HG3	9:A:1051:HOH:O	2.00	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:22:DG:N2	3:C:3:DA:C2	2.71	0.58
1:A:525:ILE:HG23	1:A:536:VAL:HG21	1.85	0.58
1:A:1:MET:HE1	1:A:133:GLU:HB2	1.85	0.58
1:A:746:ARG:HH21	7:B:102:EDO:H11	1.70	0.57
2:B:11:DC:H2''	2:B:12:DG:N7	2.20	0.56
1:A:188:ARG:HG3	1:A:228:ILE:HD11	1.86	0.56
1:A:147:HIS:HB2	1:A:150:GLU:HG3	1.88	0.56
1:A:260:LEU:HD21	1:A:323:GLY:HA2	1.88	0.55
1:A:501[A]:ARG:NH1	2:B:2:DA:OP1	2.38	0.55
1:A:605:THR:HG23	1:A:608:LEU:HD12	1.89	0.55
1:A:247:ARG:HH12	1:A:694[B]:ARG:HE	1.55	0.54
2:B:15:DC:H2''	2:B:16:DA:C8	2.43	0.54
3:C:2:DC:H2'	3:C:3:DA:C8	2.43	0.54
1:A:754:ASP:OD1	1:A:754:ASP:N	2.41	0.53
1:A:685:ARG:NH2	1:A:714:ALA:O	2.41	0.53
1:A:289:LYS:HE2	1:A:311:TYR:OH	2.09	0.53
3:C:2:DC:H2'	3:C:3:DA:N7	2.24	0.53
1:A:48:ILE:HD12	1:A:83:TRP:CE3	2.33	0.52
1:A:463:ILE:HG21	1:A:483:GLN:HB2	1.92	0.52
1:A:743[B]:ARG:NH2	9:A:916:HOH:O	2.42	0.52
1:A:145:LEU:HD23	1:A:156:PRO:HD2	1.93	0.51
3:C:11:DT:H6	3:C:11:DT:H5''	1.75	0.51
1:A:641:ARG:O	1:A:645:GLU:HG3	2.11	0.50
1:A:48:ILE:O	1:A:51:VAL:HG22	2.12	0.50
1:A:611:VAL:HG23	3:C:18:DC:H5''	1.93	0.50
1:A:364:ARG:NH2	1:A:451:SER:OG	2.46	0.49
1:A:739:PRO:O	1:A:743[B]:ARG:HG3	2.13	0.49
1:A:157:ILE:HG13	1:A:222:ARG:HG3	1.93	0.49
1:A:243:ARG:NH1	1:A:243:ARG:CG	2.73	0.49
1:A:428:CYS:HG	1:A:442:CYS:HG	1.60	0.49
1:A:579:TYR:OH	1:A:582:PHE:HB2	2.14	0.48
1:A:571:LEU:O	8:A:808:ACT:O	2.32	0.48
2:B:1:DT:O2	2:B:1:DT:H2'	2.14	0.48
1:A:575:LEU:O	8:A:808:ACT:OXT	2.32	0.48
1:A:53:LYS:HE3	1:A:54:ILE:HD13	1.94	0.48
1:A:756:ARG:HA	1:A:756:ARG:HH11	1.78	0.47
1:A:746:ARG:NH2	7:B:102:EDO:H11	2.28	0.47
3:C:10:DG:H2'	3:C:11:DT:C6	2.50	0.47
1:A:49:GLU:HA	1:A:49:GLU:OE2	2.15	0.47
1:A:354:GLU:OE2	1:A:496:TYR:OH	2.21	0.47
2:B:21:DT:H2'	2:B:22:DG:C8	2.51	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:590:THR:HG22	1:A:593[A]:LYS:HB2	1.98	0.45
1:A:332:GLN:OE1	1:A:335:ARG:NH1	2.49	0.45
1:A:153:ALA:HA	1:A:218:TYR:CZ	2.51	0.45
1:A:55:THR:HG22	1:A:64:THR:HG22	1.99	0.45
3:C:8:DT:H2"	3:C:9:DG:C8	2.52	0.45
1:A:119:ARG:HD2	1:A:123:ASP:OD2	2.16	0.45
1:A:296:THR:O	1:A:300:GLU:HG3	2.17	0.45
1:A:567:ILE:HA	1:A:567:ILE:HD12	1.87	0.44
1:A:390:LYS:HE2	1:A:536:VAL:O	2.18	0.44
1:A:22:GLU:OE1	1:A:27:LYS:HD2	2.17	0.44
1:A:41:LEU:HG	1:A:107:ILE:HD12	1.99	0.44
1:A:752:LYS:HD2	1:A:752:LYS:HA	1.40	0.44
1:A:93:VAL:HB	1:A:94:PRO:HD3	2.01	0.43
1:A:124:LYS:NZ	9:A:909:HOH:O	2.29	0.43
1:A:295:ILE:HG12	1:A:308:VAL:HG13	2.00	0.43
1:A:676:THR:HG23	1:A:680:VAL:HG12	1.99	0.43
1:A:564:LEU:HB2	9:A:1088:HOH:O	2.18	0.43
1:A:657:PRO:HB2	1:A:717:PHE:HE1	1.83	0.43
3:C:3:DA:H5"	3:C:3:DA:H8	1.84	0.43
1:A:472:ASP:OD1	1:A:474:ILE:HG12	2.18	0.43
1:A:644:LYS:NZ	1:A:754:ASP:HB2	2.34	0.43
1:A:266:ARG:NH1	1:A:347:SER:O	2.51	0.43
1:A:558:LYS:HB3	1:A:558:LYS:HE3	1.67	0.43
1:A:501[A]:ARG:NH2	2:B:2:DA:H5"	2.29	0.42
1:A:648:GLU:HG2	1:A:652:LYS:HE2	2.01	0.42
1:A:590:THR:HG22	1:A:593[B]:LYS:HB2	2.02	0.42
1:A:70:LYS:HD3	1:A:83:TRP:CZ2	2.55	0.42
1:A:218:TYR:O	1:A:222:ARG:HG2	2.20	0.41
1:A:394:ARG:NH2	1:A:590:THR:OG1	2.54	0.41
1:A:159:MET:HG2	1:A:172:THR:HB	2.03	0.41
1:A:564:LEU:CD1	1:A:579:TYR:HB2	2.51	0.41
1:A:753:GLU:HB2	1:A:755:LEU:HD21	2.02	0.41
1:A:7:TYR:HB3	1:A:16:ILE:HD13	2.02	0.41
1:A:465:LYS:HE3	1:A:465:LYS:HB3	1.91	0.40
1:A:514:ILE:HA	1:A:514:ILE:HD13	1.80	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	762/774 (98%)	737 (97%)	25 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	668/675 (99%)	649 (97%)	19 (3%)	43	62

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	21	LYS
1	A	46	SER
1	A	53	LYS
1	A	59	HIS
1	A	199	LYS
1	A	244	MET
1	A	289	LYS
1	A	304	ASN
1	A	334	SER
1	A	382	GLN
1	A	383	SER
1	A	483	GLN
1	A	516	TRP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	689	ARG
1	A	694[A]	ARG
1	A	694[B]	ARG
1	A	751	ARG
1	A	752	LYS
1	A	754	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	304	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 5 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	TRS	A	806	-	7,7,7	0.54	0	9,9,9	0.48	0
7	EDO	A	807	-	3,3,3	0.08	0	2,2,2	0.17	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	ACT	A	808	-	1,3,3	0.28	0	0,3,3	0.00	-
7	EDO	B	102	-	3,3,3	0.52	0	2,2,2	0.16	0
5	DZ4	A	805	4	29,32,32	1.47	3 (10%)	33,50,50	1.36	4 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	TRS	A	806	-	-	0/9/9/9	-
7	EDO	A	807	-	-	1/1/1/1	-
7	EDO	B	102	-	-	0/1/1/1	-
5	DZ4	A	805	4	-	5/15/34/34	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	805	DZ4	PB-N3A	4.35	1.74	1.63
5	A	805	DZ4	PA-N3A	4.33	1.74	1.63
5	A	805	DZ4	C5-C4	2.32	1.47	1.40

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	805	DZ4	N3-C2-N1	-3.15	123.75	128.68
5	A	805	DZ4	PG-O3B-PB	-2.81	122.73	132.62
5	A	805	DZ4	C2'-C1'-N9	-2.66	108.13	114.27
5	A	805	DZ4	C4-C5-N7	-2.57	106.72	109.40

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	A	805	DZ4	PA-N3A-PB-O1B
5	A	805	DZ4	PB-O3B-PG-O2G
5	A	805	DZ4	O4'-C4'-C5'-O5'
7	A	807	EDO	O1-C1-C2-O2
5	A	805	DZ4	PB-O3B-PG-O1G
5	A	805	DZ4	C3'-C4'-C5'-O5'

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	806	TRS	1	0
8	A	808	ACT	2	0
7	B	102	EDO	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	756/774 (97%)	-0.27	7 (0%) 84 82	33, 57, 100, 142	0
2	B	22/23 (95%)	-0.33	1 (4%) 33 31	48, 86, 149, 160	0
3	C	20/20 (100%)	-0.45	0 100 100	44, 127, 130, 135	0
All	All	798/817 (97%)	-0.28	8 (1%) 82 80	33, 58, 111, 160	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	752	LYS	3.8
1	A	754	ASP	3.0
1	A	751	ARG	2.7
1	A	303	GLU	2.4
1	A	134	GLU	2.3
2	B	1	DT	2.2
1	A	657	PRO	2.2
1	A	46	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MG	A	801	1/1	0.52	0.21	87,87,87,87	0
8	ACT	A	808	4/4	0.70	0.21	104,106,107,107	0
4	MG	B	101	1/1	0.77	0.09	74,74,74,74	0
7	EDO	B	102	4/4	0.82	0.21	79,81,82,82	0
6	TRS	A	806	8/8	0.84	0.22	76,80,82,84	0
7	EDO	A	807	4/4	0.86	0.31	92,93,93,94	0
5	DZ4	A	805	30/30	0.95	0.14	55,65,80,83	0
4	MG	A	802	1/1	0.96	0.08	75,75,75,75	0
4	MG	A	803	1/1	0.98	0.09	48,48,48,48	0
4	MG	A	804	1/1	0.98	0.04	57,57,57,57	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around DZ4 A 805:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.