



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 10, 2020 – 10:37 AM BST

PDB ID : 6WYR  
Title : Crystal structure of anti-Muscle Specific Kinase (MuSK) Fab, MuSK1A  
Authors : Vieni, C.; Ekiert, D.  
Deposited on : 2020-05-13  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.14.3.dev2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.14.3.dev2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	214	<div> <div>3%</div> <div>95%</div> <div>6%</div> <div>•</div> </div>
1	M	214	<div> <div>6%</div> <div>93%</div> <div>6%</div> <div>•</div> </div>
2	H	232	<div> <div>2%</div> <div>90%</div> <div>6%</div> <div>•</div> </div>
2	I	232	<div> <div>3%</div> <div>91%</div> <div>7%</div> <div>•</div> </div>
3	A	3	<div> <div>100%</div> </div>
4	B	2	<div> <div>100%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	EDO	H	308	-	-	-	X

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 14243 atoms, of which 6747 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MuSK1A light chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	L	213	Total	C	H	N	O	S	0	11	0
			3281	1039	1612	277	346	7			
1	M	212	Total	C	H	N	O	S	0	6	0
			3209	1018	1574	270	339	8			

- Molecule 2 is a protein called MuSK1A heavy chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	H	222	Total	C	H	N	O	S	0	11	0
			3401	1080	1690	287	336	8			
2	I	226	Total	C	H	N	O	S	0	17	0
			3464	1094	1716	295	349	10			

- Molecule 3 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	3	Total	C	H	N	O	0	0	0
			73	22	34	2	15			

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	B	2	Total	C	H	N	O	0	0	0
			53	16	25	2	10			

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	L	1	Total	C	H	O	0	0
			10	2	6	2		
5	L	1	Total	C	H	O	0	0
			10	2	6	2		
5	H	1	Total	C	H	O	0	0
			10	2	6	2		
5	H	1	Total	C	H	O	0	0
			10	2	6	2		
5	H	1	Total	C	H	O	0	0
			10	2	6	2		
5	H	1	Total	C	H	O	0	0
			10	2	6	2		
5	M	1	Total	C	H	O	0	0
			10	2	6	2		
5	M	1	Total	C	H	O	0	0
			10	2	6	2		
5	M	1	Total	C	H	O	0	0
			10	2	6	2		
5	M	1	Total	C	H	O	0	0
			10	2	6	2		
5	I	1	Total	C	H	O	0	0
			10	2	6	2		
5	I	1	Total	C	H	O	0	0
			10	2	6	2		
5	I	1	Total	C	H	O	0	0
			10	2	6	2		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	I	1	Total	C	H	O	0	0
			10	2	6	2		
5	I	1	Total	C	H	O	0	0
			10	2	6	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	L	101	Total	O	0	0
			101	101		
6	H	197	Total	O	0	0
			197	197		
6	M	91	Total	O	0	0
			91	91		
6	I	213	Total	O	0	0
			213	213		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: MuSK1A light chain




- Molecule 1: MuSK1A light chain



- Molecule 2: MuSK1A heavy chain



- Molecule 2: MuSK1A heavy chain



- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain B:

100%

MG1  
MG2



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	65.87Å 71.72Å 106.18Å 90.00° 104.37° 90.00°	Depositor
Resolution (Å)	49.04 – 1.80 49.04 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.5 (49.04-1.80) 91.6 (49.04-1.80)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.63 (at 1.79Å)	Xtriage
Refinement program	PHENIX 1.17.1 _3660	Depositor
R, $R_{free}$	0.180 , 0.213 0.181 , 0.213	Depositor DCC
$R_{free}$ test set	2005 reflections (2.27%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.2	Xtriage
Anisotropy	0.246	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 63.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	14243	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.16% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, PCA, NAG, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	L	0.25	0/1735	0.47	0/2362
1	M	0.26	0/1682	0.47	0/2293
2	H	0.27	0/1778	0.50	0/2421
2	I	0.27	0/1840	0.53	0/2506
All	All	0.26	0/7035	0.49	0/9582

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1669	1612	1590	5	0
1	M	1635	1574	1574	6	0
2	H	1711	1690	1666	9	0
2	I	1748	1716	1660	12	0
3	A	39	34	34	0	0
4	B	28	25	25	0	0
5	H	20	30	30	0	0
5	I	20	30	30	0	0
5	L	8	12	12	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	M	16	24	24	0	0
6	H	197	0	0	2	0
6	I	213	0	0	2	0
6	L	101	0	0	0	0
6	M	91	0	0	1	0
All	All	7496	6747	6645	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:I:58:PRO:O	2:I:80:ARG:NH1	2.36	0.58
2:H:144:LYS:NZ	6:H:404:HOH:O	2.47	0.48
1:L:13:LEU:HD21	1:L:19:ALA:HB2	1.96	0.47
1:M:107:TRP:HZ2	2:I:107:GLU:OE2	1.98	0.46
2:I:165[C]:VAL:HG23	2:I:193:LEU:HD21	1.97	0.46
1:M:116:VAL:HB	2:I:52:TRP:CD2	2.50	0.46
2:I:59:TYR:C	2:I:80:ARG:HH12	2.20	0.45
1:L:123:ARG:CZ	1:L:163[B]:ARG:HH22	2.29	0.45
1:M:125:THR:HG21	1:M:162:PRO:HB2	1.99	0.45
1:L:163[B]:ARG:HD3	1:L:194:TYR:CE2	2.53	0.44
2:I:59:TYR:O	2:I:80:ARG:NH1	2.51	0.43
2:H:37:TYR:CE2	2:H:108:VAL:HG22	2.53	0.43
1:L:107:TRP:HZ2	2:H:107:GLU:OE2	2.01	0.43
1:M:75:ARG:HB2	1:M:92:SER:O	2.18	0.43
2:I:193:LEU:HD12	2:I:193:LEU:C	2.39	0.43
2:I:229:LYS:NZ	6:I:410:HOH:O	2.51	0.43
1:M:163:ARG:NH2	6:M:409:HOH:O	2.51	0.43
2:I:63:GLY:N	2:I:80:ARG:NH1	2.67	0.42
2:H:160:TYR:CE1	2:H:165[B]:VAL:HG23	2.54	0.42
1:M:205:ALA:O	1:M:209:LYS:HG3	2.19	0.42
2:H:69:GLN:NE2	6:H:402:HOH:O	2.45	0.41
2:I:106:ARG:O	2:I:115:SER:HA	2.20	0.41
1:L:116:VAL:HB	2:H:52:TRP:CG	2.55	0.41
2:I:158:LYS:NZ	2:I:186:GLN:OE1	2.54	0.41
2:H:106:ARG:O	2:H:115:SER:HA	2.20	0.41
2:I:95:ARG:NH1	6:I:414:HOH:O	2.54	0.41
2:H:193:LEU:C	2:H:193:LEU:HD12	2.41	0.40
2:H:165[B]:VAL:HG22	2:H:215:HIS:CD2	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	222/214 (104%)	215 (97%)	6 (3%)	1 (0%)	29	15
1	M	216/214 (101%)	209 (97%)	7 (3%)	0	100	100
2	H	229/232 (99%)	225 (98%)	4 (2%)	0	100	100
2	I	241/232 (104%)	236 (98%)	5 (2%)	0	100	100
All	All	908/892 (102%)	885 (98%)	22 (2%)	1 (0%)	51	36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	L	109	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	195/186 (105%)	194 (100%)	1 (0%)	88	87
1	M	190/186 (102%)	185 (97%)	5 (3%)	46	32
2	H	196/194 (101%)	195 (100%)	1 (0%)	88	87
2	I	205/194 (106%)	204 (100%)	1 (0%)	88	87
All	All	786/760 (103%)	778 (99%)	8 (1%)	73	71

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	29	ASP
2	H	224	LYS
1	M	13	LEU
1	M	125	THR
1	M	129	GLN
1	M	186	GLU
1	M	190	LYS
2	I	111	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	PCA	I	1	2	7,8,9	2.19	2 (28%)	9,10,12	2.01	4 (44%)
2	PCA	H	1	2	7,8,9	2.18	2 (28%)	9,10,12	2.06	5 (55%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PCA	I	1	2	-	0/0/11/13	0/1/1/1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PCA	H	1	2	-	0/0/11/13	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	I	1	PCA	CD-N	4.64	1.46	1.34
2	H	1	PCA	CD-N	4.60	1.46	1.34
2	H	1	PCA	CA-N	3.36	1.50	1.46
2	I	1	PCA	CA-N	3.34	1.50	1.46

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	I	1	PCA	CA-N-CD	-3.10	102.96	113.58
2	H	1	PCA	CA-N-CD	-3.10	102.96	113.58
2	H	1	PCA	OE-CD-CG	-2.98	121.57	126.76
2	I	1	PCA	OE-CD-CG	-2.92	121.67	126.76
2	H	1	PCA	CB-CA-N	2.62	110.82	103.30
2	I	1	PCA	CB-CA-N	2.49	110.46	103.30
2	H	1	PCA	CG-CD-N	2.44	114.70	108.39
2	I	1	PCA	CG-CD-N	2.42	114.65	108.39
2	H	1	PCA	CB-CA-C	-2.00	109.94	112.70

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates

5 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NAG	A	1	3,2	14,14,15	0.34	0	17,19,21	0.43	0
3	NAG	A	2	3	14,14,15	0.20	0	17,19,21	0.46	0
3	BMA	A	3	3	11,11,12	0.53	0	15,15,17	0.87	0
4	NAG	B	1	2,4	14,14,15	0.41	0	17,19,21	0.45	0
4	NAG	B	2	4	14,14,15	0.27	0	17,19,21	0.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	1	3,2	-	2/6/23/26	0/1/1/1
3	NAG	A	2	3	-	2/6/23/26	0/1/1/1
3	BMA	A	3	3	-	2/2/19/22	0/1/1/1
4	NAG	B	1	2,4	-	0/6/23/26	0/1/1/1
4	NAG	B	2	4	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	2	NAG	O5-C5-C6-O6
3	A	2	NAG	C4-C5-C6-O6
4	B	2	NAG	O5-C5-C6-O6
4	B	2	NAG	C4-C5-C6-O6
3	A	3	BMA	O5-C5-C6-O6
3	A	3	BMA	C4-C5-C6-O6
3	A	1	NAG	C4-C5-C6-O6
3	A	1	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

## 5.6 Ligand geometry

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	EDO	M	301	-	3,3,3	0.44	0	2,2,2	0.33	0
5	EDO	L	302	-	3,3,3	0.44	0	2,2,2	0.33	0
5	EDO	M	304	-	3,3,3	0.43	0	2,2,2	0.55	0
5	EDO	H	304	-	3,3,3	0.44	0	2,2,2	0.36	0
5	EDO	I	306	-	3,3,3	0.44	0	2,2,2	0.46	0
5	EDO	I	305	-	3,3,3	0.47	0	2,2,2	0.28	0
5	EDO	I	304	-	3,3,3	0.46	0	2,2,2	0.33	0
5	EDO	I	303	-	3,3,3	0.44	0	2,2,2	0.53	0
5	EDO	M	303	-	3,3,3	0.46	0	2,2,2	0.29	0
5	EDO	H	305	-	3,3,3	0.44	0	2,2,2	0.30	0
5	EDO	M	302	-	3,3,3	0.47	0	2,2,2	0.31	0
5	EDO	I	307	-	3,3,3	0.41	0	2,2,2	0.64	0
5	EDO	L	301	-	3,3,3	0.48	0	2,2,2	0.30	0
5	EDO	H	306	-	3,3,3	0.47	0	2,2,2	0.34	0
5	EDO	H	308	-	3,3,3	0.46	0	2,2,2	0.28	0
5	EDO	H	307	-	3,3,3	0.44	0	2,2,2	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	M	301	-	-	1/1/1/1	-
5	EDO	L	302	-	-	0/1/1/1	-
5	EDO	M	304	-	-	0/1/1/1	-
5	EDO	H	304	-	-	0/1/1/1	-
5	EDO	I	306	-	-	0/1/1/1	-
5	EDO	I	305	-	-	0/1/1/1	-
5	EDO	I	304	-	-	0/1/1/1	-
5	EDO	I	303	-	-	0/1/1/1	-
5	EDO	M	303	-	-	0/1/1/1	-
5	EDO	H	305	-	-	0/1/1/1	-
5	EDO	M	302	-	-	0/1/1/1	-
5	EDO	I	307	-	-	0/1/1/1	-

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	L	301	-	-	0/1/1/1	-
5	EDO	H	306	-	-	0/1/1/1	-
5	EDO	H	308	-	-	0/1/1/1	-
5	EDO	H	307	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	M	301	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	L	213/214 (99%)	-0.13	7 (3%) 46 40	25, 43, 88, 156	0
1	M	212/214 (99%)	0.03	12 (5%) 23 19	23, 46, 88, 173	0
2	H	221/232 (95%)	-0.34	4 (1%) 68 64	20, 29, 59, 107	0
2	I	225/232 (96%)	-0.29	8 (3%) 42 37	20, 27, 58, 257	0
All	All	871/892 (97%)	-0.19	31 (3%) 42 37	20, 35, 81, 257	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	L	235	CYS	6.4
2	I	110	GLY	6.1
2	I	112(A)	TYR	6.0
1	M	235	CYS	5.7
1	L	2	TYR	5.0
1	L	114	SER	4.7
2	H	112(A)	TYR	4.6
2	I	111	ASP	4.6
2	I	230	SER	3.8
2	I	111(A)	THR	3.7
1	M	126	VAL	3.6
2	I	231	CYS	3.6
2	I	112(B)	GLY	3.5
2	H	112(B)	GLY	3.3
1	M	96	ALA	3.3
1	M	107	TRP	3.2
1	L	109	GLY	3.2
1	M	13	LEU	3.0
1	L	107	TRP	2.7
2	H	230	SER	2.5
1	M	14	SER	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	L	27	LYS	2.3
1	M	26	ASP	2.3
1	M	18	THR	2.2
1	M	27	LYS	2.2
1	M	95	GLN	2.2
2	I	109	ARG	2.2
1	L	108	ASP	2.2
2	H	67[A]	TYR	2.1
1	M	84	GLY	2.1
1	M	114	SER	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	PCA	H	1	8/9	0.91	0.10	27,61,69,75	0
2	PCA	I	1	8/9	0.92	0.14	32,66,80,80	0

## 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	BMA	A	3	11/12	0.44	0.27	142,154,182,185	0
3	NAG	A	2	14/15	0.52	0.24	114,165,316,316	0
4	NAG	B	2	14/15	0.58	0.25	127,147,173,191	0
4	NAG	B	1	14/15	0.76	0.13	64,92,111,120	0
3	NAG	A	1	14/15	0.93	0.12	37,59,74,96	0

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column

labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	EDO	M	304	4/4	0.58	0.22	83,99,111,111	0
5	EDO	M	302	4/4	0.71	0.17	43,54,59,64	0
5	EDO	H	308	4/4	0.72	0.44	100,120,124,125	0
5	EDO	I	305	4/4	0.77	0.14	31,38,45,54	0
5	EDO	M	303	4/4	0.80	0.14	54,66,79,79	0
5	EDO	M	301	4/4	0.82	0.13	55,66,79,87	0
5	EDO	L	302	4/4	0.85	0.12	42,50,61,70	0
5	EDO	H	306	4/4	0.89	0.13	32,38,46,51	0
5	EDO	L	301	4/4	0.91	0.10	30,38,46,53	0
5	EDO	H	307	4/4	0.92	0.21	29,48,58,58	0
5	EDO	I	304	4/4	0.95	0.10	34,40,45,50	0
5	EDO	I	303	4/4	0.95	0.09	32,40,48,48	0
5	EDO	I	307	4/4	0.96	0.13	23,41,74,74	0
5	EDO	I	306	4/4	0.97	0.09	29,39,48,48	0
5	EDO	H	304	4/4	0.97	0.13	33,41,64,64	0
5	EDO	H	305	4/4	0.99	0.14	26,33,61,73	0

## 6.5 Other polymers

There are no such residues in this entry.