



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 17, 2020 – 04:29 am BST

PDB ID : 2WZJ  
Title : Catalytic and UBA domain of kinase MARK2/(Par-1) K82R, T208E double mutant  
Authors : Panneerselvam, S.; Marx, A.; Mandelkow, E.-M.; Mandelkow, E.  
Deposited on : 2009-11-30  
Resolution : 2.79 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4107 (2.80-2.76)
Clashscore	141614	4575 (2.80-2.76)
Ramachandran outliers	138981	4487 (2.80-2.76)
Sidechain outliers	138945	4489 (2.80-2.76)
RSRZ outliers	127900	4027 (2.80-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	327	 2% 67% 24% 5% 5%
1	B	327	 2% 69% 23% 5% 5%
1	C	327	 2% 66% 25% 5% 5%
1	D	327	 2% 67% 26% 5% 5%
1	E	327	 5% 64% 29% 5% 5%
1	F	327	 8% 60% 32% 5% 5%

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 15326 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SERINE/THREONINE-PROTEIN KINASE MARK2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	310	Total	C	N	O	S	0	0	0
			2531	1627	432	459	13			
1	B	310	Total	C	N	O	S	0	0	0
			2531	1627	432	459	13			
1	C	310	Total	C	N	O	S	0	0	0
			2531	1627	432	459	13			
1	D	313	Total	C	N	O	S	0	0	0
			2556	1642	436	465	13			
1	E	310	Total	C	N	O	S	0	0	0
			2531	1627	432	459	13			
1	F	310	Total	C	N	O	S	0	0	0
			2531	1627	432	459	13			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	82	ARG	LYS	engineered mutation	UNP O08679
A	208	GLU	THR	engineered mutation	UNP O08679
B	82	ARG	LYS	engineered mutation	UNP O08679
B	208	GLU	THR	engineered mutation	UNP O08679
C	82	ARG	LYS	engineered mutation	UNP O08679
C	208	GLU	THR	engineered mutation	UNP O08679
D	82	ARG	LYS	engineered mutation	UNP O08679
D	208	GLU	THR	engineered mutation	UNP O08679
E	82	ARG	LYS	engineered mutation	UNP O08679
E	208	GLU	THR	engineered mutation	UNP O08679
F	82	ARG	LYS	engineered mutation	UNP O08679
F	208	GLU	THR	engineered mutation	UNP O08679


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	36	Total 36	O 36	0	0
2	B	31	Total 31	O 31	0	0
2	C	13	Total 13	O 13	0	0
2	D	24	Total 24	O 24	0	0
2	E	3	Total 3	O 3	0	0
2	F	8	Total 8	O 8	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: SERINE/THREONINE-PROTEIN KINASE MARK2




#### • Molecule 1: SERINE/THREONINE-PROTEIN KINASE MARK2




#### • Molecule 1: SERINE/THREONINE-PROTEIN KINASE MARK2








• Molecule 1: SERINE/THREONINE-PROTEIN KINASE MARK2



• Molecule 1: SERINE/THREONINE-PROTEIN KINASE MARK2



• Molecule 1: SERINE/THREONINE-PROTEIN KINASE MARK2





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	162.24Å 205.58Å 91.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.39 – 2.79 45.39 – 2.79	Depositor EDS
% Data completeness (in resolution range)	99.5 (45.39-2.79) 99.5 (45.39-2.79)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.02 (at 2.77Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.179 , 0.246 0.171 , 0.238	Depositor DCC
$R_{free}$ test set	3851 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	62.8	Xtriage
Anisotropy	0.391	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 64.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	15326	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	76.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.60% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.46	0/2582	0.59	0/3476
1	B	0.44	0/2582	0.62	0/3476
1	C	0.42	0/2582	0.57	0/3476
1	D	0.42	0/2608	0.60	0/3512
1	E	0.35	0/2582	0.52	0/3476
1	F	0.35	0/2582	0.51	0/3476
All	All	0.41	0/15518	0.57	0/20892

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2531	0	2568	72	0
1	B	2531	0	2568	72	0
1	C	2531	0	2568	74	0
1	D	2556	0	2589	93	0
1	E	2531	0	2568	68	0
1	F	2531	0	2568	84	0
2	A	36	0	0	0	0
2	B	31	0	0	0	0
2	C	13	0	0	0	0
2	D	24	0	0	4	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	E	3	0	0	0	0
2	F	8	0	0	1	0
All	All	15326	0	15429	439	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 439 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:46:GLU:HG2	1:D:54:ARG:HE	1.24	1.03
1:F:73:ILE:HG22	1:F:74:LEU:HD23	1.38	1.01
1:A:90:ASN:H	1:A:90:ASN:HD22	0.97	0.95
1:B:104:MET:HE2	1:B:196:PHE:HB3	1.49	0.90
1:C:90:ASN:H	1:C:90:ASN:HD22	1.17	0.90

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	306/327 (94%)	289 (94%)	17 (6%)	0	100	100
1	B	306/327 (94%)	293 (96%)	12 (4%)	1 (0%)	41	70
1	C	306/327 (94%)	287 (94%)	19 (6%)	0	100	100
1	D	309/327 (94%)	287 (93%)	19 (6%)	3 (1%)	15	41
1	E	306/327 (94%)	276 (90%)	30 (10%)	0	100	100
1	F	306/327 (94%)	279 (91%)	26 (8%)	1 (0%)	41	70
All	All	1839/1962 (94%)	1711 (93%)	123 (7%)	5 (0%)	41	70

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	202	PHE
1	D	47	GLN
1	D	323	TYR
1	F	73	ILE
1	B	334	SER

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	277/289 (96%)	257 (93%)	20 (7%)	14	36
1	B	277/289 (96%)	262 (95%)	15 (5%)	22	50
1	C	277/289 (96%)	254 (92%)	23 (8%)	11	29
1	D	280/289 (97%)	265 (95%)	15 (5%)	22	50
1	E	277/289 (96%)	260 (94%)	17 (6%)	18	45
1	F	277/289 (96%)	262 (95%)	15 (5%)	22	50
All	All	1665/1734 (96%)	1560 (94%)	105 (6%)	18	43

5 of 105 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	256	LYS
1	D	46	GLU
1	F	256	LYS
1	C	268	ARG
1	C	288	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 33 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	198	ASN
1	C	343	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	F	198	ASN
1	C	222	GLN
1	C	253	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	310/327 (94%)	-0.18	2 (0%) 89 88	34, 58, 99, 154	0
1	B	310/327 (94%)	-0.11	7 (2%) 60 55	34, 60, 100, 132	0
1	C	310/327 (94%)	-0.03	5 (1%) 72 69	43, 66, 100, 133	0
1	D	313/327 (95%)	-0.09	6 (1%) 66 63	37, 62, 105, 173	0
1	E	310/327 (94%)	0.23	16 (5%) 27 22	55, 88, 131, 168	0
1	F	310/327 (94%)	0.36	26 (8%) 11 7	46, 100, 149, 182	0
All	All	1863/1962 (94%)	0.03	62 (3%) 46 41	34, 71, 129, 182	0

The worst 5 of 62 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	54	ARG	4.7
1	F	70	ALA	4.6
1	F	326	PRO	4.0
1	E	244	VAL	3.8
1	E	150	LYS	3.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.