



# Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 02:17 am BST

PDB ID : 5X09  
Title : Crystal structure of subunit A mutant P235A/S238C of the A-ATP synthase from pyrococcus horikoshii OT3  
Authors : Dhirendra, S.; Gruber, G.  
Deposited on : 2017-01-20  
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1164 (2.36-2.36)
Clashscore	141614	1232 (2.36-2.36)
Ramachandran outliers	138981	1211 (2.36-2.36)
Sidechain outliers	138945	1212 (2.36-2.36)
RSRZ outliers	127900	1150 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	538	

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4342 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called V-type ATP synthase alpha chain,V-type ATP synthase alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	530	4187	2675	717	778	17	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	79	ARG	GLY	engineered mutation	UNP O57728
A	235	ALA	PRO	engineered mutation	UNP O57728
A	238	CYS	SER	engineered mutation	UNP O57728

- Molecule 2 is ACETIC ACID (three-letter code: ACY) (formula: C<sub>2</sub>H<sub>4</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	4	2	2	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula:  $C_4H_{12}NO_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 4 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula:  $C_6H_{14}O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			8	6	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	131	Total	O	0	0
			131	131		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: V-type ATP synthase alpha chain,V-type ATP synthase alpha chain



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	127.47Å 127.47Å 105.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.35 29.35 – 2.35	Depositor EDS
% Data completeness (in resolution range)	99.6 (30.00-2.35) 99.7 (29.35-2.35)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.08 (at 2.34Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, $R_{free}$	0.235 , 0.272 0.239 , 0.277	Depositor DCC
$R_{free}$ test set	1830 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	56.3	Xtriage
Anisotropy	0.024	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 57.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	4342	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.07% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACY, MPD, TRS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.34	0/4277	0.55	2/5794 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	499	ARG	NE-CZ-NH2	-5.78	117.41	120.30
1	A	373	ARG	NE-CZ-NH2	-5.48	117.56	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4187	0	4237	41	0
2	A	8	0	6	0	0
3	A	8	0	12	0	0
4	A	8	0	14	3	0
5	A	131	0	0	11	0
All	All	4342	0	4269	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.



All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:578:GLN:OE1	5:A:701:HOH:O	1.76	1.03
1:A:210:PRO:O	1:A:224:GLN:O	1.78	1.00
1:A:253:ALA:O	1:A:290:ARG:NH2	2.20	0.75
1:A:142:THR:HG21	1:A:289:GLU:O	1.90	0.72
1:A:249:LYS:HE3	4:A:604:MPD:H11	1.77	0.65
1:A:579:PHE:N	5:A:703:HOH:O	2.30	0.65
1:A:576:ASN:C	5:A:703:HOH:O	2.34	0.65
1:A:72:LEU:HD11	1:A:88:ILE:HG23	1.79	0.63
1:A:114:ARG:CZ	5:A:708:HOH:O	2.47	0.63
1:A:499:ARG:NH2	5:A:705:HOH:O	2.34	0.59
1:A:551:PRO:N	5:A:701:HOH:O	2.36	0.59
1:A:120:PHE:HB2	1:A:189:MET:CE	2.32	0.59
1:A:211:LEU:HD23	1:A:224:GLN:HG2	1.85	0.58
1:A:88:ILE:HG22	1:A:88:ILE:O	2.04	0.57
1:A:382:ARG:NH1	5:A:702:HOH:O	1.97	0.57
1:A:576:ASN:O	5:A:703:HOH:O	2.17	0.56
1:A:468:LYS:NZ	5:A:704:HOH:O	2.32	0.55
1:A:145:ILE:HD12	1:A:290:ARG:HA	1.90	0.52
1:A:74:VAL:HG13	1:A:195:VAL:CG1	2.40	0.52
1:A:77:GLY:HA3	1:A:147:HIS:CE1	2.44	0.52
1:A:210:PRO:O	1:A:211:LEU:HB3	2.10	0.51
1:A:494:VAL:HG11	1:A:531:TYR:HB2	1.93	0.50
1:A:249:LYS:CE	4:A:604:MPD:H11	2.42	0.48
1:A:142:THR:HG23	1:A:289:GLU:HB3	1.94	0.48
1:A:198:LYS:HG2	1:A:370:ARG:HG2	1.94	0.48
1:A:142:THR:CG2	1:A:289:GLU:HB3	2.44	0.48
1:A:266:ASN:HD22	1:A:266:ASN:C	2.16	0.48
1:A:109:ALA:HB1	1:A:110:PRO:HD2	1.96	0.47
1:A:343:ILE:O	1:A:343:ILE:HG23	2.14	0.47
1:A:546:GLU:HG2	1:A:582:LEU:HD22	1.98	0.46
1:A:79:ARG:HG3	1:A:79:ARG:O	2.15	0.46
1:A:142:THR:HG22	1:A:144:ILE:H	1.80	0.45
1:A:551:PRO:CD	5:A:701:HOH:O	2.64	0.45
1:A:130:VAL:HG13	1:A:151:VAL:HG21	1.98	0.44
1:A:200:PRO:HG2	1:A:377:LEU:HD11	2.01	0.43
1:A:259:ILE:O	1:A:294:ILE:O	2.38	0.42
1:A:241:THR:O	1:A:245:HIS:HB2	2.20	0.42
1:A:239:GLY:O	1:A:242:VAL:HG12	2.20	0.42
1:A:472:LEU:HD11	5:A:704:HOH:O	2.20	0.41
1:A:427:PHE:HA	1:A:428:PRO:C	2.42	0.40

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:242:VAL:HG23	4:A:604:MPD:C5	2.52	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	526/538 (98%)	500 (95%)	25 (5%)	1 (0%)	47 56

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	98	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	447/454 (98%)	443 (99%)	4 (1%)	78 87

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	83	SER
1	A	142	THR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	245	HIS
1	A	266	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	244	GLN
1	A	266	ASN
1	A	299	ASN
1	A	448	HIS
1	A	539	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	ACY	A	601	-	1,3,3	1.12	0	0,3,3	0.00	-
3	TRS	A	603	-	7,7,7	0.30	0	9,9,9	0.22	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	ACY	A	602	-	1,3,3	1.47	0	0,3,3	0.00	-
4	MPD	A	604	-	7,7,7	0.26	0	9,10,10	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TRS	A	603	-	-	3/9/9/9	-
4	MPD	A	604	-	-	1/5/5/5	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	603	TRS	C2-C-C1-O1
3	A	603	TRS	N-C-C1-O1
3	A	603	TRS	C3-C-C1-O1
4	A	604	MPD	C2-C3-C4-O4

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	604	MPD	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	530/538 (98%)	1.20	115 (21%) 0 1	39, 65, 159, 219	0

All (115) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	357	PRO	15.4
1	A	397	GLY	13.8
1	A	358	ALA	12.0
1	A	66	VAL	11.9
1	A	102	PHE	11.6
1	A	399	PHE	10.4
1	A	99	THR	10.1
1	A	400	SER	9.8
1	A	96	ARG	9.6
1	A	403	VAL	9.5
1	A	396	GLY	9.1
1	A	94	VAL	8.2
1	A	53	TYR	7.9
1	A	345	GLY	7.8
1	A	406	ASN	7.7
1	A	103	ILE	7.6
1	A	344	SER	7.6
1	A	407	THR	7.5
1	A	356	TYR	7.4
1	A	398	ASP	7.3
1	A	588	ALA	7.3
1	A	408	LEU	7.0
1	A	98	LYS	6.8
1	A	395	PRO	6.3
1	A	401	GLU	6.2
1	A	61	PRO	6.2
1	A	346	ARG	6.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	359	TYR	6.1
1	A	105	ARG	5.9
1	A	67	GLY	5.8
1	A	340	LEU	5.8
1	A	405	GLN	5.6
1	A	77	GLY	5.5
1	A	100	GLY	5.5
1	A	95	ILE	5.5
1	A	62	GLY	5.5
1	A	273	GLU	5.3
1	A	65	VAL	5.2
1	A	59	VAL	5.0
1	A	394	PRO	5.0
1	A	402	PRO	4.9
1	A	404	VAL	4.8
1	A	333	THR	4.7
1	A	70	ALA	4.7
1	A	260	GLY	4.6
1	A	78	PRO	4.6
1	A	337	ALA	4.5
1	A	266	ASN	4.4
1	A	235	ALA	4.2
1	A	259	ILE	4.1
1	A	328	LEU	4.0
1	A	389	ILE	3.9
1	A	393	SER	3.9
1	A	342	GLU	3.9
1	A	294	ILE	3.8
1	A	264	ARG	3.7
1	A	97	GLU	3.7
1	A	343	ILE	3.7
1	A	258	TYR	3.6
1	A	107	VAL	3.5
1	A	409	ARG	3.5
1	A	237	GLY	3.4
1	A	275	PHE	3.4
1	A	79	ARG	3.3
1	A	52	VAL	3.3
1	A	329	MET	3.2
1	A	302	VAL	3.2
1	A	104	ALA	3.2
1	A	360	LEU	3.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	292	VAL	3.1
1	A	236	PHE	3.1
1	A	299	ASN	3.1
1	A	68	THR	3.0
1	A	248	ALA	2.9
1	A	101	ASP	2.9
1	A	584	LYS	2.9
1	A	338	GLU	2.9
1	A	272	LEU	2.9
1	A	270	ASP	2.9
1	A	363	LYS	2.8
1	A	311	THR	2.7
1	A	110	PRO	2.7
1	A	88	ILE	2.7
1	A	51	GLN	2.6
1	A	341	ARG	2.6
1	A	503	LEU	2.6
1	A	327	ALA	2.6
1	A	331	ASP	2.5
1	A	313	ILE	2.5
1	A	355	GLY	2.5
1	A	76	LEU	2.5
1	A	218	ILE	2.5
1	A	336	TRP	2.5
1	A	121	ILE	2.4
1	A	334	SER	2.4
1	A	69	GLY	2.4
1	A	293	LEU	2.3
1	A	92	LEU	2.3
1	A	57	ALA	2.3
1	A	330	ALA	2.3
1	A	64	PRO	2.3
1	A	335	ARG	2.2
1	A	368	TYR	2.2
1	A	85	TYR	2.2
1	A	87	GLY	2.2
1	A	56	THR	2.2
1	A	106	GLY	2.2
1	A	213	THR	2.1
1	A	332	SER	2.1
1	A	581	GLU	2.1
1	A	257	ILE	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	498	LEU	2.1
1	A	295	ALA	2.0
1	A	502	TYR	2.0
1	A	230	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	MPD	A	604	8/8	0.63	0.37	78,84,88,89	0
3	TRS	A	603	8/8	0.75	0.25	90,92,95,95	0
2	ACY	A	602	4/4	0.81	0.30	85,90,92,92	0
2	ACY	A	601	4/4	0.86	0.17	86,87,88,89	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.