



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 28, 2021 – 10:14 AM EST

PDB ID : 6X1U
Title : Structure of pHis Fab (SC39-4) in complex with pHis mimetic peptide
Authors : Kalagiri, R.; Stanfield, R.; Wilson, I.A.; Hunter, T.
Deposited on : 2020-05-19
Resolution : 1.64 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.16
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.16

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.64 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	130704	3122 (1.66-1.62)
Clashscore	141614	3268 (1.66-1.62)
Ramachandran outliers	138981	3215 (1.66-1.62)
Sidechain outliers	138945	3215 (1.66-1.62)
RSRZ outliers	127900	3079 (1.66-1.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	221	<div><div>5%</div><div>90%</div><div>7%</div><div>.</div></div>
1	H	221	<div><div>4%</div><div>88%</div><div>9%</div><div>.</div></div>
2	B	213	<div><div>%</div><div>93%</div><div>7%</div></div>
2	L	213	<div><div>91%</div><div>9%</div></div>
3	C	9	<div><div>33%</div><div>11%</div><div>56%</div></div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	D	9	 33% 11% 56%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	EDO	L	301	-	-	X	-
5	ACT	L	303	-	-	X	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7500 atoms, of which 3 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SC39-4 Heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	H	214	Total	C	N	O	S	0	6	0
			1598	1005	270	311	12			
1	A	214	Total	C	N	O	S	0	5	0
			1593	1002	267	312	12			

- Molecule 2 is a protein called SC39-4 Light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	213	Total	C	N	O	S	0	2	0
			1610	1005	269	327	9			
2	B	213	Total	C	N	O	S	0	6	0
			1631	1017	273	332	9			

- Molecule 3 is a protein called ACLYana-3-pTza peptide.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	4	Total	C	N	O	P	0	0	0
			27	12	7	7	1			
3	C	4	Total	C	N	O	P	0	0	0
			27	12	7	7	1			

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	H	1	Total	C	O	0	0
			4	2	2		
4	L	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	L	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	L	1	Total C H O 7 2 3 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	H	237	Total O 237 237	0	0
6	L	219	Total O 219 219	0	0
6	A	263	Total O 263 263	0	0
6	B	244	Total O 244 244	0	0
6	D	9	Total O 9 9	0	0
6	C	7	Total O 7 7	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: SC39-4 Heavy chain





- Molecule 1: SC39-4 Heavy chain





- Molecule 2: SC39-4 Light chain





- Molecule 2: SC39-4 Light chain



- Molecule 3: ACLYana-3-pTza peptide



- Molecule 3: ACLYana-3-pTza peptide



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	67.06Å 86.11Å 167.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.80 – 1.64 46.80 – 1.64	Depositor EDS
% Data completeness (in resolution range)	98.1 (46.80-1.64) 98.1 (46.80-1.64)	Depositor EDS
R_{merge}	0.88	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.20 (at 1.64Å)	Xtriage
Refinement program	PHENIX (1.16_3549: ???)	Depositor
R, R_{free}	0.189 , 0.225 0.189 , 0.225	Depositor DCC
R_{free} test set	5819 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	20.3	Xtriage
Anisotropy	0.390	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 41.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7500	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 45.66 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.2808e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PCA, UKD, EDO, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.63	0/1640	0.78	0/2248
1	H	0.60	0/1648	0.76	0/2258
2	B	0.59	0/1684	0.72	0/2302
2	L	0.59	0/1651	0.72	0/2257
3	C	0.40	0/11	0.52	0/11
3	D	0.50	0/11	0.51	0/11
All	All	0.60	0/6645	0.75	0/9087

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1593	0	1575	10	0
1	H	1598	0	1587	16	0
2	B	1631	0	1574	9	0
2	L	1610	0	1551	20	0
3	C	27	0	10	0	0
3	D	27	0	10	0	0
4	B	4	0	6	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	H	4	0	6	0	0
4	L	4	0	6	4	0
5	B	12	0	9	1	0
5	L	8	3	6	3	0
6	A	263	0	0	3	0
6	B	244	0	0	3	0
6	C	7	0	0	0	0
6	D	9	0	0	0	0
6	H	237	0	0	5	0
6	L	219	0	0	4	0
All	All	7497	3	6340	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:129:CYS:SG	2:B:119:PRO:HG3	2.34	0.68
2:L:48:ILE:HD13	2:L:54:VAL:HA	1.79	0.65
2:B:189:LYS:HG3	6:B:579:HOH:O	1.96	0.64
2:L:189:LYS:HG3	6:L:576:HOH:O	1.98	0.63
5:B:304:ACT:H3	6:B:479:HOH:O	1.99	0.61
2:B:3:ASP:N	2:B:26:SER:HG	1.98	0.61
2:L:45:LYS:NZ	4:L:301:EDO:H11	2.15	0.61
2:L:3:ASP:N	2:L:26:SER:HG	2.02	0.58
1:H:129:CYS:SG	2:L:119:PRO:HG3	2.45	0.57
1:H:128:PRO:HB3	6:L:540:HOH:O	2.04	0.56
2:L:48:ILE:CD1	2:L:54:VAL:HG12	2.36	0.55
1:A:137:THR:HG22	1:A:187:THR:HG22	1.87	0.55
1:H:111:ILE:N	1:H:111:ILE:HD12	2.23	0.53
2:L:169:ASP:O	2:L:170:CYS:HB2	2.08	0.52
1:H:166[A]:ARG:NH1	2:L:173:ASN:OD1	2.43	0.52
1:A:128:PRO:HB3	6:B:518:HOH:O	2.13	0.48
2:B:190:GLU:OE2	2:B:207[A]:ASN:OD1	2.31	0.48
1:A:31:ARG:NH1	6:A:307:HOH:O	2.45	0.47
1:H:128:PRO:HG2	6:H:415:HOH:O	2.15	0.47
1:H:191:GLN:HA	1:H:192:PRO:C	2.36	0.47
2:B:31:ASP:O	2:B:50:LYS:HA	2.14	0.46
2:L:45:LYS:HZ3	4:L:301:EDO:H11	1.80	0.46
1:H:58:LYS:NZ	5:L:303:ACT:H1	2.30	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:45:LYS:HZ2	4:L:301:EDO:H11	1.79	0.46
1:A:112:SER:HB3	6:A:310:HOH:O	2.16	0.45
1:A:41:PRO:HG2	6:A:317:HOH:O	2.16	0.45
1:H:23[A]:THR:HG23	6:H:522:HOH:O	2.17	0.45
1:H:203:ASN:ND2	6:H:409:HOH:O	2.50	0.44
1:H:117:GLN:NE2	6:H:408:HOH:O	2.50	0.44
2:L:25:ALA:HB1	5:L:302:ACT:H3	1.99	0.44
1:A:172:ARG:NH2	1:A:176:GLY:HA2	2.32	0.44
2:L:117:ILE:HG22	2:L:204:GLN:HG3	1.98	0.44
2:B:61[A]:ARG:HD2	2:B:76:SER:O	2.18	0.44
1:A:51:ILE:HB	1:A:57:LEU:CD2	2.49	0.43
2:L:120:PRO:HD3	2:L:132:ILE:HG12	2.00	0.43
1:H:121:PRO:HB3	1:H:147:TYR:HB3	2.00	0.43
1:H:166[A]:ARG:HD3	2:L:173:ASN:ND2	2.34	0.43
1:H:22:CYS:HB3	1:H:76:VAL:HG22	2.01	0.42
1:H:166[A]:ARG:HH11	2:L:173:ASN:ND2	2.18	0.42
2:L:95:ARG:HG2	2:L:95(A):TRP:CE2	2.55	0.42
2:L:125:VAL:HG12	6:L:403:HOH:O	2.20	0.42
1:A:166:ARG:HH21	2:B:137[A]:ASN:CG	2.22	0.42
2:L:124:GLN:NE2	6:L:407:HOH:O	2.48	0.42
2:B:169:ASP:O	2:B:170:CYS:HB2	2.19	0.41
1:H:166[A]:ARG:HH11	2:L:173:ASN:HD21	1.68	0.41
6:H:610:HOH:O	5:L:303:ACT:H2	2.20	0.41
2:L:45:LYS:HD3	4:L:301:EDO:H12	2.02	0.41
1:H:137:THR:HG22	1:H:187:THR:HG22	2.03	0.40
1:A:166:ARG:NH2	2:B:137[A]:ASN:ND2	2.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	217/221 (98%)	214 (99%)	3 (1%)	0	100	100
1	H	218/221 (99%)	213 (98%)	5 (2%)	0	100	100
2	B	217/213 (102%)	212 (98%)	5 (2%)	0	100	100
2	L	213/213 (100%)	209 (98%)	4 (2%)	0	100	100
3	C	1/9 (11%)	0	1 (100%)	0	100	100
3	D	1/9 (11%)	0	1 (100%)	0	100	100
All	All	867/886 (98%)	848 (98%)	19 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	185/187 (99%)	183 (99%)	2 (1%)	73	55
1	H	186/187 (100%)	183 (98%)	3 (2%)	62	39
2	B	185/179 (103%)	184 (100%)	1 (0%)	88	80
2	L	181/179 (101%)	180 (99%)	1 (1%)	86	75
All	All	737/732 (101%)	730 (99%)	7 (1%)	76	63

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	66	ARG
1	H	133	THR
1	H	215	CYS
2	L	211	CYS
1	A	66	ARG
1	A	215	CYS
2	B	211	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	PCA	H	2	1	7,8,9	1.68	1 (14%)	9,10,12	2.02	4 (44%)
1	PCA	A	2	1	7,8,9	1.81	1 (14%)	9,10,12	1.97	5 (55%)
3	UKD	D	5	3	11,14,15	1.88	3 (27%)	10,20,22	2.42	3 (30%)
3	UKD	C	5	3	11,14,15	1.87	2 (18%)	10,20,22	2.68	2 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PCA	H	2	1	-	0/0/11/13	0/1/1/1
1	PCA	A	2	1	-	0/0/11/13	0/1/1/1
3	UKD	D	5	3	-	0/0/12/14	0/1/1/1
3	UKD	C	5	3	-	0/0/12/14	0/1/1/1

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	2	PCA	CD-N	4.54	1.46	1.34
1	H	2	PCA	CD-N	4.37	1.46	1.34
3	C	5	UKD	NE1-ND1	3.80	1.41	1.34
3	C	5	UKD	ND1-NG	3.67	1.41	1.34
3	D	5	UKD	NE1-ND1	3.62	1.40	1.34
3	D	5	UKD	ND1-NG	3.60	1.41	1.34
3	D	5	UKD	P-O4	2.45	1.54	1.49

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	5	UKD	CE2-NE1-ND1	-7.30	104.06	108.20
3	D	5	UKD	CE2-NE1-ND1	-6.19	104.69	108.20
1	A	2	PCA	CA-N-CD	-3.09	102.99	113.58
1	H	2	PCA	CA-N-CD	-3.00	103.32	113.58
1	H	2	PCA	OE-CD-CG	-2.67	122.10	126.76
1	A	2	PCA	O-C-CA	-2.52	118.17	124.78
1	H	2	PCA	CG-CD-N	2.50	114.86	108.39
1	A	2	PCA	CB-CA-N	2.50	110.47	103.30
3	D	5	UKD	O1-P-O3	2.47	116.19	107.90
1	H	2	PCA	CB-CA-N	2.44	110.30	103.30
3	C	5	UKD	CB-NG-CD2	2.41	135.49	129.82
1	A	2	PCA	OE-CD-CG	-2.37	122.64	126.76
3	D	5	UKD	O3-P-O4	-2.28	105.94	112.45
1	A	2	PCA	CG-CD-N	2.07	113.75	108.39

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	ACT	B	304	-	1,3,3	4.47	1 (100%)	0,3,3	0.00	-

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	ACT	L	302	-	1,3,3	7.55	1 (100%)	0,3,3	0.00	-
4	EDO	B	301	-	3,3,3	0.45	0	2,2,2	0.88	0
5	ACT	B	302	-	1,3,3	8.32	1 (100%)	0,3,3	0.00	-
5	ACT	L	303	-	1,3,3	4.37	1 (100%)	0,3,3	0.00	-
5	ACT	B	303	-	1,3,3	6.58	1 (100%)	0,3,3	0.00	-
4	EDO	H	301	-	3,3,3	0.71	0	2,2,2	0.25	0
4	EDO	L	301	-	3,3,3	0.42	0	2,2,2	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	H	301	-	-	0/1/1/1	-
4	EDO	B	301	-	-	1/1/1/1	-
4	EDO	L	301	-	-	0/1/1/1	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	302	ACT	CH3-C	8.32	1.59	1.48
5	L	302	ACT	CH3-C	7.55	1.58	1.48
5	B	303	ACT	CH3-C	6.58	1.57	1.48
5	B	304	ACT	CH3-C	4.47	1.54	1.48
5	L	303	ACT	CH3-C	4.37	1.54	1.48

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	301	EDO	O1-C1-C2-O2

There are no ring outliers.

4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	304	ACT	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	L	302	ACT	1	0
5	L	303	ACT	2	0
4	L	301	EDO	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	213/221 (96%)	-0.06	10 (4%) 31 29	14, 21, 57, 113	0
1	H	213/221 (96%)	-0.07	8 (3%) 40 38	14, 21, 56, 109	0
2	B	213/213 (100%)	-0.36	2 (0%) 84 85	14, 21, 35, 65	0
2	L	213/213 (100%)	-0.39	1 (0%) 91 91	15, 22, 35, 67	0
3	C	3/9 (33%)	-0.24	0 100 100	30, 30, 31, 35	0
3	D	3/9 (33%)	-0.30	0 100 100	26, 26, 31, 34	0
All	All	858/886 (96%)	-0.22	21 (2%) 59 58	14, 21, 44, 113	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	215	CYS	13.3
1	A	215	CYS	12.4
1	A	132	ASP	8.4
1	H	132	ASP	7.2
1	H	214	THR	6.9
1	A	133	THR	5.6
1	H	133	THR	5.2
2	B	211	CYS	5.0
2	L	211	CYS	4.9
1	H	130	CYS	4.5
1	A	113	SER	4.4
1	A	214	THR	4.2
1	A	130	CYS	3.9
1	H	115	SER	3.9
1	H	113	SER	3.8
1	A	189	SER	3.3
1	A	190	SER	3.3
1	H	114	SER	2.9
1	A	213	SER	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	3	ASP	2.1
1	A	131	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	PCA	A	2	8/9	0.95	0.07	20,23,29,33	0
1	PCA	H	2	8/9	0.96	0.08	25,28,41,55	0
3	UKD	C	5	14/15	0.97	0.08	17,21,25,27	0
3	UKD	D	5	14/15	0.98	0.06	17,19,23,24	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	ACT	B	303	4/4	0.84	0.21	42,44,46,51	0
5	ACT	B	304	4/4	0.86	0.19	27,41,45,48	0
5	ACT	B	302	4/4	0.90	0.23	26,34,40,44	0
5	ACT	L	303	4/4	0.91	0.15	37,41,45,46	0
5	ACT	L	302	4/4	0.91	0.17	24,34,35,41	0
4	EDO	L	301	4/4	0.91	0.10	32,33,36,42	0
4	EDO	H	301	4/4	0.93	0.12	21,24,26,31	0
4	EDO	B	301	4/4	0.96	0.07	33,37,38,41	0

6.5 Other polymers [i](#)

There are no such residues in this entry.