



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 10, 2020 – 10:47 AM BST

PDB ID : 5X3K
Title : Kfla1895 D451A mutant in complex with isomaltose
Authors : Tanaka, Y.; Chen, M.; Tagami, T.; Yao, M.; Kimura, A.
Deposited on : 2017-02-06
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	743	<div> <div>5%</div> <div>89%</div> <div>6%</div> <div>5%</div> </div>
1	B	743	<div> <div>5%</div> <div>86%</div> <div>9%</div> <div>5%</div> </div>
2	C	3	<div> <div>33%</div> <div>67%</div> </div>
3	D	2	<div> <div>100%</div> </div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 11886 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycoside hydrolase family 31.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	708	Total	C	N	O	S	0	0	0
			5590	3556	992	1030	12			
1	B	709	Total	C	N	O	S	0	0	0
			5594	3558	993	1031	12			

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	expression tag	UNP D2PPM7
A	-18	GLY	-	expression tag	UNP D2PPM7
A	-17	SER	-	expression tag	UNP D2PPM7
A	-16	SER	-	expression tag	UNP D2PPM7
A	-15	HIS	-	expression tag	UNP D2PPM7
A	-14	HIS	-	expression tag	UNP D2PPM7
A	-13	HIS	-	expression tag	UNP D2PPM7
A	-12	HIS	-	expression tag	UNP D2PPM7
A	-11	HIS	-	expression tag	UNP D2PPM7
A	-10	HIS	-	expression tag	UNP D2PPM7
A	-9	SER	-	expression tag	UNP D2PPM7
A	-8	SER	-	expression tag	UNP D2PPM7
A	-7	GLY	-	expression tag	UNP D2PPM7
A	-6	LEU	-	expression tag	UNP D2PPM7
A	-5	VAL	-	expression tag	UNP D2PPM7
A	-4	PRO	-	expression tag	UNP D2PPM7
A	-3	ARG	-	expression tag	UNP D2PPM7
A	-2	GLY	-	expression tag	UNP D2PPM7
A	-1	SER	-	expression tag	UNP D2PPM7
A	0	HIS	-	expression tag	UNP D2PPM7
A	451	ALA	ASP	engineered mutation	UNP D2PPM7
B	-19	MET	-	expression tag	UNP D2PPM7
B	-18	GLY	-	expression tag	UNP D2PPM7
B	-17	SER	-	expression tag	UNP D2PPM7
B	-16	SER	-	expression tag	UNP D2PPM7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-15	HIS	-	expression tag	UNP D2PPM7
B	-14	HIS	-	expression tag	UNP D2PPM7
B	-13	HIS	-	expression tag	UNP D2PPM7
B	-12	HIS	-	expression tag	UNP D2PPM7
B	-11	HIS	-	expression tag	UNP D2PPM7
B	-10	HIS	-	expression tag	UNP D2PPM7
B	-9	SER	-	expression tag	UNP D2PPM7
B	-8	SER	-	expression tag	UNP D2PPM7
B	-7	GLY	-	expression tag	UNP D2PPM7
B	-6	LEU	-	expression tag	UNP D2PPM7
B	-5	VAL	-	expression tag	UNP D2PPM7
B	-4	PRO	-	expression tag	UNP D2PPM7
B	-3	ARG	-	expression tag	UNP D2PPM7
B	-2	GLY	-	expression tag	UNP D2PPM7
B	-1	SER	-	expression tag	UNP D2PPM7
B	0	HIS	-	expression tag	UNP D2PPM7
B	451	ALA	ASP	engineered mutation	UNP D2PPM7

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-6)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	C	2	Total	C	O	0	1	0
			35	18	17			

- Molecule 3 is an oligosaccharide called alpha-D-glucopyranose-(1-6)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
3	D	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	350	Total	O	0	0
			350	350		
6	B	226	Total	O	0	0
			226	226		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glycoside hydrolase family 31



- Molecule 1: Glycoside hydrolase family 31



- Molecule 2: alpha-D-glucopyranose-(1-6)-alpha-D-glucopyranose



- Molecule 3: alpha-D-glucopyranose-(1-6)-alpha-D-glucopyranose

Chain D:

100%

GLC1
GLC2

4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	180.58Å 180.58Å 392.29Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.42 – 2.50 48.42 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.42-2.50) 99.9 (48.42-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.96 (at 2.51Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.182 , 0.216 0.182 , 0.217	Depositor DCC
R_{free} test set	1562 reflections (1.84%)	wwPDB-VP
Wilson B-factor (Å ²)	31.9	Xtriage
Anisotropy	0.407	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 39.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	11886	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, GOL, BGC, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	0/5764	0.48	0/7877
1	B	0.33	0/5768	0.47	0/7882
All	All	0.36	0/11532	0.48	0/15759

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	211	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5590	0	5311	30	0
1	B	5594	0	5314	38	0
2	C	35	0	32	1	0
3	D	23	0	21	0	0
4	A	45	0	0	0	0
4	B	5	0	0	0	0
5	A	18	0	24	2	0
6	A	350	0	0	5	1
6	B	226	0	0	2	1
All	All	11886	0	10702	68	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:104:HIS:HD1	1:B:113:SER:HG	1.15	0.93
1:A:41:ASP:OD2	1:A:108:GLY:N	2.10	0.83
1:B:31:VAL:H	1:B:92:THR:HG22	1.45	0.80
1:A:660:GLU:OE1	1:A:711:ARG:NH1	2.15	0.79
1:B:465:ASP:OD2	1:B:467:ARG:NH1	2.17	0.77
1:B:107:ARG:HH22	1:B:112:GLU:HB2	1.56	0.71
1:A:685:ASP:OD2	1:A:699:ARG:NH2	2.28	0.67
6:A:901:HOH:O	2:C:1[A]:GLC:O1	2.14	0.65
1:A:568:PHE:HB3	1:A:663:ILE:HG12	1.81	0.63
1:A:626:ARG:NH1	1:B:583:LEU:O	2.32	0.63
1:A:643:GLU:OE2	6:A:902:HOH:O	2.16	0.60
1:B:314:GLU:HA	1:B:372:TRP:HB2	1.85	0.59
1:A:449:LYS:HA	1:A:498:PHE:HB3	1.85	0.58
1:A:314:GLU:HA	1:A:372:TRP:HB2	1.86	0.57
1:B:449:LYS:HA	1:B:498:PHE:HB3	1.87	0.57
1:B:104:HIS:ND1	1:B:113:SER:OG	2.20	0.56
1:A:626:ARG:HD3	6:A:1206:HOH:O	2.05	0.56
1:B:385:GLN:NE2	1:B:389:ASP:OD1	2.40	0.54
1:B:1:MET:HA	1:B:112:GLU:OE2	2.08	0.54
1:B:92:THR:HG23	1:B:93:PRO:O	2.08	0.52
1:B:525:ARG:HG2	1:B:667:LEU:HD13	1.92	0.51
1:B:47:TRP:O	1:B:50:LEU:HB2	2.11	0.50
1:B:9:ILE:HG23	1:B:11:HIS:H	1.77	0.50
1:B:699:ARG:NH1	6:B:913:HOH:O	2.45	0.49
1:B:287:GLU:OE2	1:B:579:ASN:HB3	2.13	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:578:PHE:CZ	1:A:580:HIS:HA	2.48	0.48
1:A:319:GLU:HA	1:A:322:ILE:CD1	2.43	0.48
1:B:578:PHE:CZ	1:B:580:HIS:HA	2.49	0.48
1:A:482:ARG:HB2	1:A:509:HIS:CE1	2.48	0.48
1:A:585:LEU:O	1:A:592:HIS:NE2	2.38	0.48
1:A:39:GLU:OE1	1:A:39:GLU:N	2.46	0.48
1:B:277:TRP:CD2	1:B:368:LYS:HG3	2.49	0.47
1:A:169:PHE:HA	1:A:204:MET:O	2.15	0.47
1:A:467:ARG:HH12	5:A:812:GOL:C1	2.29	0.46
1:A:138:ARG:NH1	1:A:254:GLU:OE2	2.43	0.46
1:B:560:LEU:HD12	1:B:703:LEU:HD13	1.96	0.46
1:A:532:LEU:HD21	1:A:655:TYR:HB3	1.98	0.46
1:A:245:GLU:HA	1:A:246:PRO:HD3	1.85	0.45
1:A:319:GLU:HA	1:A:322:ILE:HD13	1.98	0.45
1:B:660:GLU:CD	1:B:714:ARG:HG3	2.37	0.45
1:B:319:GLU:HA	1:B:322:ILE:CD1	2.47	0.45
1:B:404:ASP:OD1	1:B:406:THR:OG1	2.24	0.45
1:A:513:TRP:HA	1:A:543:GLY:O	2.16	0.45
1:B:607:ARG:O	1:B:611:THR:HG23	2.17	0.45
1:B:47:TRP:HZ2	1:B:95:LEU:HD13	1.82	0.45
1:B:103:PHE:H	1:B:114:THR:HG22	1.82	0.45
1:B:7:HIS:CD2	1:B:23:VAL:HG11	2.53	0.44
1:B:319:GLU:HA	1:B:322:ILE:HD13	2.00	0.44
1:A:39:GLU:HG2	1:A:39:GLU:O	2.18	0.43
1:B:326:ARG:O	1:B:327:ASP:HB2	2.18	0.43
1:B:191:TYR:CE2	1:B:192:LYS:HE2	2.54	0.42
1:A:282:TRP:HB2	1:A:573:GLN:HB2	1.99	0.42
1:B:5:ARG:O	1:B:8:GLY:N	2.47	0.42
1:B:691:ARG:NH1	6:B:923:HOH:O	2.51	0.42
1:A:440:VAL:O	1:A:444:ASP:HA	2.19	0.42
1:A:467:ARG:HH12	5:A:812:GOL:H12	1.83	0.42
1:B:632:ARG:HA	1:B:633:PRO:HD3	1.93	0.42
1:B:275:PRO:HB2	1:B:277:TRP:CD1	2.55	0.41
1:A:714:ARG:CG	1:A:714:ARG:HH21	2.34	0.41
1:B:450:THR:N	1:B:498:PHE:O	2.53	0.41
1:A:340:ARG:HG3	6:A:1015:HOH:O	2.20	0.41
1:A:100:LYS:HB3	1:A:100:LYS:HE2	1.86	0.41
1:B:228:TRP:HB2	1:B:239:GLU:HB3	2.02	0.41
1:A:334:GLU:HG2	6:A:1241:HOH:O	2.20	0.41
1:A:714:ARG:HH21	1:A:714:ARG:HG2	1.85	0.40
1:B:342:GLU:CD	1:B:342:GLU:H	2.23	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:107:ARG:HG2	1:B:107:ARG:H	1.80	0.40
1:B:9:ILE:HD11	1:B:411:ARG:HB3	2.04	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:1015:HOH:O	6:B:902:HOH:O[6_557]	2.10	0.10

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	704/743 (95%)	696 (99%)	8 (1%)	0	100	100
1	B	705/743 (95%)	695 (99%)	10 (1%)	0	100	100
All	All	1409/1486 (95%)	1391 (99%)	18 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	567/588 (96%)	557 (98%)	10 (2%)	59	81

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	B	567/588 (96%)	557 (98%)	10 (2%)	59 81
All	All	1134/1176 (96%)	1114 (98%)	20 (2%)	59 81

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	35	VAL
1	A	51	GLU
1	A	73	HIS
1	A	183	LEU
1	A	202	LEU
1	A	482	ARG
1	A	513	TRP
1	A	524	PHE
1	A	549	PHE
1	A	714	ARG
1	B	32	LEU
1	B	50	LEU
1	B	80	LYS
1	B	112	GLU
1	B	113	SER
1	B	482	ARG
1	B	513	TRP
1	B	517	GLU
1	B	524	PHE
1	B	549	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	7	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

5 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	BGC	C	1[B]	2	12,12,12	0.40	0	17,17,17	0.63	0
2	GLC	C	1[A]	2	12,12,12	0.42	0	17,17,17	0.60	0
2	GLC	C	2	2	11,11,12	0.22	0	15,15,17	0.68	0
3	GLC	D	1	3	12,12,12	0.41	0	17,17,17	0.71	0
3	GLC	D	2	3	11,11,12	0.23	0	15,15,17	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	C	1[B]	2	-	0/2/22/22	0/1/1/1
2	GLC	C	1[A]	2	-	0/2/22/22	0/1/1/1
2	GLC	C	2	2	-	0/2/19/22	0/1/1/1
3	GLC	D	1	3	-	0/2/22/22	0/1/1/1
3	GLC	D	2	3	-	2/2/19/22	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	D	2	GLC	C4-C5-C6-O6
3	D	2	GLC	O5-C5-C6-O6


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1[A]	GLC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	A	806	-	4,4,4	0.14	0	6,6,6	0.10	0
5	GOL	A	813	-	5,5,5	0.38	0	5,5,5	0.26	0
4	SO4	A	809	-	4,4,4	0.15	0	6,6,6	0.10	0
4	SO4	A	811	-	4,4,4	0.13	0	6,6,6	0.09	0
4	SO4	A	808	-	4,4,4	0.15	0	6,6,6	0.06	0
4	SO4	A	805	-	4,4,4	0.14	0	6,6,6	0.17	0
5	GOL	A	814	-	5,5,5	0.36	0	5,5,5	0.34	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SO4	B	803	-	4,4,4	0.15	0	6,6,6	0.08	0
4	SO4	A	807	-	4,4,4	0.14	0	6,6,6	0.06	0
5	GOL	A	812	-	5,5,5	0.38	0	5,5,5	0.33	0
4	SO4	A	803	-	4,4,4	0.13	0	6,6,6	0.10	0
4	SO4	A	804	-	4,4,4	0.14	0	6,6,6	0.11	0
4	SO4	A	810	-	4,4,4	0.15	0	6,6,6	0.22	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	812	-	-	2/4/4/4	-
5	GOL	A	814	-	-	2/4/4/4	-
5	GOL	A	813	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	814	GOL	O1-C1-C2-O2
5	A	814	GOL	O1-C1-C2-C3
5	A	812	GOL	O1-C1-C2-C3
5	A	812	GOL	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	812	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	708/743 (95%)	-0.23	39 (5%)	25 26	10, 20, 60, 91	0
1	B	709/743 (95%)	-0.15	36 (5%)	28 29	12, 28, 70, 97	0
All	All	1417/1486 (95%)	-0.19	75 (5%)	26 28	10, 24, 67, 97	0

All (75) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	111	ALA	6.5
1	A	106	HIS	5.5
1	A	213	ASP	5.2
1	A	108	GLY	4.9
1	A	107	ARG	4.8
1	A	83	GLY	4.7
1	B	107	ARG	4.7
1	A	88	TRP	4.5
1	B	40	ALA	4.4
1	B	82	LEU	4.2
1	B	105	ALA	4.1
1	B	109	GLY	4.1
1	A	2	ILE	4.0
1	A	110	ALA	3.9
1	A	44	VAL	3.8
1	A	39	GLU	3.8
1	A	42	ARG	3.7
1	A	40	ALA	3.7
1	B	110	ALA	3.7
1	A	57	THR	3.6
1	A	52	LEU	3.6
1	B	37	ALA	3.6
1	B	83	GLY	3.5
1	A	109	GLY	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	39	GLU	3.4
1	B	80	LYS	3.4
1	B	73	HIS	3.3
1	B	108	GLY	3.3
1	B	42	ARG	3.3
1	A	116	TRP	3.2
1	B	85	ASP	3.1
1	B	106	HIS	3.1
1	B	2	ILE	3.0
1	B	48	GLY	3.0
1	A	56	ALA	3.0
1	A	38	PRO	3.0
1	A	85	ASP	3.0
1	B	52	LEU	2.9
1	B	88	TRP	2.9
1	A	86	GLY	2.9
1	B	57	THR	2.8
1	A	105	ALA	2.8
1	A	55	SER	2.8
1	A	76	GLU	2.8
1	A	212	ALA	2.8
1	B	86	GLY	2.8
1	B	32	LEU	2.7
1	A	73	HIS	2.7
1	A	41	ASP	2.7
1	B	54	LEU	2.7
1	B	340	ARG	2.7
1	B	79	ALA	2.7
1	A	112	GLU	2.6
1	B	84	ALA	2.5
1	A	43	VAL	2.5
1	B	116	TRP	2.5
1	A	104	HIS	2.5
1	B	38	PRO	2.4
1	B	342	GLU	2.4
1	B	41	ASP	2.4
1	B	51	GLU	2.4
1	A	1	MET	2.4
1	A	211	GLY	2.3
1	B	43	VAL	2.3
1	A	3	LYS	2.3
1	A	84	ALA	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	49	THR	2.2
1	A	113	SER	2.2
1	A	80	LYS	2.2
1	B	56	ALA	2.1
1	B	87	ALA	2.1
1	A	140	ARG	2.1
1	A	49	THR	2.1
1	A	82	LEU	2.1
1	B	92	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GLC	D	1	12/12	0.97	0.14	19,23,28,35	0
3	GLC	D	2	11/12	0.97	0.12	23,29,30,33	0
2	GLC	C	2	11/12	0.97	0.13	14,19,22,24	0
2	GLC	C	1[A]	12/12	0.98	0.19	22,25,35,37	12
2	BGC	C	1[B]	12/12	0.98	0.19	19,25,34,36	12




The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain D:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	GOL	A	813	6/6	0.90	0.27	34,36,44,51	0
4	SO4	A	807	5/5	0.90	0.22	60,65,76,76	0
4	SO4	B	803	5/5	0.91	0.23	58,63,75,76	0
5	GOL	A	814	6/6	0.91	0.21	23,26,38,38	0
4	SO4	A	805	5/5	0.92	0.23	49,62,65,74	0
4	SO4	A	808	5/5	0.92	0.18	41,45,64,71	0
4	SO4	A	804	5/5	0.92	0.21	49,52,67,77	0
4	SO4	A	803	5/5	0.94	0.18	44,53,66,69	0
4	SO4	A	811	5/5	0.94	0.17	44,45,68,72	0
5	GOL	A	812	6/6	0.95	0.11	30,33,35,47	0
4	SO4	A	806	5/5	0.95	0.22	48,55,60,66	0
4	SO4	A	809	5/5	0.95	0.19	39,47,58,68	0
4	SO4	A	810	5/5	0.96	0.14	29,32,36,41	0

6.5 Other polymers [i](#)

There are no such residues in this entry.