



Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 07:24 pm BST

PDB ID : 4X4R
Title : Crystal structure of the A.fulgidus CCA-adding enzyme in complex with a G70A arginyl-tRNA minihelix ending in CCACC and AMPcPP
Authors : Kuhn, C.-D.; Joshua-Tor, L.
Deposited on : 2014-12-03
Resolution : 3.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1133 (3.20-3.20)
Clashscore	141614	1253 (3.20-3.20)
Ramachandran outliers	138981	1234 (3.20-3.20)
Sidechain outliers	138945	1233 (3.20-3.20)
RSRZ outliers	127900	1095 (3.20-3.20)
RNA backbone	3102	1010 (3.50-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	457	<div> <div></div> <div>93%</div> <div></div> </div>
1	C	457	<div> <div>3%</div> <div>92%</div> <div></div> </div>
2	B	37	<div> <div>24%</div> <div>76%</div> <div>19%</div> <div>5%</div> </div>
2	D	37	<div> <div>11%</div> <div>68%</div> <div>14%</div> <div>19%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	H	2	<div><div></div><div>100%</div><div>50%</div><div>50%</div></div>

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 16960 atoms, of which 8129 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CCA-adding enzyme.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	443	Total	C	H	N	O	S	0	0	0
			7344	2354	3676	639	662	13			
1	C	437	Total	C	H	N	O	S	0	0	0
			7270	2333	3641	632	651	13			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	438	SER	-	expression tag	UNP O28126
A	439	ASN	-	expression tag	UNP O28126
A	440	SER	-	expression tag	UNP O28126
A	441	SER	-	expression tag	UNP O28126
A	442	SER	-	expression tag	UNP O28126
A	443	VAL	-	expression tag	UNP O28126
A	444	ASP	-	expression tag	UNP O28126
A	445	LYS	-	expression tag	UNP O28126
A	446	LEU	-	expression tag	UNP O28126
A	447	ALA	-	expression tag	UNP O28126
A	448	ALA	-	expression tag	UNP O28126
A	449	ALA	-	expression tag	UNP O28126
A	450	LEU	-	expression tag	UNP O28126
A	451	GLU	-	expression tag	UNP O28126
A	452	HIS	-	expression tag	UNP O28126
A	453	HIS	-	expression tag	UNP O28126
A	454	HIS	-	expression tag	UNP O28126
A	455	HIS	-	expression tag	UNP O28126
A	456	HIS	-	expression tag	UNP O28126
A	457	HIS	-	expression tag	UNP O28126
C	438	SER	-	expression tag	UNP O28126
C	439	ASN	-	expression tag	UNP O28126
C	440	SER	-	expression tag	UNP O28126
C	441	SER	-	expression tag	UNP O28126
C	442	SER	-	expression tag	UNP O28126

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	443	VAL	-	expression tag	UNP O28126
C	444	ASP	-	expression tag	UNP O28126
C	445	LYS	-	expression tag	UNP O28126
C	446	LEU	-	expression tag	UNP O28126
C	447	ALA	-	expression tag	UNP O28126
C	448	ALA	-	expression tag	UNP O28126
C	449	ALA	-	expression tag	UNP O28126
C	450	LEU	-	expression tag	UNP O28126
C	451	GLU	-	expression tag	UNP O28126
C	452	HIS	-	expression tag	UNP O28126
C	453	HIS	-	expression tag	UNP O28126
C	454	HIS	-	expression tag	UNP O28126
C	455	HIS	-	expression tag	UNP O28126
C	456	HIS	-	expression tag	UNP O28126
C	457	HIS	-	expression tag	UNP O28126


- Molecule 2 is a RNA chain called G70A tRNA minihelix ending in CCACC.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	35	Total	C	H	N	O	P	0	0	0
			1128	331	382	133	247	35			
2	D	30	Total	C	H	N	O	P	0	0	0
			964	283	327	112	212	30			

- Molecule 3 is a RNA chain called RNA (5'-D(*CP*G)-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	H	2	Total	C	H	N	O	P	0	0	0
			64	19	24	8	12	1			

- Molecule 4 is DIPHOSPHOMETHYLPHOSPHONIC ACID ADENOSYL ESTER (three-letter code: APC) (formula: C₁₁H₁₈N₅O₁₂P₃).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total	C	H	N	O	P	0	0
			45	11	14	5	12	3		
4	C	1	Total	C	H	N	O	P	0	0
			45	11	14	5	12	3		

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Mg	0	0
			1	1		
5	C	1	Total	Mg	0	0
			1	1		

- Molecule 6 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C₆H₁₃NO₄S).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
6	A	1	Total	C	H	N	O	S	0	0
			25	6	13	1	4	1		

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
7	A	1	14	3	8	3	0	0
7	A	1	14	3	8	3	0	0
7	C	1	14	3	8	3	0	0

- Molecule 8 is D(-)-TARTARIC ACID (three-letter code: TAR) (formula: $C_4H_6O_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	C	1	Total	C	H	O	0	0
			14	4	4	6		

- Molecule 9 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	C	1	Total	C	H	O	0	0
			17	4	10	3		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: CCA-adding enzyme

Chain A: 





- Molecule 1: CCA-adding enzyme

Chain C: 




- Molecule 2: G70A tRNA minihelix ending in CCACC

Chain B: 



- Molecule 2: G70A tRNA minihelix ending in CCACC

Chain D: 



- Molecule 3: RNA (5'-D(*CP*G)-3')

Chain H: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	111.10Å 217.37Å 58.56Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.91 – 3.20 29.91 – 3.20	Depositor EDS
% Data completeness (in resolution range)	99.5 (29.91-3.20) 95.5 (29.91-3.20)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.05 (at 3.18Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.2_1309)	Depositor
R, R_{free}	0.209 , 0.266 0.213 , 0.270	Depositor DCC
R_{free} test set	1230 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å ²)	49.0	Xtriage
Anisotropy	0.189	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 37.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	16960	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, TAR, MES, APC, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.25	0/3751	0.40	0/5040
1	C	0.26	0/3712	0.41	0/4987
2	B	0.43	1/830 (0.1%)	0.72	0/1288
2	D	0.45	1/707 (0.1%)	0.69	0/1093
3	H	0.16	0/44	0.52	0/67
All	All	0.29	2/9044 (0.0%)	0.48	0/12475

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	G	OP3-P	-10.58	1.48	1.61
2	D	1	G	OP3-P	-10.42	1.48	1.61

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3668	3676	3668	9	0
1	C	3629	3641	3633	9	1
2	B	746	382	383	1	0
2	D	637	327	329	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	H	40	24	24	0	1
4	A	31	14	14	0	0
4	C	31	14	14	0	0
5	A	1	0	0	0	0
5	C	1	0	0	0	0
6	A	12	13	13	0	0
7	A	12	16	16	0	0
7	C	6	8	8	0	0
8	C	10	4	4	1	0
9	C	7	10	9	0	0
All	All	8831	8129	8115	18	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:358:PHE:O	1:C:363:ARG:NH2	2.28	0.67
1:A:284:ARG:NH1	1:A:325:GLU:OE2	2.28	0.66
2:B:17:A:O2'	2:B:19:U:OP2	2.16	0.64
1:A:363:ARG:NH1	1:A:378:GLU:OE2	2.32	0.63
1:A:274:ARG:NH1	1:A:332:GLU:OE2	2.36	0.58
1:A:240:HIS:ND1	8:C:504:TAR:O1	2.40	0.51
1:A:57:SER:OG	1:A:156:LYS:NZ	2.46	0.48
1:C:224:ARG:NH1	2:D:34:C:OP1	2.47	0.48
1:C:274:ARG:NH2	1:C:433:MET:O	2.48	0.47
1:A:351:ASP:OD2	1:A:354:ASN:ND2	2.48	0.45
1:C:167:VAL:O	1:C:234:ASN:ND2	2.50	0.45
1:C:1:MET:HE2	1:C:1:MET:HA	1.99	0.44
1:C:143:GLY:N	1:C:145:GLU:OE2	2.49	0.43
1:C:252:GLY:O	1:C:257:LYS:NZ	2.44	0.43
1:C:274:ARG:NH1	1:C:332:GLU:OE2	2.51	0.42
1:A:243:ARG:NH2	1:C:350:GLU:O	2.52	0.42
1:A:79:GLY:HA3	1:A:113:PRO:HG3	2.00	0.42
1:A:22:ARG:NH1	1:A:26:GLU:OE1	2.51	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:269:LYS:NZ	3:H:2:G:O2'[1_554]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	441/457 (96%)	423 (96%)	17 (4%)	1 (0%)	47	79
1	C	435/457 (95%)	411 (94%)	23 (5%)	1 (0%)	47	79
All	All	876/914 (96%)	834 (95%)	40 (5%)	2 (0%)	47	79

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	119	GLU
1	C	119	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	393/404 (97%)	392 (100%)	1 (0%)	92	96
1	C	387/404 (96%)	386 (100%)	1 (0%)	92	96
All	All	780/808 (96%)	778 (100%)	2 (0%)	92	96

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	411	TYR
1	C	217	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	B	33/37 (89%)	4 (12%)	0
2	D	27/37 (72%)	3 (11%)	0
3	H	1/2 (50%)	0	0
All	All	61/76 (80%)	7 (11%)	0

All (7) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	B	7	G
2	B	8	G
2	B	10	A
2	B	32	G
2	D	7	G
2	D	8	G
2	D	10	A

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	APC	A	501	5	27,33,33	2.20	7 (25%)	31,52,52	1.60	4 (12%)
4	APC	C	501	5	27,33,33	2.16	7 (25%)	31,52,52	1.54	4 (12%)
6	MES	A	503	-	12,12,12	2.26	1 (8%)	14,16,16	1.71	6 (42%)
8	TAR	C	504	-	3,9,9	0.59	0	6,12,12	0.66	0
7	GOL	A	505	-	5,5,5	0.35	0	5,5,5	0.26	0
9	PEG	C	505	-	6,6,6	0.62	0	5,5,5	0.26	0
7	GOL	A	504	-	5,5,5	0.39	0	5,5,5	0.16	0
7	GOL	C	503	-	5,5,5	0.35	0	5,5,5	0.22	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	APC	A	501	5	-	2/15/38/38	0/3/3/3
4	APC	C	501	5	-	6/15/38/38	0/3/3/3
6	MES	A	503	-	-	0/6/14/14	0/1/1/1
8	TAR	C	504	-	-	4/4/12/12	-
7	GOL	A	505	-	-	2/4/4/4	-
9	PEG	C	505	-	-	0/4/4/4	-
7	GOL	A	504	-	-	2/4/4/4	-
7	GOL	C	503	-	-	2/4/4/4	-

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	501	APC	O4'-C1'	7.67	1.51	1.41
4	C	501	APC	O4'-C1'	7.62	1.51	1.41
6	A	503	MES	C8-S	-7.56	1.66	1.77
4	A	501	APC	C2'-C1'	-3.69	1.48	1.53
4	C	501	APC	C2'-C1'	-3.61	1.48	1.53
4	A	501	APC	C2'-C3'	-2.97	1.45	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	501	APC	C2-N3	2.97	1.36	1.32
4	C	501	APC	C2'-C3'	-2.95	1.45	1.53
4	C	501	APC	C2-N3	2.82	1.36	1.32
4	C	501	APC	O4'-C4'	2.81	1.51	1.45
4	A	501	APC	O4'-C4'	2.70	1.51	1.45
4	C	501	APC	PB-O2B	-2.47	1.50	1.56
4	A	501	APC	PB-O2B	-2.38	1.50	1.56
4	A	501	APC	PA-O2A	-2.35	1.50	1.56
4	C	501	APC	C6-N6	2.03	1.41	1.34

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	501	APC	N3-C2-N1	-5.57	119.98	128.68
4	A	501	APC	N3-C2-N1	-5.32	120.37	128.68
4	A	501	APC	C3'-C2'-C1'	4.40	107.60	100.98
4	C	501	APC	C3'-C2'-C1'	4.21	107.32	100.98
6	A	503	MES	C5-N4-C3	2.65	114.79	108.83
6	A	503	MES	C2-C3-N4	-2.58	106.19	110.10
4	A	501	APC	PG-O3B-PB	-2.57	123.58	132.62
6	A	503	MES	O2S-S-C8	2.41	109.81	106.92
6	A	503	MES	C6-C5-N4	-2.31	106.60	110.10
4	C	501	APC	C4-C5-N7	-2.27	107.03	109.40
4	C	501	APC	O2G-PG-O3B	2.25	112.17	104.64
4	A	501	APC	C4-C5-N7	-2.22	107.08	109.40
6	A	503	MES	O3S-S-C8	2.20	109.32	105.77
6	A	503	MES	O1S-S-C8	2.08	109.42	106.92

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	C	501	APC	PA-C3A-PB-O1B
4	C	501	APC	PA-C3A-PB-O3B
8	C	504	TAR	C1-C2-C3-O3
8	C	504	TAR	O2-C2-C3-O3
8	C	504	TAR	O2-C2-C3-C4
7	C	503	GOL	O1-C1-C2-C3
4	A	501	APC	O4'-C4'-C5'-O5'
4	C	501	APC	O4'-C4'-C5'-O5'
7	A	505	GOL	O1-C1-C2-O2
4	A	501	APC	C3'-C4'-C5'-O5'

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	C	501	APC	C3'-C4'-C5'-O5'
7	A	505	GOL	O1-C1-C2-C3
7	C	503	GOL	O1-C1-C2-O2
8	C	504	TAR	C1-C2-C3-C4
7	A	504	GOL	O1-C1-C2-O2
4	C	501	APC	PB-O3B-PG-O2G
4	C	501	APC	PB-O3B-PG-O3G
7	A	504	GOL	O1-C1-C2-C3


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	C	504	TAR	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	443/457 (96%)	-0.16	2 (0%) 91 86	14, 38, 67, 109	0
1	C	437/457 (95%)	0.04	15 (3%) 45 29	14, 48, 77, 130	0
2	B	35/37 (94%)	1.54	9 (25%) 0 0	19, 95, 169, 180	0
2	D	30/37 (81%)	1.18	4 (13%) 3 2	40, 94, 131, 147	0
3	H	2/2 (100%)	2.70	2 (100%) 0 0	106, 106, 106, 124	0
All	All	947/990 (95%)	0.05	32 (3%) 45 29	14, 43, 93, 180	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	18	G	8.6
2	B	26	G	4.7
2	B	28	G	4.5
2	B	27	C	4.4
2	B	19	U	3.9
2	B	31	C	3.7
2	D	6	C	3.7
1	C	261	GLU	3.5
1	C	139	GLY	3.4
1	C	4	GLU	3.2
2	B	17	A	3.1
1	C	1	MET	3.1
1	C	142	LYS	3.0
3	H	1	C	2.9
2	D	7	G	2.9
1	C	258	HIS	2.8
1	C	121	LYS	2.8
1	C	122	ASN	2.6
2	D	17	A	2.5
1	A	353	ARG	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	H	2	G	2.5
1	C	437	ASP	2.3
2	D	19	U	2.3
2	B	20	C	2.3
1	C	3	VAL	2.2
2	B	6	C	2.2
1	C	39	GLY	2.2
1	C	5	GLU	2.1
1	C	40	VAL	2.1
1	C	2	LYS	2.1
1	C	6	ILE	2.1
1	A	1	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ


There are no carbohydrates in this entry.


6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
8	TAR	C	504	10/10	0.79	0.34	42,60,87,89	0
9	PEG	C	505	7/7	0.84	0.22	48,64,79,79	0
7	GOL	C	503	6/6	0.84	0.34	38,56,109,109	0
6	MES	A	503	12/12	0.86	0.57	65,93,137,144	0
4	APC	C	501	31/31	0.90	0.30	56,80,119,123	0
5	MG	C	502	1/1	0.90	0.44	68,68,68,68	0
7	GOL	A	505	6/6	0.91	0.20	28,36,54,64	0
7	GOL	A	504	6/6	0.92	0.28	27,44,53,53	0
5	MG	A	502	1/1	0.92	0.22	33,33,33,33	0
4	APC	A	501	31/31	0.94	0.22	28,46,81,97	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers ⓘ

There are no such residues in this entry.